

# TEMOZOLOMIDE - temozolomide capsule

## Zybus Pharmaceuticals (USA) Inc.

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TEMOZOLOMIDE CAPSULES safely and effectively. See full prescribing information for TEMOZOLOMIDE CAPSULES.

### TEMOZOLOMIDE Capsules, for oral use

Initial U.S. Approval: 1996

### RECENT MAJOR CHANGES

Warnings and Precautions (5.5) 11/2019

### INDICATIONS AND USAGE

Temozolomide is an alkylating drug indicated for the treatment of adult patients with: (1)

- Newly diagnosed glioblastoma concomitantly with radiotherapy and then as maintenance treatment. (1.1)
- Refractory anaplastic astrocytoma who have experienced disease progression on a drug regimen containing nitrosourea and procarbazine. (1.2)

### DOSAGE AND ADMINISTRATION

- Administer orally.
- **Newly Diagnosed Glioblastoma**
  - 75 mg/m<sup>2</sup> once daily for 42 days concomitant with focal radiotherapy followed by initial maintenance dose of 150 mg/m<sup>2</sup> once daily for Days 1 to 5 of each 28-day cycle for 6 cycles. May increase maintenance dose to 200 mg/m<sup>2</sup> for cycles 2 – 6 based on toxicity. (2.1)
  - Provide Pneumocystis pneumonia (PCP) prophylaxis during concomitant phase and continue in patients who develop lymphopenia until resolution to grade 1 or less. (2.1)
- **Refractory Anaplastic Astrocytoma** : Initial dose of 150 mg/m<sup>2</sup> once daily on Days 1 to 5 of each 28-day cycle. (2.2)

### DOSAGE FORMS AND STRENGTHS

- **Capsules** : 5 mg, 20 mg, 100 mg, 140 mg, 180 mg, and 250 mg (3)

### CONTRAINDICATIONS

- History of hypersensitivity to temozolomide or any other ingredients in temozolomide capsules and dacarbazine. (4.1)

### WARNINGS AND PRECAUTIONS

- **Myelosuppression**

Monitor absolute neutrophil count (ANC) and platelet count prior to each cycle and during treatment. Geriatric patients and women have a higher risk of developing myelosuppression. (5.1) (5)

- **Myelodysplastic Syndrome and Secondary Malignancies** , including myeloid leukemia, have been observed. (5.2)

- **Pneumocystis Pneumonia (PCP)**

Closely monitor all patients, particularly those receiving steroids, for the development of lymphopenia and PCP. (5.3) (5)

- **Hepatotoxicity**

Fatal and severe hepatotoxicity have been reported. Perform liver tests at baseline, midway through the first cycle, prior to each subsequent cycle, and approximately 2 to 4 weeks after the last dose of temozolomide. (5.4) (5)

- **Embryo-Fetal Toxicity**

Can cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception. Advise male patients with pregnant partners or female partners of reproductive potential to use condoms. (5.5, 8.1, 8.3) (5)

-----**ADVERSE REACTIONS**-----

The most common adverse reactions ( $\geq 20\%$  incidence) are: alopecia, fatigue, nausea, vomiting, headache, constipation, anorexia, and convulsions. (6.1) (6)

- The most common Grade 3 to 4 hematologic laboratory abnormalities ( $\geq 10\%$  incidence) in patients with anaplastic astrocytoma are: decreased lymphocytes, decreased platelets, decreased neutrophils, and decreased leukocytes. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Zydus Pharmaceuticals (USA) Inc. at 1-877-993-8779 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch). (6)**

-----**USE IN SPECIFIC POPULATIONS**-----

- **Lactation**

Advise not to breastfeed. (8.2) (7)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 4/2021**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

- 1.1 Newly Diagnosed Glioblastoma
- 1.2 Refractory Anaplastic Astrocytoma

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma
- 2.2 Recommended Dosage and Dosage Modifications for Refractory Anaplastic Astrocytoma
- 2.3 Preparation and Administration

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Myelosuppression
- 5.2 Myelodysplastic Syndrome and Secondary Malignancies
- 5.3 Pneumocystis Pneumonia
- 5.4 Hepatotoxicity
- 5.5 Embryo-Fetal Toxicity

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

**10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

## **14 CLINICAL STUDIES**

14.1 Newly Diagnosed Glioblastoma

14.2 Refractory Anaplastic Astrocytoma

## **15 REFERENCES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Newly Diagnosed Glioblastoma**

Temozolomide capsules are indicated for the treatment of adult patients with newly diagnosed glioblastoma concomitantly with radiotherapy and then as maintenance treatment.

#### **1.2 Refractory Anaplastic Astrocytoma**

Temozolomide capsules are indicated for the treatment of adult patients with refractory anaplastic astrocytoma who have experienced disease progression on a drug regimen containing nitrosourea and procarbazine.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma**

Administer temozolomide capsules orally once daily for 42 consecutive days during the concomitant phase with focal radiotherapy and then once daily on Days 1 to 5 of each 28-day cycle for 6 cycles during the maintenance phase.

Provide *Pneumocystis* pneumonia (PCP) prophylaxis during the concomitant phase and continue in patients who develop lymphocytopenia until resolution to grade 1 or less [see *Warnings and Precautions* (5.3)].

#### **Concomitant Phase**

The recommended dosage of temozolomide capsules is 75 mg/m<sup>2</sup> orally once daily for 42 days (up to 49 days) concomitant with focal radiotherapy (60 Gy administered in 30

fractions). Focal radiotherapy includes the tumor bed or resection site with a 2-to 3-cm margin.

Obtain a complete blood count weekly. No dose reductions are recommended during the concomitant phase. The recommended dosage modifications during the concomitant phase are provided in Table 1.

**Table 1 Temozolomide Dosage Modifications During Concomitant Phase**

<b>Adverse Reaction</b>	<b>Interruption</b>	<b>Discontinuation</b>
Absolute Neutrophil Count	Withhold temozolomide if ANC is greater than or equal to $0.5 \times 10^9/L$ and less than $1.5 \times 10^9/L$ . Resume temozolomide when ANC is greater than or equal to $1.5 \times 10^9/L$ .	Discontinue temozolomide if platelet count is less than $0.5 \times 10^9/L$ .
Platelet Count	Withhold temozolomide if platelet count is greater than or equal to $10 \times 10^9/L$ and less than $100 \times 10^9/L$ . Resume temozolomide when platelet count is greater than or equal to $100 \times 10^9/L$ .	Discontinue temozolomide if platelet count is less than $10 \times 10^9/L$ .
Non-hematological Adverse Reaction (except for alopecia, nausea, vomiting)	Withhold temozolomide if Grade 2 adverse reaction occurs. Resume temozolomide when resolution to Grade 1 or less.	Discontinue temozolomide if Grade 3 or 4 adverse reaction occurs.

### **Maintenance Phase**

Beginning 4 weeks after concomitant phase completion, administer temozolomide orally once daily on Days 1 to 5 of each 28-day cycle for 6 cycles. The recommended dosage of temozolomide is as follows:

- Cycle 1: 150 mg/m<sup>2</sup>per day
- Cycles 2 to 6: May increase to 200 mg/m<sup>2</sup>per day if the following conditions are met before starting cycle 2. If the dose was not escalated at the onset of Cycle 2, **do not** increase the dose for Cycles 3 to 6.
  - Nonhematologic toxicity is grade 2 or less (except for alopecia, nausea, and vomiting)
  - ANC is greater than or equal to  $1.5 \times 10^9/L$  and
  - Platelet count is greater than or equal to  $100 \times 10^9/L$ .

Obtain a complete blood count on Day 22 and then weekly until the ANC is above  $1.5 \times 10^9/L$  and the platelet count is above  $100 \times 10^9/L$ . Do not start the next cycle until the ANC and platelet count exceed these levels.

The recommended dosage modifications during the maintenance phase are provided in Table 2. If temozolomide is withheld, reduce the dose for the next cycle by 50 mg/m<sup>2</sup>per day. Permanently discontinue temozolomide in patients who are unable to tolerate a dose of 100 mg/m<sup>2</sup>per day.

**Table 2 Temozolomide Dosage Modifications During Maintenance Treatment**

<b>Toxicity</b>	<b>Interruption and Dose Reduction</b>	<b>Discontinuation</b>
Absolute Neutrophil Count	Withhold temozolomide if ANC less than $1 \times 10^9/L$ . When ANC is above $1.5 \times 10^9/L$ , resume temozolomide at reduced dose for the next cycle.	Unable to tolerate a dose of $100 \text{ mg}/\text{m}^2$ per day.
Platelet Count	Withhold temozolomide if platelet less than $50 \times 10^9/L$ . When platelet count is above $100 \times 10^9/L$ , resume temozolomide at reduced dose for the next cycle.	Unable to tolerate a dose of $100 \text{ mg}/\text{m}^2$ per day.
Nonhematological Adverse Reaction (except for alopecia, nausea, vomiting)	Withhold temozolomide if Grade 3 adverse reaction. When resolved to grade 1 or less, resume temozolomide at reduced dose for the next cycle.	Recurrent Grade 3 after dose reduction. Grade 4 Unable to tolerate a dose of $100 \text{ mg}/\text{m}^2$ per day.

## **2.2 Recommended Dosage and Dosage Modifications for Refractory Anaplastic Astrocytoma**

The recommended initial dosage of temozolomide is  $150 \text{ mg}/\text{m}^2$  once daily on Days 1 to 5 of each 28-day cycle. Increase the temozolomide dose to  $200 \text{ mg}/\text{m}^2$  per day if the following conditions are met at the nadir and on Day 1 of the next cycle:

- ANC is greater than or equal to  $1.5 \times 10^9/L$  and
- Platelet count is greater than or equal to  $100 \times 10^9/L$

Continue temozolomide until disease progression or unacceptable toxicity. In the clinical trial, treatment could be continued for a maximum of 2 years, but the optimum duration of therapy is not known.

Obtain a complete blood count on Day 22 and then weekly until the ANC is above  $1.5 \times 10^9/L$  and the platelet count is above  $100 \times 10^9/L$ . Do not start the next cycle until the ANC and platelet count exceed these levels.

If the ANC is less than  $1 \times 10^9/L$  or the platelet count is less than  $50 \times 10^9/L$  during any cycle, reduce the temozolomide dose for the next cycle by  $50 \text{ mg}/\text{m}^2$  per day. Permanently discontinue temozolomide in patients who are unable to tolerate a dose of  $100 \text{ mg}/\text{m}^2$  per day.

## **2.3 Preparation and Administration**

Temozolomide is a cytotoxic drug. Follow applicable special handling and disposal procedures.<sup>1</sup>

### ***Temozolomide capsules***

Administer temozolomide consistently with respect to food (fasting vs. nonfasting) [see

*Clinical Pharmacology (12.3)].* To reduce nausea and vomiting, take temozolomide on an empty stomach or at bedtime and consider antiemetic therapy prior to and/or following temozolomide administration.

Swallow temozolomide capsules whole. Do not open or chew capsules.

If capsules are accidentally opened or damaged, take precautions to avoid inhalation or contact with the skin or mucous membranes. In case of powder contact, the hands should be washed.

### **3 DOSAGE FORMS AND STRENGTHS**

Temozolomide capsules, for oral administration

- Temozolomide capsules, 5 mg have white to off-white to light-pink/light-tan granular powder filled in size '5' hard gelatin capsules with green colored cap and white colored body. The capsule body is imprinted with "751" in black.
- Temozolomide capsules, 20 mg have white to off-white to light-pink/light-tan granular powder filled in size '5' hard gelatin capsules with yellow colored cap and white colored body. The capsule body is imprinted with "752" in black.
- Temozolomide capsules, 100 mg have white to off-white to light-pink/light-tan granular powder filled in size '3' hard gelatin capsules with pink colored cap and white colored body. The capsule body is imprinted with "753" in black.
- Temozolomide capsules, 140 mg have white to off-white to light-pink/light-tan granular powder filled in size '2' hard gelatin capsules with blue colored cap and white colored body. The capsule body is imprinted with "754" in black.
- Temozolomide capsules, 180 mg have white to off-white to light-pink/light-tan granular powder filled in size '0' hard gelatin capsules with orange colored cap and white colored body. The capsule body is imprinted with "755" in black.
- Temozolomide capsules, 250 mg have white to off-white to light-pink/light-tan granular powder filled in size '00' hard gelatin capsules with white colored cap and white colored body. The capsule body is imprinted with "756" in black.

### **4 CONTRAINDICATIONS**

Temozolomide is contraindicated in patients with a history of hypersensitivity reactions to:

- temozolomide or any other ingredients in temozolomide capsules; and
- dacarbazine, since both temozolomide and dacarbazine are metabolized to the same active metabolite 5-(3methyltriazene-1-yl)-imidazole-4-carboxamide.

Reactions to temozolomide have included anaphylaxis [*see Adverse Reactions (6.2)*].

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Myelosuppression**

Myelosuppression, including pancytopenia, leukopenia and anemia, some with fatal outcomes, have occurred with temozolomide [*see Adverse Reactions (6.1, 6.2)*].

Geriatric patients and women have been shown in clinical trials to have a higher risk of

developing myelosuppression.

Prior to dosing, patients must have an ANC of  $1.5 \times 10^9/L$  or greater and a platelet count of  $100 \times 10^9/L$  or greater.

For the concomitant phase with radiotherapy, obtain a complete blood count prior to initiation of treatment and weekly during treatment [see *Dosage and Administration (2.1)*].

For the 28-day treatment cycles, obtain a complete blood count prior to treatment on Day 1 and on Day 22 of each cycle. Perform complete blood counts weekly until recovery if the ANC falls below  $1.5 \times 10^9/L$  and the platelet count falls below  $100 \times 10^9/L$  [see *Dosage and Administration (2.1, 2.2)*].

## **5.2 Myelodysplastic Syndrome and Secondary Malignancies**

Cases of myelodysplastic syndrome and secondary malignancies, including myeloid leukemia, have been observed following temozolomide administration.

## **5.3 Pneumocystis Pneumonia**

Pneumocystis pneumonia (PCP) can occur in patients receiving temozolomide. The risk of PCP is increased in patients receiving steroids or with longer treatment regimens.

For patients with newly diagnosed glioblastoma, provide PCP prophylaxis for all patients during the concomitant phase. Continue in patients who experience lymphopenia until resolution to grade 1 or less [see *Dosage and Administration (2.1)*].

Monitor all patients receiving temozolomide for the development of lymphopenia and PCP.

## **5.4 Hepatotoxicity**

Fatal and severe hepatotoxicity have been reported in patients receiving temozolomide. Perform liver tests at baseline, midway through the first cycle, prior to each subsequent cycle, and approximately two to four weeks after the last dose of temozolomide.

## **5.5 Embryo-Fetal Toxicity**

Based on findings from animal studies and its mechanism of action, temozolomide can cause fetal harm when administered to a pregnant woman. Adverse developmental outcomes have been reported in both pregnant patients and pregnant partners of male patients. Oral administration of temozolomide to rats and rabbits during the period of organogenesis resulted in embryoletality and polymalformations at doses less than the maximum human dose based on body surface area.

Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with temozolomide and for at least 6 months after the final dose. Because of potential risk of genotoxic effects on sperm, advise male patients with female partners of reproductive potential to use condoms during treatment with temozolomide and for at least 3 months after the final dose. Advise male patients not to donate semen during treatment with temozolomide and for at least 3 months after the final dose [see *Use in Specific Populations (8.1, 8.3)*].

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Myelosuppression [see *Warnings and Precautions (5.1)*].
- Myelodysplastic Syndrome and Secondary Malignancies [see *Warnings and Precautions (5.2)*].
- *Pneumocystis* Pneumonia [see *Warnings and Precautions (5.3)*].
- Hepatotoxicity [see *Warnings and Precautions (5.4)*].

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### ***Newly Diagnosed Glioblastoma***

The safety of temozolomide was evaluated in Study MK-7365-051 [see *Clinical Studies (14.1)*].

Forty-nine percent (49%) of patients treated with temozolomide reported one or more severe or life-threatening reactions, most commonly fatigue (13%), convulsions (6%), headache (5%), and thrombocytopenia (5%).

The most common adverse reactions ( $\geq 20\%$ ) across the cumulative temozolomide experience were alopecia, fatigue, nausea, and vomiting. Table 3 summarizes the adverse reactions in Newly Diagnosed Glioblastoma Trial. Overall, the pattern of reactions during the maintenance phase was consistent with the known safety profile of temozolomide.

**Table 3 Adverse Reactions ( $\geq 5\%$ ) in Patients Receiving Temozolomide in Newly Diagnosed Glioblastoma Trial**

Adverse Reactions	Concomitant Phase				Maintenance Phase	
	Radiation Therapy and Temozolomide		Radiation Therapy Alone		Temozolomide	
	N=288*		N=285		N=224	
	All Grades	Grade $\geq 3$	All Grades	Grade $\geq 3$	All Grades	Grade $\geq 3$
	(%)	(%)	(%)	(%)	(%)	(%)
<b>Skin and Subcutaneous Tissue</b>						
Alopecia	69		63		55	
Rash	19	1	15		13	1
Dry Skin	2		2		5	<1
Pruritus	4		1		5	
Erythema	5		5		1	
<b>General</b>						



Fatigue	54	7	49	5	61	9
Anorexia	19	1	9	<1	27	1
Headache	19	2	17	4	23	4
Weakness	3	2	3	1	7	2
Dizziness	4	1	4		5	
<b>Gastrointestinal System</b>						
Nausea	36	1	16	<1	49	1
Vomiting	20	<1	6	<1	29	2
Constipation	18	1	6		22	
Diarrhea	6		3		10	1
Stomatitis	7		5	<1	9	1
Abdominal Pain	2	<1	1		5	<1
<b>Eye</b>						
Vision Blurred	9	1	9	1	8	
<b>Injury</b>						
Radiation Injury NOS	7		4	<1	2	
<b>Central and Peripheral Nervous System</b>						
Convulsions	6	3	7	3	11	3
Memory Impairment	3	<1	4	<1	7	1
Confusion	4	1	4	2	5	2
<b>Special Senses Other</b>						
Taste Perversion	6		2		5	
<b>Respiratory System</b>						
Coughing	5	1	1		8	<1
Dyspnea	4	2	3	1	5	<1
<b>Psychiatric</b>						
Insomnia	5		3	<1	4	
<b>Immune System</b>						
Allergic Reaction	5		2	<1	3	
<b>Platelet, Bleeding and Clotting</b>						
Thrombocytopenia	4	3	1		8	4
<b>Musculoskeletal System</b>						
Arthralgia	2	<1	1		6	

\*One patient who was randomized to radiation therapy only arm received radiation therapy and temozolomide.

NOS=not otherwise specified.

**Note:** Grade 5 (fatal) adverse reactions are included in the Grade  $\geq 3$  column.

When laboratory abnormalities and adverse reactions were combined, Grade 3 or Grade 4 neutrophil abnormalities including neutropenic reactions were observed in 8% of patients, and Grade 3 or Grade 4 platelet abnormalities including thrombocytopenic reactions, were observed in 14% of patients.

### **Refractory Anaplastic Astrocytoma**

The safety of temozolomide was evaluated in Study MK-7365-006 [see *Clinical Studies (14.2)*].

Myelosuppression (thrombocytopenia and neutropenia) was the dose-limiting adverse reaction. It usually occurred within the first few cycles of therapy and was not cumulative. Myelosuppression occurred late in the treatment cycle and returned to normal, on average, within 14 days of nadir counts. The median nadirs occurred at 26 days for platelets (range: 21-40 days) and 28 days for neutrophils (range: 1-44 days). Only 14% (22/158) of patients had a neutrophil nadir and 20% (32/158) of patients had a platelet nadir, which may have delayed the start of the next cycle. Less than 10% of patients required hospitalization, blood transfusion, or discontinuation of therapy due to myelosuppression.

The most common adverse reactions ( $\geq 20\%$ ) were nausea, vomiting, headache, fatigue, constipation, and convulsions.

Tables 4 and 5 summarize the adverse reactions and hematological laboratory abnormalities in Refractory Anaplastic Astrocytoma Trial.

**Table 4 Adverse Reactions ( $\geq 5\%$ ) in Patients Receiving Temozolomide in Refractory Anaplastic Astrocytoma Trial**

Adverse Reactions	Temozolomide N=158	
	All Reactions (%)	Grades 3-4 (%)
<b>Gastrointestinal System</b>		
Nausea	53	10
Vomiting	42	6
Constipation	33	1
Diarrhea	16	2
Abdominal pain	9	1
Anorexia	9	1
<b>General</b>		
Headache	41	6
Fatigue	34	4
Asthenia	13	6
Fever	13	2
Back pain	8	3
<b>Central and Peripheral Nervous System</b>		
Convulsions	23	5
Hemiparesis	18	6
Dizziness	12	1
Coordination abnormal	11	1
Amnesia	10	4
Insomnia	10	
Paresthesia	9	1
Somnolence	9	3
Paresis	8	3

Urinary incontinence	8	2
Ataxia	8	2
Dysphasia	7	1
Convulsions Local	6	
Gait abnormal	6	1
Confusion	5	
<b>Cardiovascular</b>		
Edema peripheral	11	1
<b>Resistance Mechanism</b>		
Infection Viral	11	
<b>Endocrine</b>		
Adrenal hypercorticism	8	
<b>Respiratory System</b>		
Upper respiratory tract infection	8	
Pharyngitis	8	
Sinusitis	6	
Coughing	5	
<b>Skin and Appendages</b>		
Rash	8	
Pruritus	8	1
<b>Urinary System</b>		
Urinary tract infection	8	
Micturition increased frequency	6	
<b>Psychiatric</b>		
Anxiety	7	1
Depression	6	
<b>Reproductive Disorders</b>		
Breast pain, female	6	
<b>Metabolic</b>		
Weight Increase	5	
<b>Musculoskeletal System</b>		
Myalgia	5	
<b>Vision</b>		
Diplopia	5	
Vision abnormal*	5	

\*This term includes blurred vision; visual deficit; vision changes; and vision troubles.

**Table 5 Grade 3 to 4 Adverse Hematologic Laboratory Abnormalities in Refractory Anaplastic Astrocytoma Trial**

	<b>Temozolomide<sup>*,†</sup></b>
Decreased lymphocytes	55%
Decreased platelets	19%
Decreased neutrophils	14%

Decreased leukocytes	11%
Decreased hemoglobin	4%

\* Change from Grade 0 to 2 at baseline to Grade 3 or 4 during treatment.

† Denominator range= 142, 158

## **Hematological Toxicities for Advanced Gliomas**

In clinical trial experience with 110 to 111 females and 169 to 174 males (depending on measurements), females experienced higher rates of Grade 4 neutropenia ( $ANC < 0.5 \times 10^9/L$ ) and thrombocytopenia ( $< 20 \times 10^9/L$ ) than males in the first cycle of therapy (12% vs. 5% and 9% vs. 3%, respectively).

In the entire safety database for which hematologic data exist (N=932), 7% (4/61) and 9.5% (6/63) of patients > 70 years experienced Grade 4 neutropenia or thrombocytopenia in the first cycle, respectively. For patients  $\leq 70$  years, 7% (62/871) and 5.5% (48/879) experienced Grade 4 neutropenia or thrombocytopenia in the first cycle, respectively. Pancytopenia, leukopenia, and anemia also occurred.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of temozolomide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to the drug exposure.

**Dermatologic:** Toxic epidermal necrolysis and Stevens-Johnson syndrome

**Immune System:** Hypersensitivity reactions, including anaphylaxis. Erythema multiforme, which resolved after discontinuation of temozolomide and, in some cases, recurred upon rechallenge.

**Hematopoietic:** Prolonged pancytopenia, which may result in aplastic anemia and fatal outcomes.

**Hepatobiliary:** Fatal and severe hepatotoxicity, elevation of liver enzymes, hyperbilirubinemia, cholestasis, and hepatitis.

**Infections:** Serious opportunistic infections, including some cases with fatal outcomes, with bacterial, viral (primary and reactivated), fungal, and protozoan organisms.

**Pulmonary:** Interstitial pneumonitis, pneumonitis, alveolitis, and pulmonary fibrosis.

**Endocrine:** Diabetes insipidus

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### **Risk Summary**

Based on its mechanism of action [see *Clinical Pharmacology (12.1)*] and findings from animal studies, temozolomide can cause fetal harm when administered to a pregnant woman. Available postmarketing reports describe cases of spontaneous abortions and congenital malformations, including polymalformations with central nervous system,

facial, cardiac, skeletal, and genitourinary system anomalies with exposure to temozolomide during pregnancy. These cases report similar adverse developmental outcomes to those observed in animal studies. Administration of temozolomide to rats and rabbits during the period of organogenesis caused numerous external, internal, and skeletal malformations at doses less than the maximum human dose based on body surface area (see *Data*). Advise pregnant women of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## **Data**

### *Animal Data*

Five consecutive days of oral administration of temozolomide at doses of 75 mg/m<sup>2</sup> and 150 mg/m<sup>2</sup> (0.38 and 0.75 times the human dose of 200 mg/m<sup>2</sup>) in rats and rabbits, respectively, during the period of organogenesis (Gestation Days 8 to 12) caused numerous malformations of the external and internal organs and skeleton in both species. In rabbits, temozolomide at the 150 mg/m<sup>2</sup> dose (0.75 times the human dose of 200 mg/m<sup>2</sup>) caused embryoletality as indicated by increased resorptions.

## **8.2 Lactation**

There are no data on the presence of temozolomide or its metabolites in human milk, the effects on a breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions, including myelosuppression from temozolomide in the breastfed children, advise women not to breastfeed during treatment with temozolomide and for at least 1 week after the final dose.

## **8.3 Females and Males of Reproductive Potential**

### ***Pregnancy Testing***

Verify pregnancy status in females of reproductive potential prior to initiating temozolomide [see *Use in Specific Populations (8.1)*].

### ***Contraception***

#### *Females*

Temozolomide can cause embryo-fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with temozolomide and for at least 6 months after the last dose.

#### *Males*

Because of the potential for embryofetal toxicity and genotoxic effects on sperm cells, advise male patients with pregnant partners or female partners of reproductive potential to use condoms during treatment with temozolomide and for at least 3 months after the final dose [see *Use in Specific Populations (8.1)*, *Nonclinical Toxicology (13.1)*].

Advise male patients not to donate semen during treatment with temozolomide and for at least 3 months after the final dose.

### ***Infertility***

Temozolomide may impair male fertility [see *Nonclinical Toxicology (13.1)*]. Limited data from male patients show changes in sperm parameters during treatment with temozolomide; however, no information is available on the duration or reversibility of these changes.

#### **8.4 Pediatric Use**

Safety and effectiveness of temozolomide have not been established in pediatric patients. Safety and effectiveness of temozolomide capsules were assessed, but not established, in 2 open-label studies in pediatric patients aged 3 to 18 years. In one study, 29 patients with recurrent brain stem glioma and 34 patients with recurrent high-grade astrocytoma were enrolled. In a second study conducted by the Children's Oncology Group (COG), 122 patients were enrolled, including patients with medulloblastoma/PNET (29), high grade astrocytoma (23), low grade astrocytoma (22), brain stem glioma (16), ependymoma (14), other CNS tumors (9), and non-CNS tumors (9). The adverse reaction profile in pediatric patients was similar to adults.

#### **8.5 Geriatric Use**

In the Newly Diagnosed Glioblastoma trial, Study MK-7365-051, 15% of patients were 65 years and older. This study did not include sufficient numbers of patients aged 65 years and older to determine differences in effectiveness from younger patients. No overall differences in safety were observed between patients  $\geq 65$  years and younger patients.

In the Refractory Anaplastic Astrocytoma trial, Study MK-7365-0006, 4% of patients were 70 years and older. This study did not include sufficient numbers of patients aged 70 years and older to determine differences in effectiveness from younger patients. Patients 70 years and older had a higher incidence of Grade 4 neutropenia (25%) and Grade 4 thrombocytopenia (20%) in the first cycle of therapy than patients less than 70 years of age [see *Warnings and Precautions (5.1)*, *Adverse Reactions (6.1)*].

#### **8.6 Renal Impairment**

No dosage adjustment is recommended for patients with creatinine clearance (CL<sub>cr</sub>) of 36 to 130 mL/min/m<sup>2</sup> [see *Clinical Pharmacology (12.3)*]. The recommended dose of temozolomide has not been established for patients with severe renal impairment (CL<sub>cr</sub> < 36 mL/min/m<sup>2</sup>) or for patients with end-stage renal disease on dialysis.

#### **8.7 Hepatic Impairment**

No dosage adjustment is recommended for patients with mild to moderate hepatic impairment (Child Pugh class A and B) [see *Clinical Pharmacology (12.3)*]. The recommended dose of temozolomide has not been established for patients with severe hepatic impairment (Child-Pugh class C).


### **10 OVERDOSAGE**

Dose-limiting toxicity was myelosuppression and was reported with any dose but is expected to be more severe at higher doses. An overdose of 2000 mg per day for 5 days was taken by one patient and the adverse reactions reported were pancytopenia, pyrexia, multi-organ failure, and death. There are reports of patients who have taken more than 5 days of treatment (up to 64 days), with adverse reactions reported

including myelosuppression, which in some cases was severe and prolonged, and infections and resulted in death. In the event of an overdose, monitor complete blood count and provide supportive measures as necessary.

## 11 DESCRIPTION

Temozolomide capsules for oral administration contain temozolomide USP, an imidazotetrazine derivative. The chemical name of temozolomide is 3, 4-dihydro-3-methyl-4-oxoimidazo [5,1-d]-as-tetrazine-8-carboxamide. The structural formula is:



Temozolomide, USP is a white to light pink/light tan powder with a molecular formula of C<sub>6</sub>H<sub>6</sub>N<sub>6</sub>O<sub>2</sub> and a molecular weight of 194.15. It is sparingly soluble in water, soluble in dimethyl sulfoxide, and practically insoluble in toluene. The molecule is stable at acidic pH (< 5) and labile at pH > 7; hence temozolomide can be administered orally. The prodrug, temozolomide, is rapidly hydrolyzed to the active 5-(3-methyltriazene-1-yl) imidazole-4-carboxamide (MTIC) at neutral and alkaline pH values, with hydrolysis taking place even faster at alkaline pH.

Each temozolomide capsule, for oral use contains either 5 mg, 20 mg, 100 mg, 140 mg, 180 mg, or 250 mg of temozolomide. The inactive ingredients for temozolomide capsules are as follows: anhydrous lactose, colloidal silicon dioxide, gelatin, stearic acid, sodium lauryl sulfate, sodium starch glycolate, tartaric acid, and titanium dioxide. The 5 mg capsule shell contains FD&C Blue 1, FD&C Red 3, and iron oxide yellow. The 20 mg capsule shell contains iron oxide yellow. The 100 mg capsule shell contains FD&C Red 3, and D&C Yellow 10. The 140 mg capsule shell contains FD&C Blue 1, and FD&C Red 3. The 180 mg capsule shell contains iron oxide black, iron oxide red, and iron oxide yellow. The capsule shells are imprinted with black pharmaceutical ink which contains following inactive ingredients: iron oxide black, potassium hydroxide, propylene glycol, shellac.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Temozolomide is not directly active but undergoes rapid nonenzymatic conversion at physiologic pH to the reactive compound 5-(3methyltriazene-1-yl)-imidazole-4-carboxamide (MTIC). The cytotoxicity of MTIC is thought to be primarily due to alkylation of DNA. Alkylation (methylation) occurs mainly at the O<sup>6</sup> and N<sup>7</sup> positions of guanine.

### 12.3 Pharmacokinetics

Following a single oral dose of 150 mg/m<sup>2</sup>, the mean C<sub>max</sub> value for temozolomide was 7.5 mcg/mL and for MTIC was 282 ng/mL.

The mean AUC value for temozolomide was 23.4 mcg·hr/mL and for MTIC was 864 ng·hr/mL. Following a single 90-minute intravenous infusion of 150 mg/m<sup>2</sup>, the mean C<sub>max</sub> value for temozolomide was 7.3 mcg/mL and for MTIC was 276 ng/mL. The mean AUC value for temozolomide was 24.6 mcg·hr/mL and for MTIC was 891 ng·hr/mL.

Temozolomide exhibits linear kinetics over the therapeutic dosing range of 75 mg/m<sup>2</sup>/day to 250 mg/m<sup>2</sup>/day.

### **Absorption**

The median T<sub>max</sub> is 1 hour.

#### *Effect of Food*

The mean C<sub>max</sub> and AUC decreased by 32% and 9%, respectively, and median T<sub>max</sub> increased by 2-fold (from 1 hour to 2.25 hours) when temozolomide capsules were administered after a modified high-fat breakfast (587 calories comprised of 1 fried egg, 2 strips of bacon, 2 slices of toast, 2 pats of butter, and 8 oz whole milk).

### **Distribution**

Temozolomide has a mean apparent volume of distribution of 0.4 L/kg (%CV=13%). The mean percent bound of drug-related total radioactivity is 15%.

### **Elimination**

Clearance of temozolomide is about 5.5 L/hr/m<sup>2</sup> and the mean elimination half-life is 1.8 hours.

### **Metabolism**

Temozolomide is spontaneously hydrolyzed at physiologic pH to the active species, MTIC and to temozolomide acid metabolite. MTIC is further hydrolyzed to 5-amino-imidazole-4-carboxamide (AIC), which is known to be an intermediate in purine and nucleic acid biosynthesis, and to methylhydrazine, which is believed to be the active alkylating species. Cytochrome P450 enzymes play only a minor role in the metabolism of temozolomide and MTIC. Relative to the AUC of temozolomide, the exposure to MTIC and AIC is 2.4% and 23%, respectively.

### **Excretion**

About 38% of the administered temozolomide total radioactive dose is recovered over 7 days: 38% in urine and 0.8% in feces. The majority of the recovery of radioactivity in urine is unchanged temozolomide (6%), AIC (12%), temozolomide acid metabolite (2.3%), and unidentified polar metabolite(s) (17%).

### **Specific Populations**

No clinically meaningful differences in the pharmacokinetics of temozolomide were observed based on age (range: 19 years to 78 years), gender, smoking status (smoker vs. non-smoker), creatinine clearance (CL<sub>cr</sub>) of 36 to 130 mL/min/m<sup>2</sup>, or mild to moderate hepatic impairment (Child Pugh class A and B). The pharmacokinetics of temozolomide has not been studied in patients with CL<sub>cr</sub> < 36 mL/min/m<sup>2</sup>, end-stage renal disease on dialysis, or severe hepatic impairment (Child-Pugh class C).

### **Drug Interaction Studies**

#### *Effect of Other Drugs on Temozolomide Pharmacokinetics*



In a multiple-dose study, administration of temozolomide capsules with ranitidine did not change the  $C_{max}$  or AUC values for temozolomide or MTIC.

A population analysis indicated that administration of valproic acid decreases the clearance of temozolomide by about 5%.

A population analysis did not demonstrate any influence of coadministered dexamethasone, prochlorperazine, phenytoin, carbamazepine, ondansetron, histamine-2-receptor antagonists, or phenobarbital on the clearance of orally administered temozolomide.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Temozolomide is carcinogenic in rats at doses less than the maximum recommended human dose. Temozolomide induced mammary carcinomas in both males and females at doses 0.13 to 0.63 times the maximum human dose (25 mg/m<sup>2</sup> to 125 mg/m<sup>2</sup>) when administered orally on 5 consecutive days every 28 days for 6 cycles. Temozolomide also induced fibrosarcomas of the heart, eye, seminal vesicles, salivary glands, abdominal cavity, uterus, and prostate, carcinomas of the seminal vesicles, schwannomas of the heart, optic nerve, and harderian gland, and adenomas of the skin, lung, pituitary, and thyroid at doses 0.5 times the maximum daily dose. Mammary tumors were also induced following 3 cycles of temozolomide at the maximum recommended daily dose.

Temozolomide is a mutagen and a clastogen. In a reverse bacterial mutagenesis assay (Ames assay), temozolomide increased revertant frequency in the absence and presence of metabolic activation. Temozolomide was clastogenic in human lymphocytes in the presence and absence of metabolic activation.

Temozolomide impairs male fertility. Temozolomide caused syncytial cells/immature sperm formation at doses of 50 mg/m<sup>2</sup> and 125 mg/m<sup>2</sup> (0.25 times and 0.63 times the human dose of 200 mg/m<sup>2</sup>) in rats and dogs, respectively, and testicular atrophy in dogs at 125 mg/m<sup>2</sup>.

### **13.2 Animal Toxicology and/or Pharmacology**

Toxicology studies in rats and dogs identified a low incidence of hemorrhage, degeneration, and necrosis of the retina at temozolomide doses equal to or greater than 125 mg/m<sup>2</sup> (0.63 times the human dose of 200 mg/m<sup>2</sup>). These changes were most commonly seen at doses where mortality was observed.

## **14 CLINICAL STUDIES**

### **14.1 Newly Diagnosed Glioblastoma**

The efficacy of temozolomide was evaluated in Study MK-7365-051, a randomized (1:1), multicenter, open-label trial. Eligible patients were required to have newly diagnosed glioblastoma. Patients were randomized to receive either radiation therapy alone or concomitant temozolomide 75 mg/m<sup>2</sup> once daily starting the first day of radiation therapy and continuing until the last day of radiation therapy for 42 days (with a


therapy and continuing until the last day of radiation therapy for 12 days (with a maximum of 49 days), followed by temozolomide 150 mg/m<sup>2</sup> or 200 mg/m<sup>2</sup> once daily on Days 1 to 5 of each 28-day cycle, starting 4 weeks after the end of radiation therapy and continuing for 6 cycles. In both arms, focal radiation therapy was delivered as 60 Gy/30 fractions and included radiation to the tumor bed or resection site with a 2- to 3-cm margin. PCP prophylaxis was required during the concomitant phase, regardless of lymphocyte count and continued until recovery of lymphocyte count to grade 1 or less. The major efficacy outcome measure was overall survival.

A total of 573 patients were randomized, 287 to temozolomide and radiation therapy and 286 to radiation therapy alone. At the time of disease progression, temozolomide was administered as salvage therapy in 161 patients of the 282 (57%) in the radiation therapy alone arm and 62 patients of the 277 (22%) in the temozolomide and radiation therapy arm.

The addition of concomitant and maintenance temozolomide to radiation therapy for the treatment of patients with newly diagnosed glioblastoma showed a statistically significant improvement in overall survival compared to radiotherapy alone (**Figure 1**). The hazard ratio (HR) for overall survival was 0.63 (95% CI: 0.52, 0.75) with a log-rank  $P < 0.0001$  in favor of the temozolomide arm. The median survival was increased by 2.5 months in the temozolomide arm.

**Figure 1**

### **Kaplan-Meier Curves for Overall Survival (ITT Population) in Newly Diagnosed Glioblastoma Trial**



### **14.2 Refractory Anaplastic Astrocytoma**

The efficacy of temozolomide was evaluated in Study MK-7365-006, a single-arm, multicenter trial. Eligible patients had anaplastic astrocytoma at first relapse and a baseline Karnofsky performance status (KPS) of 70 or greater. Patients had previously received radiation therapy and may also have previously received a nitrosourea with or without other chemotherapy. Fifty-four patients had disease progression on prior therapy with both a nitrosourea and procarbazine and their malignancy was considered

refractory to chemotherapy (refractory anaplastic astrocytoma population). Temozolomide capsules were given on Days 1 to 5 of each 28-day cycle at a starting dose of 150 mg/m<sup>2</sup>/day. If ANC was  $\geq 1.5 \times 10^9$ /L and platelet count was  $\geq 100 \times 10^9$ /L at the nadir and on Day 1 of the next cycle, the temozolomide dose was increased to 200 mg/m<sup>2</sup>/day. The major efficacy outcome measure was progression-free survival at 6 months and the additional efficacy outcome measures were overall survival and overall response rate.

In the refractory anaplastic astrocytoma population (n=54), the median age was 42 years (range: 19 to 76); 65% were male; and 72% had a KPS of >80. Sixty-three percent of patients had surgery other than a biopsy at the time of initial diagnosis. Of those patients undergoing resection, 73% underwent a subtotal resection and 27% underwent a gross total resection. Eighteen percent of patients had surgery at the time of first relapse. The median time from initial diagnosis to first relapse was 13.8 months (range: 4.2 months to 6.3 years).

In the refractory anaplastic astrocytoma population, the overall response rate (CR+PR) was 22% (12 of 54 patients) and the complete response rate was 9% (5 of 54 patients). The median duration of all responses was 50 weeks (range: 16 to 114 weeks) and the median duration of complete responses was 64 weeks (range: 52 to 114 weeks). In this population, progression-free survival at 6 months was 45% (95% CI: 31%, 58%) and progression-free survival at 12 months was 29% (95% CI: 16%, 42%). Median progression-free survival was 4.4 months. Overall survival at 6 months was 74% (95% CI: 62%, 86%) and 12-month overall survival was 65% (95% CI: 52%, 78%). Median overall survival was 15.9 months.

## **15 REFERENCES**

1. "OSHA Hazardous Drugs." *OSHA*.  
<http://www.osha.gov/SLTC/hazardousdrugs/index>.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Temozolomide is a cytotoxic drug. Follow applicable special handling and disposal procedures.<sup>1</sup>

Temozolomide capsules are supplied in amber glass bottles with child-resistant polypropylene caps containing the following capsule strengths:

Temozolomide Capsules, 5 mg have white to off-white to light-pink/light-tan granular powder filled in size '5' hard gelatin capsules with green colored cap and white colored body. The capsule bodies are imprinted with "751" in black and are supplied as follows:

NDC 68382-751-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-751-67 in bottle of 14 capsules with child-resistant closure

Temozolomide Capsules, 20 mg have white to off-white to light-pink/light-tan granular powder filled in size '5' hard gelatin capsules with yellow colored cap and white colored body. The capsule bodies are imprinted with "752" in black and are supplied as follows:

NDC 68382-752-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-752-67 in bottle of 14 capsules with child-resistant closure

Temozolomide Capsules, 100 mg have white to off-white to light-pink/light-tan granular powder filled in size '3' hard gelatin capsules with pink colored cap and white colored body. The capsule bodies are imprinted with "753" in black and are supplied as follows:

NDC 68382-753-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-753-67 in bottle of 14 capsules with child-resistant closure

Temozolomide Capsules, 140 mg have white to off-white to light-pink/light-tan granular powder filled in size '2' hard gelatin capsules with blue colored cap and white colored body. The capsule bodies are imprinted with "754" in black and are supplied as follows:

NDC 68382-754-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-754-67 in bottle of 14 capsules with child-resistant closure

Temozolomide Capsules, 180 mg have white to off-white to light-pink/light-tan granular powder filled in Size '0' hard gelatin capsules with orange colored cap and white colored body. The capsule bodies are imprinted with "755" in black and are supplied as follows:

NDC 68382-755-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-755-67 in bottle of 14 capsules with child-resistant closure

Temozolomide Capsules, 250 mg capsules have white to off-white to light-pink/light-tan granular powder filled in Size '00' hard gelatin capsules with white colored cap and white colored body. The capsule bodies are imprinted with "756" in black and are supplied as follows:

NDC 68382-756-96 in bottle of 5 capsules with child-resistant closure

NDC 68382-756-67 in bottle of 14 capsules with child-resistant closure

Store temozolomide capsules at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information).

### **Myelosuppression**

Inform patients that temozolomide can cause low blood cell counts and the need for frequent monitoring of blood cell counts. Advise patients to contact their healthcare provider immediately for bleeding, fever, or other signs of infection [see *Warnings and Precautions* (5.1)].

### **Myelodysplastic Syndrome and Secondary Malignancies**

Advise patients of the increased risk of myelodysplastic syndrome and secondary malignancies [see *Warnings and Precautions* (5.2)].

### **Pneumocystis Pneumonia**

Advise patients of the increased risk of *Pneumocystis pneumonia* and to contact their

healthcare provided immediately for new or worsening pulmonary symptoms. Inform patients that prophylaxis for *Pneumocystis pneumonia* may be needed [see *Dosage and Administration (2.1)*, *Warnings and Precautions (5.3)*].

### **Hepatotoxicity**

Advise patients of the increased risk of hepatotoxicity and to contact their healthcare provider immediately for signs or symptoms of hepatotoxicity [see *Warnings and Precautions (5.4)*].

### **Administration Instructions**

Advise patient to not open capsules. If capsules are accidentally opened or damaged, advise patients to take rigorous precautions with capsule contents to avoid inhalation or contact with the skin or mucous membranes. In case of powder contact, the hands should be washed [see *Dosage and Administration (2.3)*].

### **Embryo-Fetal Toxicity**

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females to inform their healthcare provider of a known or suspected pregnancy [see *Warnings and Precautions (5.5)*, *Use in Specific Populations (8.1)*].

Advise females of reproductive potential to use effective contraception during treatment with temozolomide and for at least 6 months after the last dose [see *Use in Specific Populations (8.3)*].

Advise male patients with pregnant partners or female partners of reproductive potential to use condoms during treatment with temozolomide and for at least 3 months after the final dose [see *Use in Specific Populations (8.3)*, *Nonclinical Toxicology (13.1)*].

Advise male patients not to donate semen during treatment with temozolomide and for at least 3 months after the final dose [see *Use in Specific Populations (8.3)*, *Nonclinical Toxicology (13.1)*].

### **Lactation**

Advise women not to breastfeed during treatment with temozolomide and for at least 1 week after the final dose [see *Use in Specific Populations (8.2)*].

### **Infertility**

Advise males of reproductive potential that temozolomide may impair fertility [see *Use in Specific Populations (8.3)*, *Nonclinical Toxicology (13.1)*].

### **Manufactured by:**

#### **Cadila Healthcare Ltd.**

Ahmedabad, India.

### **Distributed by:**

#### **Zydus Pharmaceuticals (USA) Inc.**

Pennington, NJ 08534

Rev.: 06/20

**Patient Information**  
**Temozolomide (tem" oh zol' oh mide) Capsules**

**What is the most important information I should know about temozolomide capsules?**

**Temozolomide capsules may cause birth defects .**

Females and female partners of male patients who take temozolomide capsules:

- o Avoid becoming pregnant while taking temozolomide capsules.
- o Females who can become pregnant should use an effective form of birth control (contraception) during treatment and for at least 6 months after your last dose of temozolomide capsules. Your doctor should do a pregnancy test to make sure that you are not pregnant before you start taking temozolomide capsules.
- o Tell your doctor right away if you become pregnant or think you are pregnant during treatment with temozolomide capsules.

Males taking temozolomide capsules and have a female partner who is pregnant or who can become pregnant:

- o Use a condom for birth control (contraception) during treatment and for at least 3 months after taking your final dose of temozolomide capsules.
- o Do not donate semen during treatment and for at least 3 months after your final dose of temozolomide capsules.

**See the section "What are the possible side effects of temozolomide capsules?" for more information about side effects.**

**What are temozolomide capsules?**

Temozolomide capsules are prescription medicine used to treat adults with certain brain cancer tumors.

It is not known if temozolomide capsules are safe and effective in children.

**Who should not take temozolomide capsules?**

**Do not take temozolomide capsules if you:**

- have had an allergic reaction to temozolomide or any of the other ingredients in temozolomide capsules. See the end of this leaflet for a list of ingredients in temozolomide capsules. Symptoms of an allergic reaction with temozolomide capsules may include: a red itchy rash, or a severe allergic reaction, such as trouble breathing, swelling of the face, throat, or tongue, or severe skin reaction. If you are not sure, ask your doctor.
- have had an allergic reaction to dacarbazine (DTIC), another cancer medicine.

**What should I tell my doctor before taking temozolomide capsules?**

**Tell your doctor about all of your medical conditions, including if you:**

- have kidney problems
- have liver problems
- are pregnant or plan to become pregnant. **See "What is the most important information I should know about temozolomide capsules?"**
- are breast-feeding or plan to breastfeed. It is not known if temozolomide passes into your breast milk. **Do not** breastfeed during treatment and for at least 1 week after your last dose of temozolomide capsule.

**Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your doctor if you take a medicine that contains valproic acid. Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

### **How should I take temozolomide capsules?**

Temozolomide may be taken by mouth as a capsule.

Your doctor will decide the best way for you to take temozolomide. Take temozolomide exactly as prescribed by your doctor.

There are 2 common dosing schedules for taking temozolomide capsules depending on the type of brain cancer tumor that you have.

- People with certain brain cancer tumors take or receive temozolomide capsules:

- 1 time each day for 42 days in a row (possibly 49 days depending on side effects) along with receiving radiation treatment. **This is 1 cycle of treatment .**

- After this, your doctor may prescribe 6 more cycles of temozolomide capsules as "maintenance" treatment. For each of these cycles, you take or receive temozolomide capsules 1 time each day for 5 days in a row and then you stop taking it for the next 23 days. **This is a 28-day maintenance treatment cycle.**

- People with certain other brain cancer tumors take or receive temozolomide capsules:

- 1 time each day for 5 days in a row only, and then stop taking it for the next 23 days. **This is 1 cycle of treatment (28 days).**

- Your doctor will watch your progress on temozolomide capsules and decide how long you should take it. You might take temozolomide capsules until your tumor gets worse or for possibly up to 2 years.

- If your doctor prescribes a treatment regimen that is different from the information in this leaflet, make sure you follow the instructions given to you by your doctor.

- Your doctor may change your dose of temozolomide capsules, or tell you to stop temozolomide capsules for a short period of time or permanently if you have certain side effects.

- Your doctor will decide how many treatment cycles of temozolomide capsules that you will receive, depending on how you respond to and tolerate treatment.

### **Temozolomide capsules**

- Take temozolomide capsules exactly as your doctor tells you to.

- Temozolomide capsules contain a white capsule body with a color cap and the colors vary based on the dosage strength. Your doctor may prescribe more than 1 strength of temozolomide capsules for you, so it is important that you understand how to take your medicine the right way. Be sure that you understand exactly how many capsules you need to take on each day of your treatment, and what strengths to take. **This may be different whenever you start a new cycle .**

- **Do not** take more temozolomide capsules than prescribed.

- Talk to your doctor or pharmacist before taking your dose if you are not sure how much temozolomide capsules to take. This will help to prevent taking too much temozolomide capsules and decrease your chances of getting

serious side effects.

- Take each day's dose of temozolomide capsules at one time, with a full glass of water.
- **Swallow temozolomide capsules whole.** Do not chew, open, or split the capsules.
- Take temozolomide capsules at the same time each day.
- Take temozolomide capsules the same way each time, either with food or without food.
- If temozolomide capsules are accidentally opened or damaged, be careful not to breathe in (inhale) the powder from the capsules or get the powder on your skin or mucous membranes (for example, in your nose or mouth). If contact with any of these areas happens, flush the area with water.
- To help reduce nausea and vomiting, try to take temozolomide capsules on an empty stomach or at bedtime. Your doctor may prescribe medicine to help prevent or treat nausea, or other medicines to reduce side effects with temozolomide capsules.
- See your doctor regularly to check your progress. Your doctor will check you for side effects.
- If you take more temozolomide capsules than prescribed, call your doctor or get emergency medical help right away.

### **What are the possible side effects of temozolomide capsules?**

**Temozolomide capsules can cause serious side effects, including:**

- **See "What is the most important information I should know about temozolomide capsules?"**
- **Decreased blood cell counts** . Temozolomide capsules can affect your bone marrow and cause you to have decreased blood cell counts. Decreased white blood cell count, red blood cell count and platelet count are common with temozolomide capsules but it can also be severe and lead to death.
  - o Your doctor will do blood tests regularly to check your blood cell counts before you start and during treatment with temozolomide capsules.
  - o Your doctor may need to change the dose of temozolomide capsules, or when you get it depending on your blood cell counts.
  - o **People who are age 70 or older and women have a higher risk for developing decreased blood cell counts during treatment with temozolomide capsules.**
- **Secondary cancers** . Blood problems such as myelodysplastic syndrome (MDS) and new cancers (secondary cancers), including a certain kind of leukemia, can happen in people who take temozolomide capsules. Your doctor will monitor you for this.
- **Pneumocystis pneumonia (PCP)** . PCP is an infection that people can get when their immune system is weak. Temozolomide capsules decreases white blood cells, which makes your immune system weaker and can increase your risk of getting PCP.
  - o People who are taking steroid medicines or who stay on temozolomide capsules for a longer period of time may have an increased risk of getting PCP infection.
  - o Anyone who takes temozolomide capsules will be watched carefully by their doctor for low blood cell counts and this infection.
  - o Tell your doctor if you have any of the following signs and symptoms of PCP infection: shortness of breath, or fever, chills, dry cough.



● **Liver problems. Liver problems can happen with temozolomide capsules and can sometimes be severe and lead to death .** Your doctor will do blood tests to check your liver function before you start taking temozolomide capsules, during treatment, and about 2 to 4 weeks after your last dose of temozolomide capsules.

Common side effects of temozolomide capsules include:

- hair loss
- feeling tired
- nausea and vomiting.
- headache
- constipation
- loss of appetite
- convulsions
- rash
- diarrhea
- unable to move (paralysis) on one side of the body
- weakness
- fever
- dizziness
- coordination problems
- viral infection
- memory loss
- sleep problems

Temozolomide capsules can affect fertility in males and may affect your ability to father a child. Talk with your doctor if fertility is a concern for you.

Tell your doctor about any side effect that bothers you or that does not go away.

These are not all the possible side effects with temozolomide capsules. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store temozolomide capsules?**

- Store temozolomide capsules at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep temozolomide capsules and all medicines out of the reach of children.**

### **General information about the safe and effective use of temozolomide capsules.**

Medicines are sometimes prescribed for purposes other than those listed in the Patient Information leaflet. Do not use temozolomide capsules for a condition for which it was not prescribed. Do not give temozolomide capsules to other people, even if they have the same symptoms that you have. It may harm them.

This leaflet summarizes the most important information about temozolomide capsules. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about temozolomide capsules that is written for health professionals.

Please address medical inquiries to [MedicalAffairs@zydususa.com](mailto:MedicalAffairs@zydususa.com) or Tel.: 1-877-993-8779.

**What are the ingredients in temozolomide capsules?**

Active ingredient: temozolomide USP.

Inactive ingredients: anhydrous lactose, colloidal silicon dioxide, gelatin, stearic acid, sodium lauryl sulfate, sodium starch glycolate, tartaric acid, and titanium dioxide. The 5 mg capsule shell contains FD&C Blue 1, FD&C Red 3, and iron oxide yellow. The 20 mg capsule shell contains iron oxide yellow. The 100 mg capsule shell contains FD&C Red 3, and D&C Yellow 10. The 140 mg capsule shell contains FD&C Blue 1, and FD&C Red 3. The 180 mg capsule shell contains iron oxide black, iron oxide red, and iron oxide yellow. The capsule shells are imprinted with black pharmaceutical ink which contains following inactive ingredients: iron oxide black, potassium hydroxide, propylene glycol, shellac.

**Manufactured by:  
Cadila Healthcare Ltd.  
Ahmedabad, India.**

**Distributed by:  
Zydus Pharmaceuticals (USA) Inc.  
Pennington, NJ 08534**

Rev.: 06/20

This Patient Information has been approved by the U.S. Food and Drug Administration.

**Temozolomide (tem" oh zol' oh mide) Capsules****PHARMACIST:**

Dispense enclosed Patient Package Insert to each patient.

**PHARMACIST INFORMATION SHEET****IMPORTANT DISPENSING INFORMATION**

**For every patient, dispense temozolomide capsules in a separate vial or in its original package, making sure each container lists the strength per capsule and that patients take the appropriate number of capsules from each package or vial. Please see the dispensing instructions below for more information.**

**What is temozolomide capsules?** [see Full Prescribing Information, Indications and Usage (1)]. Temozolomide capsules is an oral alkylating agent for the treatment of newly diagnosed glioblastoma multiforme and refractory anaplastic astrocytoma.

**How is temozolomide capsules dosed?** [see Full Prescribing Information, Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma (2.1), Recommended Dosage and Dosage Modifications for Refractory Anaplastic Astrocytoma (2.2)].

The physician calculates the daily dose of temozolomide capsules for a given patient based on the patient's body surface area (BSA). Round off the resulting dose to the nearest 5 mg. An example of the dosing may be as follows: the initial daily dose of temozolomide capsules in milligrams is the BSA multiplied by mg/m<sup>2</sup>/day (e.g., a patient with a BSA of 1.84 is 1.84 x 75 mg = 138, or 140 mg/day). Adjust the dose for

subsequent cycles according to nadir neutrophil and platelet counts in the previous cycle and at the time of initiating the next cycle.

**How might the dose of temozolomide capsules be modified for Refractory Anaplastic Astrocytoma?** *[see Full Prescribing Information, Recommended Dosage and Dosage Modifications for Refractory Anaplastic Astrocytoma (2.2)].*

The initial dose is 150 mg/m<sup>2</sup> orally once daily for 5 consecutive days per 28-day treatment cycle. Increase the temozolomide capsules dose to 200 mg/m<sup>2</sup>/day for 5 consecutive days per 28-day treatment cycle if both the nadir and day of dosing (Day 29, Day 1 of next cycle) absolute neutrophil counts (ANC) are greater than or equal to 1.5 x 10<sup>9</sup>/L (1500/μL) and both the nadir and Day 29, Day 1 of next cycle platelet counts are greater than or equal to 100 x 10<sup>9</sup>/L (100,000/μL). During treatment, obtain a complete blood count on Day 22 (21 days after the first dose), and weekly until the ANC is above 1.5 x 10<sup>9</sup>/L (1500/μL) and the platelet count exceeds 100 x 10<sup>9</sup>/L (100,000/μL). Do not start the next cycle of temozolomide capsules until the ANC and platelet count exceed these levels. If the ANC falls to less than 1.0 x 10<sup>9</sup>/L (1000/μL) or the platelet count is less than 50 x 10<sup>9</sup>/L (50,000/μL) during any cycle, reduce the dose for the next cycle by 50 mg/m<sup>2</sup>. Permanently discontinue temozolomide capsules in patients who are unable to tolerate a dose of 100 mg/m<sup>2</sup> per day.

Patients should continue taking temozolomide capsules until their physician determines that their disease has progressed or until unacceptable side effects or toxicities occur. In the clinical trial, treatment could be continued for a maximum of 2 years, but the optimum duration of therapy is not known. Physicians may alter the treatment regimen for a given patient.

**Dosing for Patients with Newly Diagnosed Glioblastoma Multiforme** *[see Full Prescribing Information, Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma (2.1)].*

#### *Concomitant Phase Treatment Schedule*

Administer temozolomide capsules orally at 75 mg/m<sup>2</sup> daily for 42 days concomitant with focal radiotherapy (60 Gy administered in 30 fractions), followed by maintenance temozolomide capsules for 6 cycles. No dose reductions are recommended; however, dose interruptions may occur based on patient tolerance. Continue the temozolomide capsules dose throughout the 42-day concomitant period up to 49 days if all of the following conditions are met: absolute neutrophil count greater than or equal to 1.5 x 10<sup>9</sup>/L, platelet count greater than or equal to 100 x 10<sup>9</sup>/L, and nonhematological adverse reactions less than or equal to Grade 1 (except for alopecia, nausea and vomiting). During treatment, obtain a complete blood count weekly. Interrupt or discontinue temozolomide dosing during the concomitant phase according to the hematological and nonhematological toxicity criteria *[see Table 1 in the Full Prescribing Information, Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma (2.1)]*. Pneumocystis pneumonia (PCP) prophylaxis is required during the concomitant administration of temozolomide capsules and radiotherapy, and should be continued in patients who develop lymphocytopenia until resolution to Grade 1 or less.

#### *Maintenance Phase Treatment Schedule*

Four weeks after completing the temozolomide capsules and radiotherapy phase, administer temozolomide capsules for an additional 6 cycles of maintenance treatment. Dosage in Cycle 1 (maintenance) is 150 mg/m<sup>2</sup> once daily for 5 days followed by 23 days

without treatment. At the start of Cycle 2, escalate the dose to 200 mg/m<sup>2</sup>, if the nonhematologic adverse reactions for Cycle 1 are Grade less than or equal to 2 (except for alopecia, nausea and vomiting), absolute neutrophil count (ANC) is greater than or equal to 1.5 x 10<sup>9</sup>/L, and the platelet count is greater than or equal to 100 x 10<sup>9</sup>/L. If the dose was not escalated at Cycle 2, do not escalate the dose in subsequent cycles. Maintain the dose at 200 mg/m<sup>2</sup> per day for the first 5 days of each subsequent cycle except if toxicity occurs.

During treatment, obtain a complete blood count on Day 22 (21 days after the first dose) and weekly until the ANC is above 1.5 x 10<sup>9</sup>/L (1500/μL) and the platelet count exceeds 100 x 10<sup>9</sup>/L

(100,000/μL). Do not start the next cycle of temozolomide capsules until the ANC and platelet count exceed these levels. Base dose reductions during the next cycle on the lowest blood counts and worst nonhematologic adverse reactions during the previous cycle. Apply dose reductions or discontinuations during the maintenance phase [see *Table 2 in the Full Prescribing Information, Recommended Dosage and Dosage Modifications for Newly Diagnosed Glioblastoma (2.1)*].

**How is temozolomide capsules taken?** [See *Full Prescribing Information, Preparation and Administration, temozolomide capsules (2.3)*].

Advise patients to take each day's dose with a full glass of water, preferably on an empty stomach or at bedtime. Taking the medication on an empty stomach or at bedtime may help ease nausea. If patients are also taking anti-nausea or other medications to relieve the side effects associated with temozolomide capsules, advise them to take these medications prior to and/or following administration of temozolomide capsules. Advise patients that temozolomide capsules should be swallowed whole and **NEVER CHEWED**. Advise patients that they **SHOULD NOT** open or split the capsules. If capsules are accidentally opened or damaged, advise patients to take rigorous precautions with the capsule contents to avoid inhalation or contact with the skin or mucous membranes. In case of powder contact, advise the patients to wash their hands. Advise patients to keep this medication away from children.

**What should the patient avoid during treatment with temozolomide capsules?** [see *Full Prescribing Information, Use in Specific Populations, Pregnancy (8.1), Lactation (8.2), Females and Males of Reproductive Potential (8.3)*].

There are no dietary restrictions for patients taking temozolomide capsules. Temozolomide capsules may affect testicular function and may cause birth defects. Advise male patients to exercise adequate birth control measures. Advise female patients to avoid becoming pregnant while receiving this drug. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment and for at least 6 months after the last dose. Advise males of reproductive potential to use condoms during treatment and for at least 3 months after the last dose. Advise male patients not to donate semen during treatment with temozolomide capsules and for at least 3 months after the final dose. It is not known whether temozolomide is excreted into breast milk. Because of the potential for serious adverse reactions in breastfed children, advise women not to breastfeed while taking temozolomide capsules and for at least 1 week after the last dose.

**What are the side effects of temozolomide capsules?** [see *Full Prescribing*

*Information, Adverse Reactions (6)].*

Alopecia, fatigue, nausea, and vomiting are the most common side effects associated with temozolomide capsules. Noncumulative myelosuppression is the dose-limiting toxicity. Patients should be evaluated periodically by their physician to monitor blood counts.

**Other commonly reported side effects reported by patients taking temozolomide capsules** are headache, constipation, anorexia, and convulsions.

**How is temozolomide capsules supplied?** [*see Full Prescribing Information, How Supplied/Storage and Handling (16)].*

Temozolomide capsules are available in 5 mg, 20 mg, 100 mg, 140 mg, 180 mg, and 250 mg strengths. The capsules contain a white capsule body with a color cap, and the colors vary based on the dosage strength.

<b><u>Temozolomide Capsule Strength</u></b>	<b><u>Color</u></b>
5 mg	Green Cap
20 mg	Yellow Cap
100 mg	Pink Cap
140 mg	Blue Cap
180 mg	Orange Cap
250 mg	White Cap

The 5 mg, 20 mg, 100 mg, 140 mg, 180 mg and 250 mg capsule strengths are available in 5-count and 14-count packages.

TEMODAR is also available for injection in single-dose glass vials containing 100 mg temozolomide.

### **How is temozolomide capsules dispensed?**

Dispense each strength of temozolomide capsules in a separate vial or in its original package (one strength per one container). Follow the instructions below:

Based on the dose prescribed, determine the number of each strength of temozolomide capsules needed for the full 42- or 5-day cycle as prescribed by the physician. For example, in a 5-day cycle, 275 mg/day would be dispensed as five 250 mg capsules, five 20 mg capsules and five 5 mg capsules. Label each container with the appropriate number of capsules to be taken each day. Dispense to the patient, making sure each container lists the strength (mg) per capsule and that he or she understands to take the appropriate number of capsules of temozolomide from each package or vial to equal the total daily dose prescribed by the physician.

### **How can temozolomide capsules be ordered?**

Temozolomide capsules can be ordered from your wholesaler. It is important to understand if temozolomide capsules is being used as part of a 42-day regimen or as part of a 5-day course. Remember to order enough temozolomide capsules for the appropriate cycle.

For example:

- a 5-day course of 360 mg/day would require the following to be ordered: two 5-count

packages of 180 mg capsules.

- a 42-day course of 140 mg/day would require the following to be ordered: three 14-count packages of 140 mg capsules.

<b><u>Temozolomide capsules</u></b>	<b><u>NDC Number</u></b>
Sachets:	68382-751-96
5 mg capsules (5 count)	
5 mg capsules (14 count)	68382-751-67
20 mg capsules (5 count)	68382-752-96
20 mg capsules (14 count)	68382-752-67
100 mg capsules (5 count)	68382-753-96
100 mg capsules (14 count)	68382-753-67
140 mg capsules (5 count)	68382-754-96
140 mg capsules (14 count)	68382-754-67
180 mg capsules (5 count)	68382-755-96
180 mg capsules (14 count)	68382-755-67
250 mg capsules (5 count)	68382-756-96
250 mg capsules (14 count)	68382-756-67

**References:**

"OSHA Hazardous Drugs." OSHA. <http://www.osha.gov/SLTC/hazardousdrugs/index>.

TEMODAR is the registered trademark of Merck Sharp & Dohme Corp.

**Manufactured by:**

**Cadila Healthcare Ltd.**

Ahmedabad, India.

**Distributed by:**

**Zyodus Pharmaceuticals (USA) Inc.**

Pennington, NJ 08534

Rev.: 04/21

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 68382-751-96 in bottle of 5 capsules

Temozolomide Capsules, 5 mg

Rx only

NDC 68382-751-96

**Temozolomide**  
Capsules

**5 mg**

751

For Oral Administration

Caution: Cytotoxic agent-Special Handling

zydus  
pharmaceuticals

5 CAPSULES  
Rx only

Each capsule contains: 5 mg Temozolomide, USP.  
Usual Dosage: See package insert for recommendations regarding the use of varying capsule strengths in establishing a daily regimen. Read accompanying directions carefully.  
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].  
Dispense in tight, light-resistant containers as defined in USP/NF.  
**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.**  
Code No.: GUJDRUG0251932  
Manufactured by: Cadila Healthcare Ltd. Matoda, Ahmedabad, India  
Distributed by: Zydus Pharmaceuticals (USA) Inc. Pennington, NJ 08534  
Rev: 06/17

NDC 68382-752-96 in bottle of 5 capsules

Temozolomide Capsules, 20 mg

Rx only

NDC 68382-752-96

**Temozolomide**  
Capsules

**20 mg**

752

For Oral Administration

Caution: Cytotoxic agent-Special Handling

zydus  
pharmaceuticals

5 CAPSULES  
Rx only

Each capsule contains: 20 mg Temozolomide, USP.  
Usual Dosage: See package insert for recommendations regarding the use of varying capsule strengths in establishing a daily regimen. Read accompanying directions carefully.  
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].  
Dispense in tight, light-resistant containers as defined in USP/NF.  
**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.**  
Code No.: GUJDRUG0251932  
Manufactured by: Cadila Healthcare Ltd. Matoda, Ahmedabad, India  
Distributed by: Zydus Pharmaceuticals (USA) Inc. Pennington, NJ 08534  
Rev: 06/17

NDC 68382-753-96 in bottle of 5 capsules

Temozolomide Capsules, 100 mg

Rx only

NDC 68382-753-96

**Temozolomide**  
Capsules

**100 mg**

753

For Oral Administration

Caution: Cytotoxic agent-Special Handling

zydus  
pharmaceuticals

5 CAPSULES  
Rx only

Each capsule contains: 100 mg Temozolomide, USP.  
Usual Dosage: See package insert for recommendations regarding the use of varying capsule strengths in establishing a daily regimen. Read accompanying directions carefully.  
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].  
Dispense in tight, light-resistant containers as defined in USP/NF.  
**KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.**  
Code No.: GUJDRUG0251932  
Manufactured by: Cadila Healthcare Ltd. Matoda, Ahmedabad, India  
Distributed by: Zydus Pharmaceuticals (USA) Inc. Pennington, NJ 08534  
Rev: 06/17

NDC 68382-754-96 in bottle of 5 capsules

Temozolomide Capsules, 140 mg

Rx only



NDC 68382-755-96 in bottle of 5 capsules

Temozolomide Capsules, 180 mg

Rx only



NDC 68382-756-96 in bottle of 5 capsules

Temozolomide Capsules, 250 mg

Rx only



## TEMOZOLOMIDE

temozolomide capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-751
<b>Route of Administration</b>	ORAL		



## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TEMOZOLOMIDE (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

Color	GREEN (GREEN) , WHITE (WHITE)	Score	no score
Shape	CAPSULE (CAPSULE)	Size	11mm
Flavor		Imprint Code	751
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-751-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-751-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206750	10/26/2017	

## TEMOZOLOMIDE

temozolomide capsule

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-752
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>TEMOZOLOMIDE</b> (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	20 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE B POTATO</b> (UNII: 27NA468985)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TARTARIC ACID</b> (UNII: W4888I119H)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	YELLOW (YELLOW) , WHITE (WHITE)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	752
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68382-752-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-752-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA206750	10/26/2017	

# TEMOZOLOMIDE

temozolomide capsule

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-753
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TEMOZOLOMIDE (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	100 mg

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)	
TARTARIC ACID (UNII: W4888I119H)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
GELATIN (UNII: 2G86QN327L)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

## Product Characteristics

<b>Color</b>	PINK (PINK) , WHITE (WHITE)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	753
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-753-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-753-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206750	10/26/2017	

## TEMOZOLOMIDE

temozolomide capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68382-754
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TEMOZOLOMIDE (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	140 mg

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	BLUE (BLUE) , WHITE (WHITE)	Score	no score
Shape	CAPSULE (CAPSULE)	Size	18mm
Flavor		Imprint Code	754
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:68382-754-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-754-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206750	10/26/2017	

## TEMOZOLOMIDE

temozolomide capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68382-755
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TEMOZOLOMIDE (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	180

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3S5YLH9PMK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

### Product Characteristics

Color	ORANGE (ORANGE) , WHITE (WHITE)	Score	no score
Shape	CAPSULE (CAPSULE)	Size	21mm

<b>Flavor</b>		<b>Imprint Code</b>	755	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-755-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-755-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA206750	10/26/2017		

## TEMOZOLOMIDE

temozolomide capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-756
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TEMOZOLOMIDE (UNII: YF1K15M17Y) (TEMOZOLOMIDE - UNII:YF1K15M17Y)	TEMOZOLOMIDE	250 mg

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE (WHITE) , WHITE (WHITE)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	24mm
<b>Flavor</b>		<b>Imprint Code</b>	756
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-756-96	5 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	
2	NDC:68382-756-67	14 in 1 BOTTLE; Type 0: Not a Combination Product	10/26/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206750	10/26/2017	

**Labeler** - Zydus Pharmaceuticals (USA) Inc. (156861945)

**Registrant** - Zydus Pharmaceuticals (USA) Inc. (156861945)

## Establishment

Name	Address	ID/FEI	Business Operations
Cadila Healthcare Limited		863362789	ANALYSIS(68382-751, 68382-752, 68382-753, 68382-754, 68382-755, 68382-756) , MANUFACTURE(68382-751, 68382-752, 68382-753, 68382-754, 68382-755, 68382-756)

Revised: 4/2021

Zydus Pharmaceuticals (USA) Inc.