ANTIMONIUM IODATUM- antimony triiodide pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Antimonium iodatum 30C

Antimonium iodatum 30C Less than 10 ⁻¹² mg antimony per pellet

(**contains 0.443 mg of the active ingredient per pellet)

Productive cough with thick mucus*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen

If pregnant or breast-feeding ask a health professional before use

Keep out of reach of children

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

1-800-BOIRON-1 (1-800-264-7661), BoironUSA.com Info@boiron.com Distributed by Boiron, Inc. Newtown Square, PA 19073





Drug Facts

Active ingredient***: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).


Uses: See symptoms on front panel.


Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: Do not use if pellet dispenser seal is broken.

Drug Facts (continued) **Inactive ingredients:** lactose, sucrose







Drug Facts

Active ingredient***: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).


Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: Do not use if pellet dispenser seal is broken.

Drug Facts (continued) **Inactive ingredients:** lactose, sucrose **Questions or comments?** BoironUSA.com Info@Boiron.com 1-800-BOIRON-1 (1-800-264-7661) **C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details. Distributed by Boiron Inc. Newtown Square, PA 19073



	llet			
Product Informa	tion			
Product Type	HUMAN OTC I	DRUG Item Coc	de (Source)	NDC:0220-0377
Route of Administra	oration ORAL			
Active Ingredient	Active Moiety			
	Ingredient Name		Basis of Strength	Strength
ANTIMONY TRIIODIDE UNII:069647RPT5)	(UNII: 52ZE5Y9Y82) (ANTI	MONY CATION (3+) -	ANTIMONY TRIIODIDE	30 [hp_C] in 30 [hp_C]
Inactive Ingredie	nts			
	Strength			
SUCROSE (UNII: C151H8 LACTOSE, UNSPECIFIE				
Product Characte	eristics			
Color	white	Score		
Shape 	ROUND		Size	
Flavor Contains		Imprint Code		
Packaging	Package Description		Marketing Start	Marketing End
Packaging # Item Code	Package Desc	ription	Date	Date

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
unapproved homeopathic		03/03/1983			

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment

Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-0377)

Revised: 7/2023

Boiron