

MUCINEX FAST-MAX DM MAX- dextromethorphan hydrobromide and guaifenesin solution
RB Health (US) LLC

Maximum Strength Mucinex® Fast-Max®

DM Max

Drug Facts

<i>Active ingredients (in each 20 mL)</i>	<i>Purposes</i>
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate ¹, xanthan gum

¹ may contain this ingredient

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 180 mL Bottle Label

MAXIMUM STRENGTH

NDC 63824-019-66

MUCINEX®

FAST-MAX®

DMMAX

Dextromethorphan HBr-Cough Suppressant

Guaifenesin-Expectorant

- **Controls Cough**

- **Relieves Chest Congestion**
- **Thins & Loosens Mucus**
- **4 Hour Dosing**

6 FL OZ (180mL)

FOR AGES 12+

120115

3024382

MAXIMUM STRENGTH

NDC 83824-019-86

2952020 3024382

Mucinex
FAST-MAX

DM MAX

Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant

- ✓ Controls Cough
- ✓ Relieves Chest Congestion
- ✓ Thins & Loosens Mucus
- ✓ 4 Hour Dosing

6 FL OZ (180mL)

FOR AGES 12+

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Maximum Strength per 4-Hour Dose
 Tamper evident: Do not use if neckband on bottle cap is broken or missing.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

3 63824 01866 9

From the RB Health Unit
 Pharmacy #10784-0224
 POC# 713 76111

LOT: 3284/122
 EXP: N/A
 MFG: N/A

Drug Facts

Active ingredients (in each 20 mL)
 Dextromethorphan HBr 20 mg, Cough suppressant
 Guafenesin 400 mg, Expectorant

Uses

- Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and mucus and make coughs more productive.
- Temporarily relieves:
 - Cough due to minor throat and bronchial irritation as may occur with the common cold or other irritants.
 - The intensity of coughing.
 - The impulse to cough to help you get to sleep.

Warnings

Do not use

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (antidepressant for depression, psychiatric, or related conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before using this product.

Drug Facts (continued)

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.
- cough that occurs with too much phlegm (mucus).

When using this product do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Do not take more than 6 doses in any 24-hour period.
- Measure only with dosing cup provided.
- Do not use dosing cup with other products.

Drug Facts (continued)

- dose as follows or as directed by a doctor:
 - adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
 - children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20°-25°C (68°-77°F)
- do not refrigerate

Inactive ingredients salicylic acid, sodium bisulfite, FD&C Red No. 40, flavors, glycerin (veg), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbic, saccharin, titanium dioxide, hydroxyethylcellulose, xanthan gum may contain this ingredient.

Questions?
 1-866-MUCINEX (1-866-682-4039)
 You may also report side effects to this phone number.

HEALTH • FITNESS • HOME

Please visit our website
www.mucinex.com
 Patents: www.tb.com/patents

MUCINEX FAST-MAX DM MAX

dextromethorphan hydrobromide and guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-019
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-019-66	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	07/18/2012	
2	NDC:63824-019-22	2 in 1 CARTON	07/01/2020	09/01/2024
2		180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/18/2012	

