

**SINUS CONGESTION AND PAIN- acetaminophen, guaifenesin, phenylephrine  
hcl tablet, film coated  
Kroger Company**

-----  
**Kroger 44-527C-SCP**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
  - headache
  - nasal congestion
  - minor aches and pains
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- liver disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

### ***Questions or comments?***

**1-800-632-6900**

### ***Principal display panel***

**COMPARE TO the active ingredients of TYLENOL® SINUS SEVERE** \*See side panel

NDC 30142-727-08

Kroger®

OUR PHARMACIST RECOMMENDED

**SEVERE DAYTIME**

**Sinus Congestion & Pain**

**Acetaminophen**

Pain Reliever/Fever Reducer

Guaifenesin

Expectorant  
Phenylephrine HCl  
Nasal Decongestant

**NON-DROWSY**

Sinus Headache  
Sinus Pressure  
Nasal Congestion  
Mucus & Chest Congestion

actual  
size

**24**  
CAPLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

\*Tylenol® is a registered trademark  
of Johnson & Johnson Corporation,  
New Brunswick, NJ 08933. Johnson  
& Johnson Corporation is not  
affiliated with The Kroger Co.  
or this product.

50844      REV0922B52708

**DISTRIBUTED BY  
THE KROGER CO.  
CINCINNATI, OHIO 45202**

**QUALITY GUARANTEE  
800-632-6900**

**[www.kroger.com](http://www.kroger.com)**

DISTRIBUTED BY  
**THE KROGER CO.**  
 CINCINNATI, OHIO 45202  
**QUALITY GUARANTEE**  
 800-632-6900  
 www.kroger.com



**Drug Facts (continued)**  
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 Inactive ingredients: corn starch, croscopolone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide  
**Questions or comments?** 1-800-632-6900

**Drug Facts (continued)**  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive  
 ■ If you are not sure whether a drug contains acetaminophen (prescription or nonprescription), ask a doctor or pharmacist.  
 ■ If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ If you have ever had an allergic reaction to this product or any of its ingredients:  
 ■ Ask a doctor before use if you have:  
 ■ heart disease  
 ■ liver disease  
 ■ high blood pressure  
 ■ diabetes  
 ■ thyroid disease  
 ■ difficulty in urination due to enlargement of the prostate gland  
 ■ cough that occurs with too much phlegm (mucus)  
**Warnings (continued)**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ blisters  
 ■ rash  
 ■ skin redness  
 If a skin reaction occurs, stop use and seek medical help right away.  
 Do not use:  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ If you have ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)**  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
 When using this product do not exceed recommended dosage.  
 Stop use and ask a doctor if:  
 ■ nervousness, dizziness, or sleeplessness occur  
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present  
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use. Contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions**  
 ■ do not take more than directed  
 ■ adults and children 12 years and over:  
 ■ take 2 caplets every 4 hours  
 ■ swallow whole - do not crush, chew, or dissolve  
 ■ do not take more than 10 caplets in 24 hours  
 ■ children under 12 years: ask a doctor  
**Other information**  
 ■ each caplet contains: sodium 3 mg  
 ■ see end flap for expiration date and lot number

**Drug Facts (continued)**  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
 When using this product do not exceed recommended dosage.  
 Stop use and ask a doctor if:  
 ■ nervousness, dizziness, or sleeplessness occur  
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present  
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use. Contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions**  
 ■ do not take more than directed  
 ■ adults and children 12 years and over:  
 ■ take 2 caplets every 4 hours  
 ■ swallow whole - do not crush, chew, or dissolve  
 ■ do not take more than 10 caplets in 24 hours  
 ■ children under 12 years: ask a doctor  
**Other information**  
 ■ each caplet contains: sodium 3 mg  
 ■ see end flap for expiration date and lot number

**Drug Facts**  
**Active ingredients (in each caplet)**  
 Acetaminophen 325 mg  
 Pain reliever/fever reducer  
 Guaifenesin 200 mg  
 Expectorant  
 Phenylephrine HCl 5 mg  
 Nasal decongestant  
**Purpose**  
 ■ temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:  
 ■ headache  
 ■ nasal congestion  
 ■ minor aches and pains  
 ■ sinus congestion and pressure  
 ■ temporarily reduces fever  
**Uses**  
 ■ temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:  
 ■ headache  
 ■ nasal congestion  
 ■ minor aches and pains  
 ■ sinus congestion and pressure  
 ■ temporarily reduces fever  
**Drug Facts (continued)**

COMPARE TO the active ingredients in **TYLENOL® SINUS SEVERE** \*See side panel

NDC 30142-727-08

**SEVERE** DayTime

# Sinus Congestion & Pain

**Acetaminophen**  
 Pain Reliever/Fever Reducer  
**Guaifenesin**  
 Expectorant  
**Phenylephrine HCl**  
 Nasal Decongestant

actual size

**NON-DROWSY**  
 Sinus Headache  
 Sinus Pressure  
 Nasal Congestion  
 Mucus & Chest Congestion

**24 CAPLETS**

**Kroger 44-527**

**SINUS CONGESTION AND PAIN**  
 acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

**No Print/No Varnish**  
 Lot and Expiration No.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Recycle logo  
 PLEASE RECYCLE  
 B-1050-527C-08-R  
 REV0922852708

50844 REV0922852708

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-727
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	44;527
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:30142-727-08	2 in 1 CARTON	05/31/2019	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2019	

**Labeler** - Kroger Company (006999528)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(30142-727) , pack(30142-727)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(30142-727)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(30142-727)

Revised: 4/2024

Kroger Company