

**EXCHANGE SELECT CLEARLAX- polyethylene glycol 3350 powder, for solution  
Army & Air Force Exchange Service**

-----  
**Perrigo Company ClearLax® Drug Facts**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

**Purpose**

Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product**

you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - do not combine with starch-based thickeners used for difficulty swallowing
  - ensure that the powder is fully dissolved before drinking
  - do not drink if there are any clumps
  - do not use no more than 7 days
- children 16 years of age or under: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

### **Inactive ingredients**

none

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

exchange select™

Compare to the active ingredient of MiraLAX®

ClearLAX®

Polyethylene Glycol 3350

Powder for Solution, Osmotic Laxative

UNFLAVORED GRIT FREE POWDER

Relieves Occasional Constipation/Irregularity

Softens Stool

30 ONCE-DAILY DOSES

NET WT 17.9 OZ (510 g)

quality value

exchange **select**™

Compare to the  
active ingredient of  
MiraLAX®\*

# ClearLAX®

**Polyethylene Glycol 3350**  
**Powder for Solution, Osmotic Laxative**

UNFLAVORED  
GRIT FREE  
POWDER

Relieves Occasional  
Constipation/Irregularity  
Softens Stool

30  
ONCE-DAILY  
DOSES

NET WT 17.9 OZ (510 g)

: 306D8 CZ F1

quality  
value

Polyethylene Glycol 3350 increases  
frequency of bowel movements and softens the stool.

Dissolves in Any Beverage

## Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line).....	Osmotic Laxative

**Use** ■ relieves occasional constipation (irregularity)  
■ generally produces a bowel movement in 1 to 3 days

## Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have** ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if** ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week

## Drug Facts (continued)

■ adults and children 17 years of age and older: ■ use once a day ■ fill to top of white section in cap which is marked to indicate the correct dose (17 g) ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ do not combine with starch-based thickeners used for difficulty swallowing ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ do not use more than 7 days ■ children 16 years of age or under: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

■ tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**

1-800-719-9260

• Sugar Free



■ you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed unless advised by your doctor ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) ▶

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX®.

**SATISFACTION GUARANTEED  
OR YOUR MONEY BACK**

Packaged For:  
Your Military Exchanges  
By: Perrigo Company  
Allagan, MI USA 49010



306D8 CZ B3

## EXCHANGE SELECT CLEARLAX

polyethylene glycol 3350 powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-306
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-306-03	510 g in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/08/2015	

**Labeler** - Army & Air Force Exchange Service (001695568)

