

**EXCHANGE SELECT CLEARLAX- polyethylene glycol 3350 powder, for solution
Army & Air Force Exchange Service**

Perrigo Company ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient of MiraLax®

ClearLax®

Polyethylene Glycol 3350

Powder for Solution, Osmotic Laxative

UNFLAVORED GRIT FREE POWDER

Relieves Occasional Constipation/Irregularity

Softens Stool

30 ONCE-DAILY DOSES

NET WT 17.9 OZ (510 g)

quality value



Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool. Dissolves in Any Beverage

Drug Facts		Drug Facts (continued)	
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line).....	Purpose Osmotic Laxative	Directions	
Use		<ul style="list-style-type: none"> ■ do not take more than directed unless advised by your doctor ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) ■ adults and children 17 years of age and older: <ul style="list-style-type: none"> ■ fill to top of white section in cap which is marked to indicate the correct dose (17 g) ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor 	
Warnings		Other information	
<p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p>		<ul style="list-style-type: none"> ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if printed foil seal under cap is missing, open or broken <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-719-0260</p> <p>*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX®.</p> <p>SATISFACTION GUARANTEED OR YOUR MONEY BACK</p> <p>Packaged For: Your Military Exchanges By: Perrigo Company Allagan, MI USA 49010</p>	
		<p>• Sugar Free</p>	

EXCHANGE SELECT CLEARLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-306
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-306-03	510 g in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2015	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA090685	10/08/2015		

Labeler - Army & Air Force Exchange Service (001695568)

Revised: 5/2021

Army & Air Force Exchange Service