LOW DOSE ASPIRIN- aspirin tablet, coated H E B

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for safety coated 81 mg aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to his product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stool
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

• store between 15-30°C (59-86°F)

Inactive ingredients

D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch (corn), silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triethyl citrate

Questions or comments?

Call toll free:1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Bayer® Low Dose Aspirin active ingredient†

EASY OPEN CAP

Low Dose

Aspirin 81 mg

Pain reliever (NSAID)

- Enteric Safety Coated
- Aspirin Regimen±

THIS PACKAGE NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN ENTERIC COATED TABLETS

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

±Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

≠Aspirin is not appropriate for everyone, so be sure you talk to your doctor before you begin an aspirin regimen.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

MADE WITH

PRIDE AND CARE FOR H-E-B®

SAN ANTONIO,TX 78204

Product Label



Active ingredient (in each tablet) spirin 81 mg (NSAID **Orug Facts**

for the temporary relief of minor aches and pains or as recommended by Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. eadaches or other symptoms

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

Call toll free: 1-877-753-3935 Monday-Friday 9AM-5PM EST

Questions or comments?

Talk to your doctor or other healthcare provider before using this product for

*Coating helps protect against stomach upset our heart. Aspirin is not right for everyone. ask your doctor about other uses for safety coated 81 mg aspirin

Reye's syndrome: Children and teenagers who have or are recovering from chicker pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur, consult a doctor because ms could be an early sign of Reye's syndrome, a rare but serious illness. argy alert. Aspirin may cause a severe allergic reaction, which may include

■ hives ■ facial swelling ■ shock

store between 15-30°C (59-86°F)

Other information

asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

take other drugs containing prescription or nonprescription NSAIDs (aspirin

take a blood thinning (anticoagulant) or steroid drug

■ have 3 or more alcoholic drinks every day while using this product

ibuprofen, naproxen, or others)

take more or for a longer time than directed

 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: consult a doctor (1-800-222-1222) right away

especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may pregnant or breast-feeding, ask a health professional before use. It is ringing in the ears or loss of hearing occurs ese could be signs of a serious condition. out of reach of children. In case of overdose, get medical help or contact problems in the unborn child or complications during

Do not use If you are allergic to aspirin or any other pain reliever/fever reduce

If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn

Ask a doctor before use if

Drug Facts (continued under

AND CARE FOR H-E-B®

AN ANTONIO, TX 7820

 an allergic reaction occurs. Seek medical help right away.
 you experience any of the following signs of stomach bleeding: ■ have stomach pain that does not get better pain gets worse or lasts more than 10 days ever gets worse or lasts more than 3 days feel faint ■ vomit blood ■ have bloody or black stools redness or swelling is present

new symptoms occur

Ask a doctor or pharmacist before use if you are taking a prescription drug

8

PLD-0567A

Stop use and ask a doctor if

■ gout ■ diabetes

■ you are taking a diuretic
■ you have asthma you have high blood pressure, heart disease, liver cirrhosis, or kidney disease Drug Facts (continued)

H-E-B Low Dose Aspirin 81 mg

aury

Inactive ingredients D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch (com), silicon dioxide, sodium bicarbonate, sodium

ate, stearic acid, talc, titanium dioxide, triethyl citrate

LOW DOSE ASPIRIN

aspirin tablet, coated

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:37808-867

ORAL Route of Administration

Active Ingredient/Active Moiety

Basis of Strength Strength Ingredient Name ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) **ASPIRIN** 81 mg

Inactive Ingredients Ingredient Name Strength D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) FD&C YELLOW NO. 6 (UNII: H77VEI93A8) METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)

STARCH, CORN (UNII: 08232NY3SJ)

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

SODIUM BICARBONATE (UNII: 8MDF5V39QO)

SODIUM LAURYL SULFATE (UNII: 368GB5141J)

STEARIC ACID (UNII: 4ELV7Z65AP)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

| Product Characterist | roduct Characteristics | | | | |
|-----------------------------|------------------------|--------------|----------|--|--|
| Color | yellow | Score | no score | | |
| Shape | ROUND | Size | 7mm | | |
| Flavor | | Imprint Code | E | | |
| Contains | | | | | |

| ı | P | ackaging | | | | |
|---|---|----------------------|---|-------------------------|-----------------------|--|
| | # | Item Code | Package Description | Marketing Start Date | Marketing End Date | |
| | | NDC:37808- 867-30 | 300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 05/29/2020 | | |

| Marketing In | larketing Information | | | | |
|-----------------------|---|-------------------------|-----------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| OTC Monograph Drug | M013 | 05/29/2020 | | | |
| | | | | | |

Labeler - H E B (007924756)

Revised: 4/2024 HEB