DG HEALTH ADULT TUSSIN MULTI SYMPTOM COLD- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution Dolgencorp, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dolgencorp, LLC Adult Tussin Multi-Symptom Cold Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg
Guaifenesin, USP 200 mg
Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- · symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- · this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-888-309-9030

Principal Display Panel

Compare to the active ingredients of Robitussin® Multi-Symptom Cold

For Ages 12 & Over

Adult

Tussin

Dextromethorphan HBr

Cough Suppressant

Guaifenesin

Expectorant

Phenylephrine HCI

Nasal Decongestant

Multi-Symptom Cold

Relieves:

Cough

Nasal Congestion

Mucus

Non Drowsy

CF

4 FL OZ (118 mL)



Dextromethorphan HBr

Cough Suppressant

Phenylephrine HCl Nasal Decongestant

Multi-Symptom

Nasal Congestion

4 FL OZ (118 mL)

Guaifenesin

Cold

Relieves:

Mucus

Non Drowsy

DG™ health

Adult

Dextromethorphan HBr

Cough Suppressant

Phenylephrine HCl Nasal Decongestant

Multi-Symptom

Nasal Congestion

Guaifenesin **Expectorant**

Cold

Relieves:

Cough

Mucus

Non Drowsy

Compare to the active ingredients of Multi-Symptom Cold

For Ages 12 & Over

Drug Facts

For Ages 12 & Over

Active ingredients Purposes

Nasal Decongestant Phenylephrine HCI expectorant

Cough Suppressant uissn

DG" health

(in each 10 mL)
Dextromethorphan HBr,
USP 20 mg......C
Guaifenesin, USP 200 mg...
Phenylephrine HCl, .Cough suppressantNasal decongestant

USP 10 mg...

- Uses helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
 temporarily relieves these symptoms occurring with a cold:
- nasal congestion
 cough due to minor throat and

bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, sek a doctor or pharmacits before taking ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
heart disease high blood pressure
thyroid disease diabetes
trouble urinating due to an enlarged
prostate gland

- cough that occurs with too much
- phleam (mucus)
- prineign (mucus)

 = cough that lasts or is chronic such as occurs
 with smoking, asthma, chronic bronchitis,
 or emphysema

When using this product do not use more

- Stop use and ask a doctor if
 you get nervous, dizzy, or sleepless
 symptoms do not get better within 7 days
 or are accompanied by fever
 cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough

may be a sign of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- DIFECTIONS
 do not take more than 6 doses in any
 24-hour period
 measure only with dosing cup provided
 keep dosing cup with product
 ml = millitter
 this adult product is not intended for use
 in shill deny under 10 years of one

- in children under 12 years of age

Ш	age	dose
	adults and children	10 mL
Ш	12 years and over	every 4 hours
Ш	children under 12 years	do not use

Other information

- each 10 mL contains; sodium 5 mg
 store at 20-25°C (68-77°F). Do not refrigerate.
- Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium berzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE.TN 37072

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

100% Satisfaction Guaranteed! If you're not satisfied with this product for any reason, please call us so we can make you 100% satisfied.

(888) 309-9030

PARENTS: Learn about teen medicine abuse

www.StopMedicineAbuse.org





4 FL OZ (118 mL)



LOT NO.

EXP.

: 51626 VT C7



DG HEALTH ADULT TUSSIN MULTI SYMPTOM COLD

dextromethorphan hydrobromide, quaifenesin, phenylephrine hydrochloride solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-980
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		

Product Characteristics			
Color	RED (clear)	Score	
Shape		Size	
Flavor	FRUIT (Menthol)	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-980- 26	1 in 1 CARTON	11/09/2018	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/09/2018	

Labeler - Dolgencorp, LLC (068331990)

Revised: 4/2021 Dolgencorp, LLC