

**SEVERE SINUS CONGESTION AND PAIN NON-DROWSY, DAYTIME-
acetaminophen, guaifenesin and phenylephrine hcl tablet, film coated
Valu Merchandisers Company**

Best Choice 44-527C Delisted

Active ingredients (in each caplet)

Acetaminophen 325 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Expectorant
Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
 - minor aches and pains
 - nasal congestion
 - sinus congestion and pressure
 - headache
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over

- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

Best Choice®

**COMPARE TO THE ACTIVE INGREDIENTS
IN TYLENOL® SINUS SEVERE***

**Non-Drowsy
Severe Daytime
Sinus
Congestion & Pain**

ACETAMINOPHEN - Pain Reliever/Fever Reducer

Guaifenesin - Expectorant

Phenylephrine HCl - Nasal Decongestant

Relieves Headache, Fever,
Chest Congestion, Sinus
Pressure, Nasal Congestion

Pseudoephedrine Free

Actual Size

**24 Cool Caplets
with Cool Blast Flavor**

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® SINUS SEVERE. 50844 REV0820H52708

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106

Best Choice®
100% Guaranteed

www.bestchoicebrand.com

Best Choice®

COMPARE TO THE ACTIVE INGREDIENTS
IN **TYLENOL® SINUS SEVERE***

Non-Drowsy Severe Daytime Sinus Congestion & Pain

Actual Size



ACETAMINOPHEN - Pain Reliever/Fever Reducer
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

Relieves Headache, Fever,
Chest Congestion, Sinus
Pressure, Nasal Congestion
Pseudoephedrine Free

24 Cool Caplets
with Cool Blast Flavor

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

PROUDLY DISTRIBUTED BY:
VALU MERCHANTISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106
www.bestchoicebrand.com

No print/No varnish
Lot & Exp date

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® SINUS SEVERE. 50844 REV0820H52708

Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION
Active Ingredients
(in each caplet)
Acetaminophen 325 mg Pain reliever/fever reducer
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)
Uses
temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
headache
nasal congestion
minor aches and pains
sinus congestion and pressure

Drug Facts (continued)
Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Drug Facts (continued)
see end flap for expiration date and lot number
Inactive ingredients
corn starch, croscopolone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose.

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask a doctor or pharmacist before taking this product.
if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have:
cough that occurs with too much phlegm (mucus) or persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

headache that lasts. These could be signs of a serious condition.
if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
do not take more than directed
adults and children 12 years and over
take 2 caplets every 4 hours
swallow whole - do not crush, chew, or dissolve
do not take more than 10 caplets in 24 hours
children under 12 years: ask a doctor

Drug Facts (continued)

polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-426-9391

BEST CHOICE



B-2115-527C-08-SCR
REV0820H52708

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ heart disease ■ thyroid disease ■ diabetes ■ liver disease ■ high blood pressure ■ difficulty in urination due to enlargement of the prostate gland <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive ■ temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours</p> <ul style="list-style-type: none"> ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ blisters ■ rash ■ skin reddening <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p>
---	---

44-527C

SEVERE SINUS CONGESTION AND PAIN NON-DROWSY, DAYTIME

acetaminophen, guaifenesin and phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-527
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;527
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-527-08	2 in 1 CARTON	08/06/2005	08/29/2025
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/06/2005	08/29/2025

Labeler - Valu Merchandisers Company (868703513)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63941-527) , pack(63941-527)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63941-527)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63941-527)

Revised: 1/2024

Valu Merchandisers Company