

LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet
Discount Drug Mart

Loperamide Hydrochloride

Drug Facts

Active ingredient (in each caplet)

Loperamide Hydrochloride USP, 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert

Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert

Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store between 20° - 25°C (68° - 77°F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions?

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

**DISTRIBUTED BY DRUG MART-FOOD FAIR
MEDINA, OHIO 44256**

PRINCIPAL DISPLAY PANEL - 12 Caplet Blister Pack Carton

DISCOUNT
drug
mart

FOOD FAIR

MAY CONTAIN
ANTI-THEFT
DEVICE

See New Warnings
and Directions

LOPERAMIDE HYDROCHLORIDE
TABLETS USP, 2 mg

ANTI-DIARRHEAL

Anti-Diarrheal

Controls The
Symptoms Of Diarrhea

†COMPARE TO THE ACTIVE INGREDIENT OF IMODIUM[®] A-D

12 Caplets*

Each caplet (*capsule-shaped tablet)
contains Loperamide Hydrochloride USP, 2 mg



5197303



5197303

Drug Facts (continued)

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, dizziness, drowsiness or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over: 2 caplets after the first loose stool, 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

Drug Facts (continued)

Other information

- store between 20° - 25°C (68° - 77°F)
- see side panel for full instructions and expiration date
- TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D & C yellow 6, 10 aluminum lake, FD & C blue no. 1 aluminum lake, hydroxyethylcellulose, pregelatinized starch, vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions? Call toll-free Monday-Friday 8:30am to 5pm EST at 1-800-406-7984.

Drug Facts

Active ingredient (in each caplet)

Loperamide Hydrochloride USP, 2 mg

Purpose

Anti-diarrheal

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings

Allergy alert Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert Taking more than directed can cause serious heart problems or death

LOPERAMIDE HYDROCHLORIDE
TABLETS USP, 2 mg
ANTI-DIARRHEAL



MAY CONTAIN
**ANTI-THEFT
DEVICE**

See New Warnings
and Directions

LOPERAMIDE HYDROCHLORIDE
TABLETS USP, 2 mg
ANTI-DIARRHEAL

Anti-Diarrheal

Controls The
Symptoms Of Diarrhea



¹COMPARE TO THE ACTIVE INGREDIENT OF IMODIUM® A-D

12 Caplets*

Each caplet (*capsule-shaped tablet) contains Loperamide Hydrochloride USP, 2 mg

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D.

Expiration Date:

Lot No.:

NON VARNISH

Keep the carton. It contains important information.

R0419

DISTRIBUTED BY DRUG MART-FOOD FAIR
MEDINA, OHIO 44256 54
www.discount-drugmart.com



SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE
WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON
FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO:
DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-123
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77T135393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	9mm
Flavor		Imprint Code	123
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-123-12	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993	

2	NDC:53943-123-24	24 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/01/1993
---	------------------	---	------------

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	02/01/1993	

Labeler - Discount Drug Mart (047741335)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(53943-123)

Revised: 1/2024

Discount Drug Mart