

CHILDRENS ROBITUSSIN 12 HOUR COUGH RELIEF- dextromethorphan polistirex suspension, extended release
GlaxoSmithKline Consumer Healthcare Holdings (US) LLC

Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions

- **shake bottle well before use**
- measure only with dosing cup provided. Do not use dosing cup with other products

- dose as follows or as directed by doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: **sodium 5 mg**
- store at 20–25°C (68–77°F)
- dosing cup provided

Inactive ingredients (Grape flavor)

D&C red no. 30, FD&C blue no. 1, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Inactive ingredients (Orange flavor)

D&C red no. 30, D&C yellow no. 10, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions?

call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**.

You may also report side effects to this number.

For most recent product information, **visit www.robitussin.com**

Distributed by: Pfizer, Madison, NJ 07940 USA © 2016 Pfizer Inc.

PRINCIPAL DISPLAY PANEL - 89 mL Bottle Carton - Orange

NEW!

**Children's
Robitussin®**

**For adults & children
AGES 4 & OVER**

EXTENDED-RELEASE

12 Hour

Cough Relief

**DEXTROMETHORPHAN POLISTIREX EXTENDED-
RELEASE ORAL SUSPENSION (Cough Suppressant)**

12 Hour

Cough Relief

DAY or NIGHT

Orange Flavored Liquid

Alcohol-Free

3 FL OZ (89 mL)

**Contains sodium metabisulfite, a sulfite
that may cause allergic-type reactions**

Robitussin

EXTENDED-RELEASE

**12 Hour
Cough Relief**

Children's

NEW!

Robitussin

For adults & children **AGES 4 & OVER**

EXTENDED-RELEASE

**12 Hour
Cough Relief**

DEXTROMETHORPHAN POLISTIREX EXTENDED-
RELEASE ORAL SUSPENSION (Cough Suppressant)

**12 Hour
Cough Relief**



DAY or NIGHT

Orange Flavored Liquid
Alcohol-Free

3 FL OZ (89 mL)

Contains sodium metabisulfite, a sulfite
that may cause allergic-type reactions



Actual Size



LOT:

EXP:

Imprinted with "sealed for your protection" is broken or missing.

TAMPER EVIDENT: Do not use if cap is opened or cracked

Drug Facts

Active ingredient (in each 5 mL) Purpose
Dextromethorphan polistirex equivalent to 30 mg Cough suppressant
dextromethorphan hydrobromide, USP

Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have
■ chronic cough that lasts as occurs with smoking, asthma, or emphysema
■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if
■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
■ cough lasts more than 7 days, cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions
■ shake bottle well before use
■ measure only with dosing cup provided. Do not use dosing cup with other products
■ dose as follows or as directed by doctor
■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Drug Facts (continued)

Other information
■ each 5 mL contains: sodium 5 mg
■ store at 20-25°C (68-77°F)
■ dosing cup provided

Inactive ingredients D&C red no. 30, D&C yellow no. 10, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions? call weekdays from 9 AM to 5 PM EST at 1-800-762-4675. You may also report side effects to this number.

DOSING

SHAKE WELL BEFORE USE.

Measure only with dosing cup provided. Do not use dosing cup with other products. mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.

Contains NO fever reducer or pain reliever

DOSING CUP INCLUDED

Distributed by: Pfizer, Hudson, NJ 07940 USA © 2016 Pfizer Inc. For most recent product information, visit www.robitussin.com

Should be 18 or older to purchase

PARENTS:
Learn about teen medicine abuse

www.StopMedicineAbuse.org

PAA070657

LB8435 02/16 Rev 00

PRINCIPAL DISPLAY PANEL - 89 mL Bottle Carton - Grape

NEW!

Children's Robitussin®

For adults & children AGES 4 & OVER

EXTENDED-RELEASE

12 Hour Cough Relief

DEXTROMETHORPHAN POLISTIREX EXTENDED-

RELEASE ORAL SUSPENSION (Cough Suppressant)

12 Hour
Cough Relief

DAY or NIGHT

Grape Flavored Liquid
Alcohol-Free

3 FL OZ (89 mL)

**Contains sodium metabisulfite, a sulfite
that may cause allergic-type reactions**

Robitussin

EXTENDED-RELEASE

12 Hour Cough Relief

Children's

NEW!

Robitussin

For adults & children **AGES 4 & OVER**

EXTENDED-RELEASE

12 Hour Cough Relief

DEXTROMETHORPHAN POLISTIREX EXTENDED-
RELEASE ORAL SUSPENSION (Cough Suppressant)

12 Hour Cough Relief

DAY or NIGHT



Grape Flavored Liquid
Alcohol-Free

3 FL OZ (89 mL)

Contains sodium metabisulfite, a sulfite
that may cause allergic-type reactions




Actual Size



3 0031 8726 10 5

LOT:

EXP:



Imprinted with "sealed for your protection" is broken or missing.

TAMPER EVIDENT: Do not use if carton is opened or deformed

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Dextromethorphan polistirex equivalent to 30 mg
 dextromethorphan hydrobromide, USPCough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have
 ■ chronic cough that lasts as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Step use and ask a doctor if
 ■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
 ■ cough lasts more than 7 days, cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions
 ■ shake bottle well before use
 ■ measure only with dosing cup provided. Do not use dosing cup with other products
 ■ dose as follows or as directed by doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Drug Facts (continued)

Other information
 ■ each 5 mL contains sodium 5 mg
 ■ store at 20-25°C (68-77°F)
 ■ dosing cup provided

Inactive ingredients D&C red no. 30, FD&C blue no. 1, flavor, glycerin, high fructose corn syrup, methylparaben, polyorbate 90, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions? call weekdays from 9 AM to 5 PM EST at 1-800-762-4675. You may also report side effects to this number.

DOSE
 SHAKE WELL BEFORE USE.
 Measure only with dosing cup provided.
 Do not use dosing cup with other products.
 mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.
 Contains **NO** fever reducer or pain reliever
DOSEING CUP INCLUDED

Distributed by Pfizer, Madison, NJ 07941 USA © 2016 Pfizer Inc.
 For most recent product information, visit www.robitussin.com

Should be 18 or older to purchase

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

PAA070659

LB8432 02/16 Rev 00

CHILDRENS ROBITUSSIN 12 HOUR COUGH RELIEF

dextromethorphan polistirex suspension, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-8725
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLISTIREX (UNII: 5H9W9GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031-8725-10	1 in 1 CARTON	07/05/2016	
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	07/05/2016	

CHILDRENS ROBITUSSIN 12 HOUR COUGH RELIEF

dextromethorphan polistirex suspension, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-8726
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLISTIREX (UNII: 5H9W9GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031-8726-10	1 in 1 CARTON	07/05/2016	
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	07/05/2016	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)

Revised: 12/2022

GlaxoSmithKline Consumer Healthcare Holdings (US) LLC