

**REMODULIN- treprostinil injection, solution**  
**STERILE DILUENT FOR REMODULIN- water injection, solution**  
**United Therapeutics Corporation**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use REMODULIN safely and effectively. See full prescribing information for REMODULIN.**

**REMODULIN® (treprostinil) Injection, for subcutaneous or intravenous use**  
**Initial U.S. Approval: May 2002**

----- **INDICATIONS AND USAGE** -----

Remodulin is a prostacyclin mimetic indicated for:

- Treatment of pulmonary arterial hypertension (PAH; WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%). (1.1)
- Patients who require transition from epoprostenol, to reduce the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition. (1.2)

----- **DOSAGE AND ADMINISTRATION** -----

PAH WHO Group 1 in patients with NYHA Class II-IV symptoms:

- Initial dose for patients new to prostacyclin infusion therapy: 1.25 ng/kg/min; increase based on clinical response (increments of 1.25 ng/kg/min per week for the first 4 weeks of treatment, later 2.5 ng/kg/min per week). Avoid abrupt cessation. (2.2, 2.4)
- Mild to moderate hepatic insufficiency: Decrease initial dose to 0.625 ng/kg/min.  
Severe hepatic insufficiency: No studies performed. (2.5)

Transition from Epoprostenol:

Increase the Remodulin dose gradually as the epoprostenol dose is decreased, based on constant observation of response. (2.7)

Administration:

Continuous subcutaneous infusion is the preferred mode. Use intravenous (IV) infusion if subcutaneous infusion is not tolerated. (2.1, 2.6)

----- **DOSAGE FORMS AND STRENGTHS** -----

- Remodulin is supplied in 20-mL vials containing 2 mg (0.1 mg/mL), 4 mg (0.2 mg/mL), 8 mg (0.4 mg/mL), 20 mg (1 mg/mL), 50 mg (2.5 mg/mL), 100 mg (5 mg/mL), 200 mg (10 mg/mL), or 400 mg (20 mg/mL) of treprostinil. (3)

----- **CONTRAINDICATIONS** -----

None (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Chronic intravenous infusions delivered using an external infusion pump with an indwelling central venous catheter are associated with the risk of blood stream infections (BSIs) and sepsis, which may be fatal. (5.1)
- Do not abruptly lower the dose or withdraw dosing. (5.2)
- Remodulin may cause symptomatic hypotension. (5.4)
- Remodulin inhibits platelet aggregation and increases the risk of bleeding. (5.5)

----- **ADVERSE REACTIONS** -----

Most common adverse reactions (incidence ~QZgt;3%) reported in clinical studies with Remodulin: subcutaneous infusion site pain and reaction, headache, diarrhea, nausea, jaw pain, vasodilatation, edema, and hypotension. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact United Therapeutics Corp. at 1-866-458-6479 or contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

----- **DRUG INTERACTIONS** -----

- Remodulin dosage adjustment may be necessary if inhibitors or inducers of CYP2C8 are added or

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

- 1.1 Pulmonary Arterial Hypertension
- 1.2 Pulmonary Arterial Hypertension in Patients Requiring Transition from Epoprostenol

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 General
- 2.2 Initial Dose for Patients New to Prostacyclin Infusion Therapy
- 2.3 Initial Dose for Patients Transitioning to an Implantable Intravenous Infusion Pump
- 2.4 Dosage Adjustments
- 2.5 Patients with Hepatic Insufficiency
- 2.6 Administration
- 2.7 Patients Requiring Transition from Epoprostenol

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Risk of Catheter-Related Bloodstream Infection
- 5.2 Worsening PAH upon Abrupt Withdrawal or Sudden Large Dose Reduction
- 5.3 Patients with Hepatic Insufficiency
- 5.4 Risk of Symptomatic Hypotension
- 5.5 Risk of Bleeding

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Post-Marketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 Effect of CYP2C8 Inhibitors and Inducers on Treprostinil

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Patients with Hepatic Insufficiency
- 8.7 Patients with Renal Impairment

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

- 14.1 Clinical Trials in Pulmonary Arterial Hypertension (PAH)
- 14.2 Flolan-To-Remodulin Transition Study

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

**FULL PRESCRIBING INFORMATION**

**1 INDICATIONS AND USAGE**

**1.1 Pulmonary Arterial Hypertension**

Remodulin is indicated for the treatment of pulmonary arterial hypertension (PAH; WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%) [see *Clinical Studies (14.1)*].

**1.2 Pulmonary Arterial Hypertension in Patients Requiring Transition from Epoprostenol**

In patients with PAH requiring transition from epoprostenol, Remodulin is indicated to diminish the rate of clinical deterioration. Consider the risks and benefits of each drug prior to transition.

**2 DOSAGE AND ADMINISTRATION**

**2.1 General**

Remodulin can be administered with or without further dilution with Sterile Diluent for Remodulin or similar approved high-pH glycine diluent (e.g., Sterile Diluent for Flolan or Sterile Diluent for Epoprostenol), Sterile Water for Injection, or 0.9% Sodium Chloride Injection prior to administration. See Table 1 below for storage and administration time limits for the different diluents.

Diluted Remodulin has been shown to be stable at ambient temperature when stored for up to 14 days using high-pH glycine diluent at concentrations as low as 0.004 mg/mL (4,000 ng/mL).

**Table 1: Selection of Diluent**

Diluent	Storage Limits	Administration Limits
None	See Section 16	16 weeks at 40~QZ #176;C

Sterile Diluents for Remodulin, Flolan, or Epoprostenol	14 days at room temperature	48 hours at 40~QZ #176;C
Sterile Water for Injection 0.9% Sodium Chloride for Injection	4 hours at room temperature or 24 hours refrigerated	48 hours at 40~QZ #176;C

## 2.2 Initial Dose for Patients New to Prostacyclin Infusion Therapy

Remodulin is indicated for subcutaneous (SC) or intravenous (IV) use only as a continuous infusion. Remodulin is preferably infused subcutaneously, but can be administered by a central intravenous line if the subcutaneous route is not tolerated because of severe site pain or reaction. The infusion rate is initiated at 1.25 ng/kg/min. If this initial dose cannot be tolerated because of systemic effects, reduce the infusion rate to 0.625 ng/kg/min.

## 2.3 Initial Dose for Patients Transitioning to an Implantable Intravenous Infusion Pump

The initial dose of Remodulin should be the same as the current dose the patient is receiving using the external infusion pump at the time of transition.

## 2.4 Dosage Adjustments

The goal of chronic dosage adjustments is to establish a dose at which PAH symptoms are improved, while minimizing excessive pharmacologic effects of Remodulin (headache, nausea, emesis, restlessness, anxiety, and infusion site pain or reaction).

The infusion rate should be increased in increments of 1.25 ng/kg/min per week for the first four weeks of treatment and then 2.5 ng/kg/min per week for the remaining duration of infusion, depending on clinical response. Dosage adjustments may be undertaken more often if tolerated. Avoid abrupt cessation of infusion [see *Warnings and Precautions* (5.2)]. Restarting a Remodulin infusion within a few hours after an interruption can be done using the same dose rate. Interruptions for longer periods may require the dose of Remodulin to be re-titrated.

## 2.5 Patients with Hepatic Insufficiency

In patients with mild or moderate hepatic insufficiency, decrease the initial dose of Remodulin to 0.625 ng/kg/min ideal body weight. Remodulin has not been studied in patients with severe hepatic insufficiency [see *Warnings and Precautions* (5.3), *Use in Specific Populations* (8.6), and *Clinical Pharmacology* (12.3)].

## 2.6 Administration

Inspect parenteral drug products for particulate matter and discoloration prior to administration whenever solution and container permit. If either particulate matter or discoloration is noted, do not use.

### Preparation

Remodulin is administered by subcutaneous or intravenous infusion at a calculated rate based on a patient's dose (ng/kg/min), weight (kg), and the Remodulin concentration (mg/mL).

For administration of **Undiluted Remodulin** the rate is calculated using the following formula:

$$\text{Undiluted Infusion Rate (mL/hour)} = \frac{\text{Dose (ng/kg/min)} \times \text{Weight (kg)} \times 0.00006^*}{\text{Remodulin Vial Strength (mg/mL)}}$$

\* Conversion factor of 0.00006 = 60 min/hour; 0.000001 mg/ng

For administration of **Diluted Remodulin**, the concentration is calculated using the following formula:

Step 1

$$\text{Diluted Remodulin Concentration (mg/mL)} = \frac{\text{Dose (ng/kg/min)} \times \text{Weight (kg)} \times 0.00006}{\text{Infusion Rate (mL/hour)}}$$

The volume of Remodulin Injection needed to make the required diluted Remodulin concentration for the given reservoir size can then be calculated using the following formula:

Step 2

$$\text{Volume of Remodulin Injection (mL)} = \frac{\text{Diluted Remodulin Concentration (mg/mL)} \times \text{Remodulin Vial Strength (mg/mL)}}{\text{Total Volume of Diluted Remodulin Solution in Reservoir (mL)}}$$

The calculated volume of Remodulin Injection is then added to the reservoir along with the sufficient volume of diluent to achieve the desired total volume in the reservoir.

### Subcutaneous Infusion

Remodulin is administered subcutaneously by continuous infusion, via a subcutaneous catheter, using an infusion pump designed for subcutaneous drug delivery. The infusion pump should: (1) be adjustable to approximately 0.002 mL/hour, (2) have occlusion/no delivery, low battery, programming error and motor malfunction alarms, (3) have delivery accuracy of ~QZ #177;6% or better, (4) be positive pressure-driven, and (5) have a reservoir made of polyvinyl chloride, polypropylene or glass. Alternatively, use an infusion pump cleared for use with Remodulin. To avoid potential interruptions in drug

delivery, the patient must have immediate access to a backup infusion pump and subcutaneous infusion sets.

## Intravenous Infusion

### *External Intravenous Infusion Pump:*

Remodulin is administered intravenously by continuous infusion via a surgically placed indwelling central venous catheter using an external infusion pump designed for intravenous drug delivery. If clinically necessary, a temporary peripheral intravenous cannula, preferably placed in a large vein, may be used for short term administration of Remodulin. Use of a peripheral intravenous infusion for more than a few hours increases the risk of thrombophlebitis. The infusion pump used to administer Remodulin should: (1) have occlusion/no delivery, low battery, programming error and motor malfunction alarms, (2) have delivery accuracy of ~QZ #177;6% or better, (3) be positive pressure driven, and (4) have a reservoir made of polyvinyl chloride, polypropylene or glass. Alternatively, use an infusion pump cleared for use with Remodulin. To avoid potential interruptions in drug delivery, the patient must have immediate access to a backup infusion pump and infusion sets.

Infusion sets with an in-line 0.22- or 0.2-micron pore size filter should be used.

### *Implantable Intravenous Infusion Pump:*

Use an implantable intravenous infusion pump approved for use with Remodulin, such as the Implantable System for Remodulin~QZ #174; (ISR). Refer to the pump manufacturer~QZapos;s manual for specific instructions regarding preparation, programing, implantation, and refilling.

## **2.7 Patients Requiring Transition from Epoprostenol**

Transition from epoprostenol to Remodulin is accomplished by initiating the infusion of Remodulin and increasing it, while simultaneously reducing the dose of intravenous epoprostenol. The transition to Remodulin should take place in a hospital with constant observation of response (e.g., walk distance and signs and symptoms of disease progression). Initiate Remodulin at a recommended dose of 10% of the current epoprostenol dose, and then escalate as the epoprostenol dose is decreased (see Table 2 for recommended dose titrations).

Patients are individually titrated to a dose that allows transition from epoprostenol therapy to Remodulin while balancing prostacyclin-limiting adverse events. Treat increases in the patient~QZapos;s symptoms of PAH first with increases in the dose of Remodulin. Treat side effects normally associated with prostacyclin and prostacyclin analogs first by decreasing the dose of epoprostenol.

**Table 2: Recommended Transition Dose Changes**

Step	Epoprostenol Dose	Remodulin Dose
1	Unchanged	10% Starting Epoprostenol Dose
2	80% Starting Epoprostenol Dose	30% Starting Epoprostenol Dose
3	60% Starting Epoprostenol Dose	50% Starting Epoprostenol Dose

4	40% Starting Epoprostenol Dose	70% Starting Epoprostenol Dose
5	20% Starting Epoprostenol Dose	90% Starting Epoprostenol Dose
6	5% Starting Epoprostenol Dose	110% Starting Epoprostenol Dose
7	0	110% Starting Epoprostenol Dose + additional 5-10% increments as needed

### 3 DOSAGE FORMS AND STRENGTHS

20-mL vial containing 2 mg treprostinil (0.1 mg per mL).

20-mL vial containing 4 mg treprostinil (0.2 mg per mL).

20-mL vial containing 8 mg treprostinil (0.4 mg per mL).

20-mL vial containing 20 mg treprostinil (1 mg per mL).

20-mL vial containing 50 mg treprostinil (2.5 mg per mL).

20-mL vial containing 100 mg treprostinil (5 mg per mL).

20-mL vial containing 200 mg treprostinil (10 mg per mL).

20-mL vial containing 400 mg treprostinil (20 mg per mL).

### 4 CONTRAINDICATIONS

None

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Risk of Catheter-Related Bloodstream Infection

Chronic intravenous infusions of Remodulin delivered using an external infusion pump with an indwelling central venous catheter are associated with the risk of blood stream infections (BSIs) and sepsis, which may be fatal. Therefore, continuous subcutaneous infusion is the preferred mode of administration.

In an open-label study of IV treprostinil (n=47) using an external infusion pump, there were seven catheter-related line infections during approximately 35 patient years, or about 1 BSI event per 5 years of use. A CDC survey of seven sites that used IV treprostinil for the treatment of PAH found approximately 1 BSI (defined as any positive blood culture) event per 3 years of use. Administration of IV Remodulin with a high pH glycine diluent has been associated with a lower incidence of BSIs when compared to neutral diluents (sterile water, 0.9% sodium chloride) when used along with catheter care guidelines.

In an open-label study of an implantable pump (n=60), there were two blood stream

infections (BSIs) related to the implant procedure during approximately 265 patient years.

## 5.2 Worsening PAH upon Abrupt Withdrawal or Sudden Large Dose Reduction

Avoid abrupt withdrawal or sudden large reductions in dosage of Remodulin, which may result in worsening of PAH symptoms.

## 5.3 Patients with Hepatic Insufficiency

Titrate Remodulin slowly in patients with hepatic insufficiency, because such patients will likely be exposed to greater systemic concentrations relative to patients with normal hepatic function [see *Dosage and Administration* (2.5), *Use in Specific Populations* (8.6), and *Clinical Pharmacology* (12.3)].

## 5.4 Risk of Symptomatic Hypotension

Treprostinil is a pulmonary and systemic vasodilator. In patients with low systemic arterial pressure, treatment with Remodulin may produce symptomatic hypotension.

## 5.5 Risk of Bleeding

Remodulin inhibits platelet aggregation and increases the risk of bleeding.

# 6 ADVERSE REACTIONS

The following adverse reactions are discussed elsewhere in labeling: Infections associated with intravenous administration [see *Warnings and Precautions* (5.1)].

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

### Adverse Events with Subcutaneously Administered Remodulin

Patients receiving Remodulin as a subcutaneous infusion reported a wide range of adverse events, many potentially related to the underlying disease (dyspnea, fatigue, chest pain, right ventricular heart failure, and pallor). During clinical trials with subcutaneous infusion of Remodulin, infusion site pain and reaction were the most common adverse events among those treated with Remodulin. Infusion site reaction was defined as any local adverse event other than pain or bleeding/bruising at the infusion site and included symptoms such as erythema, induration, or rash. Infusion site reactions were sometimes severe and could lead to discontinuation of treatment.

**Table 3: Percentages of Subjects Reporting Subcutaneous Infusion Site Adverse Events**

	Reaction		Pain	
	Placebo	Remodulin	Placebo	Remodulin
Severe	1	38	2	39



Requiring narcotics*	NA†	NA†	1	32
Leading to discontinuation	0	3	0	7

\* based on prescriptions for narcotics, not actual use

† medications used to treat infusion site pain were not distinguished from those used to treat site reactions

Other adverse events included diarrhea, jaw pain, edema, vasodilatation, and nausea, and these are generally considered to be related to the pharmacologic effects of Remodulin, whether administered subcutaneously or intravenously.

### Adverse Reactions during Chronic Dosing

Table 4 lists adverse reactions that occurred at a rate of at least 3% more frequent in patients treated with subcutaneous Remodulin than with placebo in controlled trials in PAH.

**Table 4: Adverse Reactions in Controlled 12-Week Studies of Subcutaneous Remodulin and at least 3% more frequent than on Placebo**

<b>Adverse Reaction</b>	<b>Remodulin (N=236) Percent of Patients</b>	<b>Placebo (N=233) Percent of Patients</b>
Infusion Site Pain	85	27
Infusion Site Reaction	83	27
Headache	27	23
Diarrhea	25	16
Nausea	22	18
Rash	14	11
Jaw Pain	13	5
Vasodilatation	11	5
Edema	9	3

Reported adverse reactions (at least 3% more frequent on drug than on placebo) are included with the exception of those too general to be informative, and those not plausibly attributable to the use of the drug, because they were associated with the condition being treated or are very common in the treated population.

While hypotension occurred in both groups, the event was experienced twice as frequently in the Remodulin group as compared to the placebo group (4% in Remodulin treatment group versus 2% in placebo-controlled group). As a potent vasodilator, hypotension is possible with the administration of Remodulin.

The safety of Remodulin was also studied in a long-term, open-label extension study in which 860 patients were dosed for a mean duration of 1.6 years, with a maximum exposure of 4.6 years. Twenty-nine (29%) percent achieved a dose of at least 40

ng/kg/min (max: 290 ng/kg/min). The safety profile during this chronic dosing study was similar to that observed in the 12-week placebo-controlled study except for the following suspected adverse drug reactions (occurring in at least 3% of patients): anorexia, vomiting, infusion site infection, asthenia, and abdominal pain.

### **Adverse Events Attributable to the Drug Delivery System**

In controlled studies of Remodulin administered subcutaneously, there were no reports of infection related to the drug delivery system. There were 187 infusion system complications reported in 28% of patients (23% Remodulin, 33% placebo); 173 (93%) were pump related and 14 (7%) related to the infusion set. Eight of these patients (4 Remodulin, 4 placebo) reported non-serious adverse events resulting from infusion system complications. Adverse events resulting from problems with the delivery systems were typically related to either symptoms of excess Remodulin (e.g., nausea) or return of PAH symptoms (e.g., dyspnea). These events were generally resolved by correcting the delivery system pump or infusion set problem, such as replacing the syringe or battery, reprogramming the pump, or straightening a crimped infusion line. Adverse events resulting from problems with the delivery system did not lead to clinical instability or rapid deterioration. In addition to these adverse events due to the drug delivery system during subcutaneous administration, the following adverse events may be attributable to the IV mode of infusion including arm swelling, paresthesia, hematoma, and pain [see *Warnings and Precautions (5.1)*].

## **6.2 Post-Marketing Experience**

In addition to adverse reactions reported from clinical trials, the following events have been identified during post-approval use of Remodulin. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. The following events have been chosen for inclusion because of a combination of their seriousness, frequency of reporting, and potential connection to Remodulin. These events are thrombophlebitis associated with peripheral intravenous infusion, thrombocytopenia, bone pain, pruritus, dizziness, arthralgia, myalgia/muscle spasm, and pain in extremity. In addition, generalized rashes, sometimes macular or papular in nature, and cellulitis have been infrequently reported.

## **7 DRUG INTERACTIONS**

### **7.1 Effect of CYP2C8 Inhibitors and Inducers on Treprostinil**

Dose adjustment of treprostinil may be necessary when co-administered with CYP2C8 inducers or inhibitors. Human pharmacokinetic studies with an oral formulation of treprostinil (treprostinil diolamine) indicated that co-administration of the cytochrome P450 (CYP) 2C8 enzyme inhibitor gemfibrozil increases exposure (both  $C_{max}$  and AUC) to treprostinil. Co-administration of the CYP2C8 enzyme inducer rifampin decreases exposure to treprostinil. It has not been determined if the changes in exposure of treprostinil with inhibitors or inducers of CYP2C8 observed for the oral administration of treprostinil would be similar for treprostinil administered via the parenteral route [see *Clinical Pharmacology (12.3)*].

## **8 USE IN SPECIFIC POPULATIONS**

## 8.1 Pregnancy

### Risk Summary

Limited case reports of treprostinil use in pregnant women are insufficient to inform a drug-associated risk of adverse developmental outcomes. However, there are risks to the mother and the fetus associated with pulmonary arterial hypertension (*see Clinical Considerations*). In animal studies, no adverse reproductive and developmental effects were seen in rats at about 123 and 48 times the human exposure based on  $C_{\max}$  and AUC, respectively. In rabbits, external fetal and soft tissue malformations and skeletal malformations were observed at about 7 and 5 times the human exposure based on  $C_{\max}$  and AUC, respectively (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

### Clinical Considerations

#### *Disease-associated maternal and embryo-fetal risk*

Pulmonary arterial hypertension is associated with an increased risk of maternal and fetal mortality.

### Data

Animal reproduction studies have been conducted with treprostinil via continuous subcutaneous administration and with treprostinil diolamine administered orally. In pregnant rats, continuous subcutaneous infusions of treprostinil during organogenesis and late gestational development, at doses as high as 900 ng treprostinil/kg/min (about 117 times the starting human subcutaneous infusion rate, on a  $\text{ng}/\text{m}^2$  basis and about 16 times the average rate achieved in clinical trials), resulted in no evidence of harm to the fetus. In pregnant rabbits, effects of continuous subcutaneous infusions of treprostinil during organogenesis were limited to an increased incidence of fetal skeletal variations (bilateral full rib or right rudimentary rib on lumbar 1) associated with maternal toxicity (reduction in body weight and food consumption) at a dose of 150 ng treprostinil/kg/min (about 41 times the starting human subcutaneous infusion rate, on a  $\text{ng}/\text{m}^2$  basis, and 5 times the average rate used in clinical trials). In rats, continuous subcutaneous infusion of treprostinil from implantation to the end of lactation, at doses of up to 450 ng treprostinil/kg/min, did not affect the growth and development of offspring. In studies with orally administered treprostinil diolamine, no adverse effect doses for fetal viability/growth, fetal development (teratogenicity), and postnatal development were determined in rats. In pregnant rats, no evidence of harm to the fetus was observed following oral administration of treprostinil diolamine at the highest dose tested (20 mg/kg/day), which represents about 123 and 48 times the human exposure, when based on  $C_{\max}$  and AUC of the average subcutaneous infusion rate achieved in clinical trials, respectively. In pregnant rabbits, external fetal and soft tissue malformations and fetal skeletal malformation occurred. The dose at which no adverse effects were seen (0.5 mg/kg/day) represents about 7 and 5 times the human exposure, when based on  $C_{\max}$  and AUC of the average subcutaneous infusion rate achieved in clinical trials, respectively. No treprostinil treatment-related effects on labor

and delivery were seen in animal studies. Animal reproduction studies are not always predictive of human response.

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of treprostinil in human milk, the effects on the breastfed infant, or the effects on milk production.

## **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established. Clinical studies of Remodulin did not include sufficient numbers of patients aged ~QZ #8804;16 years to determine whether they respond differently from older patients.

## **8.5 Geriatric Use**

Clinical studies of Remodulin did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## **8.6 Patients with Hepatic Insufficiency**

Remodulin clearance is reduced in patients with hepatic insufficiency. In patients with mild or moderate hepatic insufficiency, decrease the initial dose of Remodulin to 0.625 ng/kg/min ideal body weight, and monitor closely. Remodulin has not been studied in patients with severe hepatic insufficiency [see *Dosage and Administration* (2.5), *Warnings and Precautions* (5.3), and *Clinical Pharmacology* (12.3)].

## **8.7 Patients with Renal Impairment**

No dose adjustments are required in patients with renal impairment. Treprostinil is not cleared by dialysis [see *Clinical Pharmacology* (12.3)].

## **10 OVERDOSAGE**

Signs and symptoms of overdose with Remodulin during clinical trials are extensions of its dose-limiting pharmacologic effects and include flushing, headache, hypotension, nausea, vomiting, and diarrhea. Most events were self-limiting and resolved with reduction or withholding of Remodulin.

In controlled clinical trials using an external infusion pump, seven patients received some level of overdose and in open-label follow-on treatment seven additional patients received an overdose; these occurrences resulted from accidental bolus administration of Remodulin, errors in pump programmed rate of administration, and prescription of an incorrect dose. In only two cases did excess delivery of Remodulin produce an event of substantial hemodynamic concern (hypotension, near-syncope).

One pediatric patient was accidentally administered 7.5 mg of Remodulin via a central venous catheter. Symptoms included flushing, headache, nausea, vomiting,

hypotension, and seizure-like activity with loss of consciousness lasting several minutes. The patient subsequently recovered.


## 11 DESCRIPTION

Remodulin (treprostinil) Injection is a sterile solution of treprostinil, a prostacyclin mimetic, formulated for subcutaneous or intravenous administration. Remodulin is supplied in 20-mL multidose vials in eight strengths, containing 2 mg (0.1 mg/mL), 4 mg (0.2 mg/mL), 8 mg (0.4 mg/mL), 20 mg (1 mg/mL), 50 mg (2.5 mg/mL), 100 mg (5 mg/mL), 200 mg (10 mg/mL), or 400 mg (20 mg/mL) of treprostinil. Each mL also contains 5.3 mg sodium chloride (except for the 10 mg/mL and 20 mg/mL strengths, which contain 4.0 mg sodium chloride), 3 mg metacresol, 6.3 mg sodium citrate dihydrate, and water for injection. Sodium hydroxide and hydrochloric acid may be added to adjust pH between 6.0 and 7.2.

Treprostinil is chemically stable at room temperature and neutral pH.

Treprostinil is (1*R*,2*R*,3*aS*,9*aS*)-[[2,3,3*a*,4,9,9*a*-hexahydro-2-hydroxy-1-[(3*S*)-3-hydroxyoctyl]-1*H*-benz[*f*]inden-5-yl]oxy]acetic acid. Treprostinil has a molecular weight of 390.52 and a molecular formula of C<sub>23</sub>H<sub>34</sub>O<sub>5</sub>.

The structural formula of treprostinil is:



Sterile Diluent for Remodulin is a high-pH (pH~10.4) glycine diluent supplied in a 50-mL vial containing 50 mL of Sterile Diluent for Remodulin. Each vial contains 94 mg glycine, 73.3 mg sodium chloride, sodium hydroxide (to adjust pH), and water for injection.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The major pharmacologic actions of treprostinil are direct vasodilation of pulmonary and systemic arterial vascular beds, and inhibition of platelet aggregation.

### 12.2 Pharmacodynamics

In animals, the vasodilatory effects reduce right and left ventricular afterload and increase cardiac output and stroke volume. Other studies have shown that treprostinil causes a dose-related negative inotropic and lusitropic effect. No major effects on cardiac conduction have been observed.

Treprostinil produces vasodilation and tachycardia. Single doses of treprostinil up to 84 mcg by inhalation produce modest and short-lasting effects on QTc, but this is apt to be

an artifact of the rapidly changing heart rate. Treprostinil administered by the subcutaneous or intravenous routes has the potential to generate concentrations many-fold greater than those generated via the inhaled route; the effect on the QTc interval when treprostinil is administered parenterally has not been established.

### **12.3 Pharmacokinetics**

The pharmacokinetics of continuous subcutaneous Remodulin are linear over the dose range of 2.5 to 125 ng/kg/min (corresponding to plasma concentrations of about 260 pg/mL to 18,250 pg/mL) and can be described by a two-compartment model. Dose proportionality at infusion rates greater than 125 ng/kg/min has not been studied.

Subcutaneous and intravenous administration of Remodulin demonstrated bioequivalence at steady state at a dose of 10 ng/kg/min.

#### **Absorption**

Remodulin is relatively rapidly and completely absorbed after subcutaneous infusion, with an absolute bioavailability approximating 100%. Steady-state concentrations occurred in approximately 10 hours. Concentrations in patients treated with an average dose of 9.3 ng/kg/min were approximately 2,000 ng/L.

#### **Distribution**

The volume of distribution of the drug in the central compartment is approximately 14 L/70 kg ideal body weight. Remodulin at *in vitro* concentrations well above what is clinically relevant was 91% bound to human plasma protein.

#### **Metabolism and Excretion**

Treprostinil is substantially metabolized by the liver, primarily by CYP2C8. In a study conducted in healthy volunteers using [<sup>14</sup>C] treprostinil, 79% and 13% of the subcutaneous dose was recovered in the urine and feces, respectively, over 10 days. Only 4% was excreted as unchanged treprostinil in the urine. Five metabolites were detected in the urine, ranging from 10% to 16% and representing 64% of the dose administered. Four of the metabolites are products of oxidation of the 3-hydroxyoctyl side chain and one is a glucuroconjugated derivative (treprostinil glucuronide). The identified metabolites do not appear to have activity.

The elimination of treprostinil (following subcutaneous administration) is biphasic, with a terminal elimination half-life of approximately 4 hours using a two-compartment model. Systemic clearance is approximately 30 L/hour for a 70 kg person.

Based on *in vitro* studies treprostinil does not inhibit or induce major CYP enzymes.

#### **Specific Populations**

##### ***Hepatic Insufficiency***

In patients with portopulmonary hypertension and mild (n=4) or moderate (n=5) hepatic insufficiency, Remodulin at a subcutaneous dose of 10 ng/kg/min for 150 minutes had a C<sub>max</sub> that was 2-fold and 4-fold, respectively, and an AUC<sub>0-~QZ#8734</sub> that was 3-fold and 5-fold, respectively, values observed in healthy subjects. Clearance in patients with hepatic insufficiency was reduced by up to 80% compared to healthy adults.

##### ***Renal Impairment***

In patients with severe renal impairment requiring dialysis (n=8), administration of a single 1 mg dose of orally administered treprostinil pre- and post-dialysis resulted in an AUC<sub>0-inf</sub> that was not significantly altered compared to healthy subjects.

## **Drug Interaction Studies**

### ***Effect of CYP2C8 Inhibitors and Inducers on Treprostinil***

Co-administration of an oral formulation of treprostinil (treprostinil diolamine) with gemfibrozil (600 mg twice a day), a CYP2C8 enzyme inhibitor, doubles the AUC and C<sub>max</sub> of treprostinil in healthy adults. Co-administration of an oral formulation of treprostinil (treprostinil diolamine) with rifampin (600 mg/day), a CYP2C8 enzyme inducer, decreases AUC of treprostinil by 22%.

### ***Effect of Treprostinil on Cytochrome P450 Enzymes***

*In vitro* studies of human hepatic microsomes showed that treprostinil does not inhibit cytochrome P450 (CYP) isoenzymes CYP1A2, CYP2A6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A. Additionally, treprostinil does not induce CYP1A2, CYP2B6, CYP2C9, CYP2C19, and CYP3A isoenzymes.

### ***Effect of Other Drugs on Treprostinil***

Human pharmacokinetic studies with an oral formulation of treprostinil (treprostinil diolamine) indicated that co-administration of the cytochrome P450 (CYP) 2C8 enzyme inhibitor gemfibrozil increases exposure (both C<sub>max</sub> and AUC) to treprostinil. Co-administration of the CYP2C8 enzyme inducer rifampin decreases exposure to treprostinil.

Drug interaction studies have been carried out with treprostinil (oral or subcutaneous) co-administered with acetaminophen (4 g/day), esomeprazole (40 mg/day), bosentan (250 mg/day), sildenafil (60 mg/day), warfarin (25 mg/day), and fluconazole (200 mg/day), respectively, in healthy volunteers. These studies did not show a clinically significant effect on the pharmacokinetics of treprostinil. Treprostinil does not affect the pharmacokinetics or pharmacodynamics of warfarin. The pharmacokinetics of R- and S-warfarin and the INR in healthy subjects given a single 25 mg dose of warfarin were unaffected by continuous subcutaneous infusion of treprostinil at an infusion rate of 10 ng/kg/min.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

A two-year rat carcinogenicity study was performed with treprostinil inhalation at target doses of 5.26, 10.6, and 34.1 mcg/kg/day. There was no evidence for carcinogenic potential associated with treprostinil inhalation in rats at systemic exposure levels up to about 34 and 1 times the human exposure, when based on C<sub>max</sub> and AUC of the average subcutaneous infusion rate achieved in clinical trials, respectively. *In vitro* and *in vivo* genetic toxicology studies did not demonstrate any mutagenic or clastogenic effects of treprostinil. Treprostinil sodium did not affect fertility or mating performance of male or female rats given continuous subcutaneous (sc) infusions at rates of up to 450 ng treprostinil/kg/min [about 59 times the recommended starting human sc infusion rate (1.25 ng/kg/min) and 8 times the average rate (9.3 ng/kg/min) achieved in clinical

trials, on a ng/m<sup>2</sup> basis]. In this study, males were dosed from 10 weeks prior to mating and through the 2-week mating period. Females were dosed from 2 weeks prior to mating until gestational day 6.

Treprostinil diolamine did not demonstrate any carcinogenic effects in mouse or rat carcinogenicity studies. Oral administration of treprostinil diolamine to Tg.rasH2 mice at 0, 5, 10, and 20 mg/kg/day in males and 0, 3, 7.5, and 15 mg/kg/day in females daily for 26 weeks did not significantly increase the incidence of tumors. The exposures, when based on AUC, obtained at the highest dose levels used in males and females are about 7- and 15-fold, respectively, the human exposure of the average subcutaneous infusion rate achieved in clinical trials. Oral administration of treprostinil diolamine to Sprague Dawley rats at 0, 1, 3, and 10 mg/kg/day daily for 104 weeks did not significantly increase the incidence of tumors. The exposures obtained at the highest dose levels used in males and females are about 18- and 26-fold, respectively, the human exposure of the average subcutaneous infusion rate achieved in clinical trials.

Treprostinil diolamine was tested *in vivo* in a rat micronucleus assay and did not induce an increased incidence of micronucleated polychromatic erythrocytes.

**14 CLINICAL STUDIES**

**14.1 Clinical Trials in Pulmonary Arterial Hypertension (PAH)**

Two 12-week, multicenter, randomized, double-blind studies compared continuous subcutaneous infusion of Remodulin to placebo in a total of 470 patients with NYHA Class II (11%), III (81%), or IV (7%) PAH. PAH was idiopathic/heritable in 58% of patients, associated with connective tissue diseases in 19%, and the result of congenital systemic-to-pulmonary shunts in 23%. The mean age was 45 (range 9 to 75 years). About 81% were female and 84% were Caucasian. Pulmonary hypertension had been diagnosed for a mean of 3.8 years. The primary endpoint of the studies was change in 6-minute walking distance, a standard measure of exercise capacity. There were many assessments of symptoms related to heart failure, but local discomfort and pain associated with Remodulin may have substantially unblinded those assessments. The 6-minute walking distance and an associated subjective measurement of shortness of breath during the walk (Borg dyspnea score) were administered by a person not participating in other aspects of the study. Remodulin was administered as a subcutaneous infusion, described in Section 2, DOSAGE AND ADMINISTRATION, and the dose averaged 9.3 ng/kg/min at Week 12. Few subjects received doses greater than 40 ng/kg/min. Background therapy, determined by the investigators, could include anticoagulants, oral vasodilators, diuretics, digoxin, and oxygen, but not an endothelin receptor antagonist or epoprostenol. The two studies were identical in design and conducted simultaneously, and the results were analyzed both pooled and individually.

**Hemodynamic Effects**

As shown in Table 5, chronic therapy with Remodulin resulted in small hemodynamic changes consistent with pulmonary and systemic vasodilation.

**Table 5: Hemodynamics during Chronic Administration of Remodulin in Patients with PAH in 12-Week Studies**

		Mean change from
--	--	------------------



Hemodynamic Parameter	Baseline		baseline at Week 12	
	Remodulin (N=204-231)	Placebo (N=215-235)	Remodulin (N=163-199)	Placebo (N=182-215)
CI (L/min/m <sup>2</sup> )	2.4 ~QZ #177; 0.88	2.2 ~QZ #177; 0.74	+0.12 ~QZ #177; 0.58*	-0.06 ~QZ #177; 0.55
PAPm (mmHg)	62 ~QZ #177; 17.6	60 ~QZ #177; 14.8	-2.3 ~QZ #177; 7.3*	+0.7 ~QZ #177; 8.5
RAPm (mmHg)	10 ~QZ #177; 5.7	10 ~QZ #177; 5.9	-0.5 ~QZ #177; 5.0*	+1.4 ~QZ #177; 4.8
PVRI (mmHg/L/min/m <sup>2</sup> )	26 ~QZ #177; 13	25 ~QZ #177; 13	-3.5 ~QZ #177; 8.2*	+1.2 ~QZ #177; 7.9
SVRI (mmHg/L/min/m <sup>2</sup> )	38 ~QZ #177; 15	39 ~QZ #177; 15	-3.5 ~QZ #177; 12*	-0.80 ~QZ #177; 12
SvO <sub>2</sub> (%)	62 ~QZ #177; 100	60 ~QZ #177; 11	+2.0 ~QZ #177; 10*	-1.4 ~QZ #177; 8.8
SAPm (mmHg)	90 ~QZ #177; 14	91 ~QZ #177; 14	-1.7 ~QZ #177; 12	-1.0 ~QZ #177; 13
HR (bpm)	82 ~QZ #177; 13	82 ~QZ #177; 15	-0.5 ~QZ #177; 11	-0.8 ~QZ #177; 11

CI, cardiac index; PAPm, mean pulmonary arterial pressure; PVRI, pulmonary vascular resistance indexed; RAPm, mean right atrial pressure; SAPm, mean systemic arterial pressure; SVRI, systemic vascular resistance indexed; SvO<sub>2</sub>, mixed venous oxygen saturation; HR, heart rate

\* Denotes statistically significant difference between Remodulin and placebo, p~QZlt;0.05.

## Clinical Effects

The effect of Remodulin on 6-minute walk, the primary endpoint of the 12-week studies, was small and did not achieve conventional levels of statistical significance. For the combined populations, the median change from baseline on Remodulin was 10 meters and the median change from baseline on placebo was 0 meters from a baseline of approximately 345 meters. Although it was not the primary endpoint of the study, the Borg dyspnea score was significantly improved by Remodulin during the 6-minute walk, and Remodulin also had a significant effect, compared with placebo, on an assessment that combined walking distance with the Borg dyspnea score. Remodulin also consistently improved indices of dyspnea, fatigue, and signs and symptoms of pulmonary hypertension, but these indices were difficult to interpret in the context of


incomplete blinding to treatment assignment resulting from infusion site symptoms.

## 14.2 Flolan-To-Remodulin Transition Study

In an 8-week, multicenter, randomized, double-blind, placebo-controlled study, patients on stable doses of Flolan were randomly withdrawn from Flolan to placebo or Remodulin. Fourteen Remodulin and 8 placebo patients completed the study. The primary endpoint of the study was the time to clinical deterioration, defined as either an increase in Flolan dose, hospitalization due to PAH, or death. No patients died during the study.

During the study period, Remodulin effectively prevented clinical deterioration in patients transitioning from Flolan therapy compared to placebo (Figure 1). Thirteen of 14 patients in the Remodulin arm were able to transition from Flolan successfully, compared to only 1 of 8 patients in the placebo arm ( $p=0.0002$ ).

**Figure 1: Time to Clinical Deterioration for PAH Patients Transitioned from Flolan to Remodulin or Placebo in an 8-Week Study**



## 16 HOW SUPPLIED/STORAGE AND HANDLING

Remodulin is supplied in 20-mL multidose vials as sterile solutions in water for injection, individually packaged in cartons. Unopened vials of Remodulin are stable until the date indicated when stored at 25~QZ #176;C (77~QZ #176;F), with excursions permitted to 2-30~QZ #176;C (36-86~QZ #176;F). A single vial of Remodulin should be used for no more than 30 days after the initial introduction into the vial.

Remodulin Injection is supplied as:

Remodulin	Concentration	NDC
-----------	---------------	-----

2 mg / 20 mL	0.1 mg/mL	66302-111-01
4 mg / 20 mL	0.2 mg/mL	66302-112-01
8 mg / 20 mL	0.4 mg/mL	66302-114-01
20 mg / 20 mL	1 mg/mL	66302-101-01
50 mg / 20 mL	2.5 mg/mL	66302-102-01
100 mg / 20 mL	5 mg/mL	66302-105-01
200 mg / 20 mL	10 mg/mL	66302-110-01
400 mg / 20 mL	20 mg/mL	66302-120-01

Sterile Diluent for Remodulin is supplied separately as:  
50-mL vial, carton of 1 (NDC 66302-150-50).

## 17 PATIENT COUNSELING INFORMATION

### *Interruption of Therapy*

Advise patients and caregivers to seek medical attention if they experience signs or symptoms of abrupt withdrawal of therapy or suspect a pump malfunction [see *Warnings and Precautions (5.2)*].

### *Overdose*

Inform patients and their caregivers to seek medical attention if they experience signs or symptoms of Remodulin overdose [see *Overdosage (10)*].

~QZ #169; Copyright 2023 United Therapeutics Corp. All rights reserved.

REMODULIN manufactured for:

United Therapeutics Corp.  
Research Triangle Park, NC 27709

## PRINCIPAL DISPLAY PANEL - 0.1 mg/mL Vial Carton


REMODULIN~QZ #174;  
(treprostinil) Injection

2 mg/20 mL  
(0.1 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION



## PRINCIPAL DISPLAY PANEL - 0.2 mg/mL Vial Carton


REMODULIN~QZ #174;  
(treprostinil) Injection

4 mg/20 mL  
(0.2 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION



## PRINCIPAL DISPLAY PANEL - 0.4 mg/mL Vial Carton

REMODULIN~QZ #174;  
(treprostinil) Injection


8 mg/20 mL  
(0.4 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION





## PRINCIPAL DISPLAY PANEL - 2.5 mg/mL Vial Carton

REMODULIN~QZ #174;  
(treprostinil) Injection


50 mg/20 mL  
(2.5 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION





## PRINCIPAL DISPLAY PANEL - 5 mg/mL Vial Carton

REMODULIN~QZ #174;  
(treprostinil) Injection


100 mg/20 mL  
(5 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION





## PRINCIPAL DISPLAY PANEL - 10 mg/mL Vial Carton

REMODULIN~QZ #174;  
(treprostinil) Injection


200 mg/20 mL  
(10 mg/mL)

For Subcutaneous  
and Intravenous Infusion

20 mL multidose vial

United  
Therapeutics  
CORPORATION





## PRINCIPAL DISPLAY PANEL - 50 mL Vial Carton

NDC 66302-150-50

STERILE DILUENT  
FOR REMODULIN~QZ #174;

1 x 50 mL  
VIAL

Contains drug diluent for use only  
with Intravenous Infusion of  
REMODULIN~QZ #174; (treprostinil) Injection


Each vial contains 94 mg glycine,  
73.3 mg sodium chloride, sodium  
hydroxide (added to adjust pH), and  
Water for Injection.

For dilution information see package

For dilution information see package insert for REMODULIN~QZ#174; (treprostinil) Injection.

Store at 20 - 25~QZ#176;C (68 - 77~QZ#176;F), excursions permitted to 15 - 30~QZ#176;C (59 - 86~QZ#176;F) [See USP Controlled Room Temperature].

DO NOT FREEZE



REMODULIN

treprostinil injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-111	
Route of Administration	INTRAVENOUS, SUBCUTANEOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)		treprostinil	2 mg in 20 mL	
Inactive Ingredients				
Ingredient Name			Strength	
trisodium citrate dihydrate (UNII: B22547B95K)				
sodium chloride (UNII: 451W47IQ8X)				
metacresol (UNII: GGO4Y809LO)				
sodium hydroxide (UNII: 55X04QC32I)				
water (UNII: 059QF0KO0R)				
hydrochloric acid (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-111-01	1 in 1 BOX	09/28/2023	03/26/2024
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021272	09/28/2023	03/26/2024

REMODULIN			
treprostinil injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-112
Route of Administration	INTRAVENOUS, SUBCUTANEOUS		
Active Ingredient/Active Moiety			

Ingredient Name		Basis of Strength	Strength	
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)		treprostinil	4 mg in 20 mL	
Inactive Ingredients				
Ingredient Name			Strength	
trisodium citrate dihydrate (UNII: B22547B95K)				
sodium chloride (UNII: 451W47IQ8X)				
metacresol (UNII: GGO4Y809LO)				
sodium hydroxide (UNII: 55X04QC32I)				
water (UNII: 059QF0KO0R)				
hydrochloric acid (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-112-01	1 in 1 BOX	09/28/2023	03/26/2024
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021272	09/28/2023	03/26/2024

REMODULIN

treprostinil injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-114
Route of Administration	INTRAVENOUS, SUBCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)	treprostinil	8 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
trisodium citrate dihydrate (UNII: B22547B95K)	
sodium chloride (UNII: 451W47IQ8X)	
metacresol (UNII: GGO4Y809LO)	

<b>sodium hydroxide</b> (UNII: 55X04QC32I)				
<b>water</b> (UNII: 059QF0KO0R)				
<b>hydrochloric acid</b> (UNII: QTT17582CB)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-114-01	1 in 1 BOX	09/28/2023	
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021272	09/28/2023	

REMODULIN				
treprostinil injection, solution				
Product Information				
Product Type		HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-101
Route of Administration		INTRAVENOUS, SUBCUTANEOUS		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)			treprostinil	20 mg in 20 mL
Inactive Ingredients				
Ingredient Name				Strength
trisodium citrate dihydrate (UNII: B22547B95K)				
sodium chloride (UNII: 451W47IQ8X)				
metacresol (UNII: GGO4Y809LO)				
sodium hydroxide (UNII: 55X04QC32I)				
water (UNII: 059QF0KO0R)				
hydrochloric acid (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-101-01	1 in 1 BOX	05/22/2002	

1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA021272		05/22/2002	

REMODULIN				
treprostinil injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-102	
Route of Administration	INTRAVENOUS, SUBCUTANEOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)		treprostinil	50 mg in 20 mL	
Inactive Ingredients				
Ingredient Name			Strength	
trisodium citrate dihydrate (UNII: B22547B95K)				
sodium chloride (UNII: 451W47IQ8X)				
metacresol (UNII: GGO4Y809LO)				
sodium hydroxide (UNII: 55X04QC32I)				
water (UNII: 059QF0KO0R)				
hydrochloric acid (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-102-01	1 in 1 BOX	05/22/2002	
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NDA	NDA021272		05/22/2002	



## REMODULIN

treprostinil injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:66302-105
<b>Route of Administration</b>	INTRAVENOUS, SUBCUTANEOUS		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)	treprostinil	100 mg in 20 mL

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
trisodium citrate dihydrate (UNII: B22547B95K)	
sodium chloride (UNII: 451W47IQ8X)	
metacresol (UNII: GGO4Y809LO)	
sodium hydroxide (UNII: 55X04QC32I)	
water (UNII: 059QF0KO0R)	
hydrochloric acid (UNII: QTT17582CB)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-105-01	1 in 1 BOX	05/22/2002	
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA021272	05/22/2002	

## REMODULIN

treprostinil injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:66302-110
<b>Route of Administration</b>	INTRAVENOUS, SUBCUTANEOUS		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)			treprostinil	200 mg in 20 mL
Inactive Ingredients				
Ingredient Name				Strength
trisodium citrate dihydrate (UNII: B22547B95K)				
sodium chloride (UNII: 451W47IQ8X)				
metacresol (UNII: GGO4Y809LO)				
sodium hydroxide (UNII: 55X04QC32I)				
water (UNII: 059QF0KO0R)				
hydrochloric acid (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-110-01	1 in 1 BOX	05/22/2002	
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021272	05/22/2002	

REMODULIN			
treprostinil injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66302-120
Route of Administration	INTRAVENOUS, SUBCUTANEOUS		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
treprostinil (UNII: RUM6K67ESG) (treprostinil - UNII:RUM6K67ESG)		treprostinil	400 mg in 20 mL
Inactive Ingredients			
Ingredient Name			Strength
trisodium citrate dihydrate (UNII: B22547B95K)			

<b>sodium chloride</b> (UNII: 451W47IQ8X)	
<b>metacresol</b> (UNII: GGO4Y809LO)	
<b>sodium hydroxide</b> (UNII: 55X04QC32I)	
<b>water</b> (UNII: 059QF0KO0R)	
<b>hydrochloric acid</b> (UNII: QTT17582CB)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-120-01	1 in 1 BOX	07/30/2021	01/26/2022
1		20 mL in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021272	07/30/2021	01/26/2022

## STERILE DILUENT FOR REMODULIN

water injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:66302-150
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>water</b> (UNII: 059QF0KO0R) (water - UNII:059QF0KO0R)	water	1 mL in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>glycine</b> (UNII: TE7660XO1C)	
<b>sodium chloride</b> (UNII: 451W47IQ8X)	
<b>sodium hydroxide</b> (UNII: 55X04QC32I)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66302-150-50	1 in 1 BOX	12/22/2014	
		50 mL in 1 VIAL; Type 0: Not a Combination		

1	50 mL in 1 VIAL; Type O: Not a Combination Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA021272	12/22/2014	

**Labeler** - United Therapeutics Corporation (965460025)

Revised: 6/2025

United Therapeutics Corporation