

**HEALTH MART NAPROXEN SODIUM- naproxen sodium tablet, film coated
McKesson**

McKesson Naproxen Sodium Tablets, 220 mg Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 caplet every 8 to 12 hours while symptoms last• for the first dose you may take 2 caplets within the first hour• do not exceed 2 caplets in any 8- to 12-hour period• do not exceed 3 caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

compare to Aleve® active ingredient

naproxen sodium tablets, 220 mg

pain reliever/fever reducer (NSAID)

ACTUAL SIZE

strength to last 12 hours – gluten free

(**capsule-shaped tablets)



DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active Ingredient (in each caplet) **Purposes**
Naproxen sodium, 220 mg Pain reliever/
fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
minor pain of arthritis muscular aches
backache menstrual cramps
headache toothache
the common cold
temporarily reduces fever

Warnings
Allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling
asthma (wheezing) rash blisters
skin reddening
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

E3 98 8479E :

EXP.

LOT NO.



Drug Facts (continued)

Do not use
if you have ever had an allergic reaction to any other pain reliever/fever reducer
right before or after heart surgery
Ask a doctor before use if
the stomach bleeding warning applies to you
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
you are taking a diuretic
you have problems or serious side effects from taking pain relievers or fever reducers
you have asthma

Ask a doctor or pharmacist before use if you are
under a doctor's care for any serious condition
taking any other drug
When using this product
take with food or milk if stomach upset occurs
the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
you experience any of the following signs of stomach bleeding:
feel faint
vomit blood
have bloody or black stools
have stomach pain that does not get better
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
you have difficulty swallowing
it feels like the pill is stuck in your throat
redness or swelling is present in the painful area
any new symptoms appear


Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
do not take more than directed
the smallest effective dose should be used
drink a full glass of water with each dose
Adults and children 12 years and older
take 1 caplet every 8 to 12 hours while symptoms last
for the first dose you may take 2 caplets within the first hour
do not exceed 2 caplets in any 8- to 12-hour period
do not exceed 3 caplets in a 24-hour period
Children under 12 years
ask a doctor

Other Information
each caplet contains: sodium 21 mg
sims at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
Inactive Ingredients FD&C Blue no. 2 aluminum lake, hypromellose, mannitol, stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

MCKESSON
Distributed by McKesson
One First Street, San Francisco, CA 94104
www.mckesson.com/healthcare/brand
Bayer HealthCare LLC, distributor of Aleve®.
This product is not manufactured or distributed by
Bayer HealthCare LLC, distributor of Aleve®.
KEEP CARTON FOR REFERENCE
Questions or comments?
1-800-719-9260
Drug Facts (continued)



NDC 62011-0017-2
compare to Aleve® active ingredient
Health Mart PHARMACY
naproxen sodium
tablets, 220 mg
pain reliever/fever reducer (NSAID)
strength to last 12 hours • gluten free
100 caplets** (**capsule-shaped tablets)



HEALTH MART NAPROXEN SODIUM
naproxen sodium tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0017
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg	
Inactive Ingredients				
Ingredient Name			Strength	
HYPROMELLOSES (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)				
POVIDONES (UNII: FZ989GH94E)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
Product Characteristics				
Color	BLUE (Light Blue)	Score	no score	
Shape	CAPSULE (oval)	Size	12mm	
Flavor		Imprint Code	L368	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0017-1	1 in 1 CARTON	11/21/2011	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:62011-0017-2	1 in 1 CARTON	11/21/2011	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:62011-0017-3	1 in 1 CARTON	11/13/2016	
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074661	11/21/2011		

Labeler - McKesson (177667227)

Revised: 11/2016

McKesson