SEVERE COLD PLUS FLU DAYTIME NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate CVS Pharmacy,Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### CVS Health™ Severe Cold & Flu Daytime and Nighttime

#### SEVERE COLD + FLU DAYTIME

### **Drug Facts**

Active ingredients (in each packet)	Purposes
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20	Cough suppressant
mg	
Guaifenesin 400 mg	Expectorant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - sore throat
  - cough
  - nasal congestion
  - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus
- temporarily reduces fever

# Warnings

## Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 packets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

#### **ALLERGY ALERT**

Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- Skin reddening
- Blisters

### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

# Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough with excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

# When using this product do not exceed recommended dosage

# Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be

- signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than the recommended dose
- take every 4 hours; do not exceed 6 packets in 24 hours or as directed by a doctor
- adults and children 12 years and over: dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

### Other information

- each packet contains: potassium 6 mg
- store at room temperature

### **Inactive ingredients**

anhydrous citric acid, flavors, maltodextrin, potassium citrate, silica, sucralose, sucrose, yellow#6

#### Questions or comments?

1-866-467-2748

#### SEVERE COLD + FLU NIGHTTIME

## **Drug Facts**

Active ingredients (in each packet)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20	Cough suppressant
mg	
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine hydrochloride 10 mg	Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - headache

- minor aches and pains
- cough
- sore throat
- nasal congestion
- sinus congestion and pressure
- runny nose
- sneezing
- temporarily reduces fever

### **Warnings**

### Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 5 packets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### **ALLERGY ALERT**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters

Is a skin or general allergic reaction occurs, stop use and seek medical help right away

# Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use to sedate children.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

# Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than the recommended dose
- take every 4 hours; do not exceed 5 packets in 24 hours or as directed by a doctor
- adults and children 12 years and over: dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

### Other information

- each packet contains: potassium 6 mg
- store at room temperature

### **Inactive ingredients**

anhydrous citric acid, flavors, maltodextrin, potassium citrate, silica, sucralose, sucrose, yellow#6

#### Questions or comments?

1-866-467-2748

#### PRINCIPAL DISPLAY PANEL-

Compare to Alka-Seltzer PLUS® Severe Cold +Flu Day and Night Active Ingredients\*

#### **DAYTIME**

#### **SEVERE COLD & FLU**

Acetaminophen / Pain reliever-fever reducer

Dextromethorphan HBr / Cough suppressant

Guaifenesin / Expectorant Phenylephrine HCI / Nasal decongestant

#### Relieves:

- Nasal Congestion Headache
- Sore Throat Body Ache Cough
- Mucus Chest Congestion Fever

## **Honey Lemon Flavor**

**Naturally and Artificially Flavored** 

#### **NIGHTTIME**

#### Severe Cold & Flu

Acetaminophen / Pain reliever-fever reducer

Dextromethorphan HBr / Cough suppressant Doxylamine succinate / Antihistamine Phenylephrine HCl / Nasal decongestant

#### Relieves:

- Nasal Congestion
  Headache
- Sore Throat Body Ache
- Cough Runny Nose Fever

# **Honey Lemon Flavor**

#### TAMPER EVIDENT: DO NOT USE IF INDIVIDUAL POUCH IS TORN OR OPEN

# DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

6 DAY PACKETS + 6 NIGHT PACKETS 12 TOTAL

Distributed by: CVS Pharmacy, Inc.

One CVS Drive Woonsocket, RI 02895

©2017 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

**Money Back Guarantee** 

\*These products is not manufactured or distributed by Bayer Healthcare, LLC, distributor of Alka-Seltzer plus® Severe Cold +Flu Day and Night.



### SEVERE COLD PLUS FLU DAYTIME NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate kit

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-546

Packaging			
# Item Cod	e Package Description	Marketing Start Date	Marketing End Date
1 NDC:69842-5	1 in 1 CARTON; Type 0: Not a Combination Product	07/14/2017	

Quantity of Parts				
Part #	Package Quantity	Total Product Quantity		
Part 1	1 CARTON	6		
Part 2	1 CARTON	6		

### Part 1 of 2

### SEVERE COLD PLUS FLU DAYTIME

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride powder, for solution

#### **Product Information**

Route of Administration ORAL

#### **Active Ingredient/Active Moiety Strength Ingredient Name Basis of Strength** ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) **ACETAMINOPHEN** 500 mg **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) DEXTROMETHORPHAN 20 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) **GUAIFENES IN** 400 mg PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -PHENYLEPHRINE 10 mg UNII:1WS297W6MV) **HYDROCHLORIDE**

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM CITRATE (UNII: EE900NI6FF)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1		6 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/14/2017	

# Part 2 of 2

# **SEVERE COLD PLUS FLU NIGHTTIME**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride powder, for solution

# **Product Information**

**Route of Administration** ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg	
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
POTASSIUM CITRATE (UNII: EE90ONI6FF)			
SILICON DIOXIDE (UNII: ETJ7Z 6XBU4)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

I	SUCROSE (UNII: C151H8M554)	
ı	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

	Packaging				
	#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
l	1		6 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	07/14/2017		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	07/14/2017		

Labeler - CVS Pharmacy,Inc. (062312574)

Revised: 9/2023 CVS Pharmacy,Inc.