

LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet
Select Brand

Drug Facts

ACTIVE INGREDIENT (IN EACH CAPLET)

Loperamide HCl USP, 2 mg

PURPOSE

Anti-diarrheal

USES

Controls symptoms of diarrhea, including Travelers' Diarrhea

WARNINGS

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use

If you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

Taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- diarrhea lasts for more than 2 days
- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

OTHER INFORMATION

- store between 20° – 25° C (68° – 77° F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

INACTIVE INGREDIENTS

Anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no.10, FD&C blue no.1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

QUESTIONS?

Call **1800-406-7984**

PRINCIPAL DISPLAY PANEL

select brand®

NDC 15127-338-12

Loperamide Hydrochloride Tablets USP, 2 mg

ANTI-DIARRHEAL


Controls The Symptoms Of Diarrhea

■ **Compare to the active ingredient of Imodium® A-D**

12 Caplets*

Each Caplet (*capsule-shaped tablet) contains Loperamide HCl USP, 2 mg

5048836/R1109



5048836



5048836

Questions? call 1-800-406-7984

Inactive ingredients anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10, FD&C blue no. 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Other information
■ store between 20° - 25° C (68° - 77° F) ■ see side panel for lot number and expiration date
■ TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS.
DO NOT USE IF ANY ARE TORN OR BROKEN.

12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Directions
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

When using this product
■ dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.
■ Stop use and ask a doctor if ■ diarrhea lasts for more than 2 days ■ symptoms get worse
■ you get abdominal swelling or bulging. These may be signs of a serious condition.
■ If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Ask a doctor before use if you have
■ fever ■ mucus in the stool ■ a history of liver disease
Ask a doctor or pharmacist before use if you are taking antibiotics

Drug Facts (continued)

Warnings
Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Use controls symptoms of diarrhea, including Travelers' Diarrhea
Active ingredient (in each caplet) Loperamide HCl USP, 2 mg
Purpose Anti-diarrheal

ANTI-DIARRHEAL

Loperamide Hydrochloride
Tablets USP, 2 mg

select brand
the lower price name brand

select brand
the lower price name brand

NDC 15127-338-12

Loperamide Hydrochloride
Tablets USP, 2 mg

ANTI-DIARRHEAL

Controls The Symptoms Of Diarrhea

*Compare to the active ingredient of Imodium® A-D



12 Caplets*
Each caplet (*capsule-shaped tablet) contains Loperamide HCl USP, 2 mg

Keep the carton.
It contains important information.

*This product is not manufactured or distributed by McNeil-PPC, Inc., distributor of Imodium® A-D. Imodium® is a registered trademark of Johnson & Johnson.

PI1109
Distributed by:
SELECT BRAND® DISTRIBUTORS
Pine Bluff, AR 71603 USA

UPC PROOF OF PURCHASE



0 15127 100420 6

Expiration Date:

NON VARNISH

Batch No.

LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:15127-338
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
TRISTEARIN (UNII: P6OCJ2551R)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	green	Score	2 pieces
Shape	CAPSULE	Size	9mm
Flavor		Imprint Code	123
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:15127-338-66	6 in 1 BLISTER PACK		
2	NDC:15127-338-12	12 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	02/01/1993	

Labeler - Select Brand (043562370)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(15127-338)

Revised: 9/2012

Select Brand