

VANCOMYCIN HYDROCHLORIDE- vancomycin hydrochloride capsule

Orient Pharma Co.,Ltd Yunlin Plant

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VANCOMYCIN HYDROCHLORIDE CAPSULES safely and effectively. See full prescribing information for VANCOMYCIN HYDROCHLORIDE CAPSULES. VANCOMYCIN HYDROCHLORIDE capsules, for oral use Initial U.S. Approval: 1986 To reduce the development of drug-resistant bacteria and maintain the effectiveness of vancomycin hydrochloride capsules and other antibacterial drugs, vancomycin hydrochloride capsules should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria.

INDICATIONS AND USAGE

Vancomycin hydrochloride capsules are glycopeptide antibacterial indicated in adult and pediatric patients (less than 18 years of age) for the treatment of: (1)

- *C. difficile* -associated diarrhea
- Enterocolitis caused by *Staphylococcus aureus* (including methicillin- resistant strains)

Limitations of Use: (1) (5.1)

- Parenteral administration of vancomycin is not effective for the above infections; therefore, vancomycin hydrochloride capsules must be given orally for these infections.
- Orally administered vancomycin hydrochloride capsules is not effective for other types of infections.
- To reduce the development of drug-resistant bacteria and maintain the effectiveness of vancomycin hydrochloride capsules and other antibacterial drugs, vancomycin hydrochloride capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. (1)

DOSAGE AND ADMINISTRATION

- *C. difficile*- associated diarrhea:
 - Adult Patients (18 years of age or greater): 125 mg orally 4 times daily for 10 days. (2.1)
 - Pediatric Patients (less than 18 years of age): 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g. (2.2)
- Staphylococcal enterocolitis:
 - Adult Patients (18 years of age or greater): 500 mg to 2 g orally in 3 or 4 divided doses for 7 to 10 days. (2.1)
 - Pediatric Patients (less than 18 years of age): 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g. (2.2)

DOSAGE FORMS AND STRENGTHS

- 125 mg capsules (3)
- 250 mg capsules (3)

CONTRAINDICATIONS

Hypersensitivity to vancomycin (4)

WARNINGS AND PRECAUTIONS

- **Vancomycin hydrochloride capsules must be given orally for treatment of staphylococcal enterocolitis and *C. difficile* -associated diarrhea. Orally administered vancomycin hydrochloride capsules are not effective for other types of infections.** (5.1)
- Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of vancomycin hydrochloride capsules for active *C. difficile* -associated diarrhea. Monitoring of serum concentrations may be appropriate in some instances. (5.2)
- Nephrotoxicity has occurred following oral vancomycin hydrochloride capsules therapy and can occur either during or after completion of therapy. The risk is increased in geriatric patients (5.3) Monitor renal function.
- Ototoxicity has occurred in patients receiving vancomycin hydrochloride capsules. (5.4) Assessment of auditory function may be appropriate in some instances.
- Severe Dermatologic Reactions: Discontinue VANCOMYCIN at the first appearance of skin rashes, mucosal lesions, or blisters. (5.5)

- Prescribing vancomycin hydrochloride capsules in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria. (5.6)
- The 125 mg strength capsules contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. (5.7)

-----ADVERSE REACTIONS-----

The most common adverse reactions ($\geq 10\%$) were nausea (17%), abdominal pain (15 %), and hypokalemia (13%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Orient Pharma Co., Ltd. at +886-5-631-1331 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch .

-----DRUG INTERACTIONS-----

No drug interaction studies have been conducted. (7)

-----USE IN SPECIFIC POPULATIONS-----

- **Pediatrics:** Safety and effectiveness in patients <18 years of age have not been established. (8.4)
- **Geriatrics:** In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with vancomycin hydrochloride capsules to detect potential vancomycin induced nephrotoxicity. (5.3) (6.1) (8.5) (14.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 6/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Adults

2.2 Pediatric Patients (less than 18 years of age)

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Oral Use Only

5.2 Potential for Systemic Absorption

5.3 Nephrotoxicity

5.4 Ototoxicity

5.5 Severe Dermatologic Reactions

5.6 Development of Drug-Resistant Bacteria

5.7 Risk of Allergic Reactions due to Tartrazine

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Diarrhea Associated With *Clostridium Difficile*

16 HOW SUPPLIED/STORAGE AND HANDLING

15 REFERENCES

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Vancomycin Hydrochloride Capsules are indicated for the treatment of *C. difficile* - associated diarrhea. Vancomycin Hydrochloride Capsules is also used for the treatment of enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains). in adult and pediatric patients less than 18 years of age.

Limitations of Use

- Parenteral administration of vancomycin is not effective for the above infections; therefore, Vancomycin Hydrochloride Capsules must be given orally for these infections.
- Orally administered Vancomycin Hydrochloride Capsules is not effective for other types of infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin Hydrochloride Capsules and other antibacterial drugs, Vancomycin Hydrochloride Capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Adults

Vancomycin Hydrochloride Capsules are used in treating *C. difficile* -associated diarrhea and staphylococcal enterocolitis.

- *C. difficile*- associated diarrhea: The recommended dose is 125 mg administered orally 4 times daily for 10 days.
- Staphylococcal enterocolitis: Total daily dosage is 500 mg to 2 g administered orally in 3 or 4 divided doses for 7 to 10 days.

2.2 Pediatric Patients (less than 18 years of age)

For both *C. difficile*-associated diarrhea and staphylococcal enterocolitis, the usual daily dosage is 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g.

3 DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride Capsules USP 125 mg* CAPSULES have a blue cap and yellow body imprinted with "OP64" on both cap and body in white ink.

Vancomycin Hydrochloride Capsules USP 250 mg* CAPSULES have a blue cap and dark-yellow body imprinted with "OP65" on both cap and body in white ink.

*Equivalent to vancomycin.

4 CONTRAINDICATIONS

Vancomycin Hydrochloride Capsules are contraindicated in patients with known hypersensitivity to vancomycin.

5 WARNINGS AND PRECAUTIONS

5.1 Oral Use Only

Vancomycin Hydrochloride Capsules for the treatment of colitis is for oral use only and is not systemically absorbed. Vancomycin Hydrochloride Capsules must be given orally for treatment of staphylococcal enterocolitis and *Clostridium difficile*-associated diarrhea. Orally administered Vancomycin Hydrochloride Capsules are not effective for other types of infections.

Parenteral administration of vancomycin is not effective for treatment of staphylococcal enterocolitis and *C. difficile*-associated diarrhea. If parenteral vancomycin therapy is desired, use an intravenous preparation of vancomycin and consult the package insert accompanying that preparation .

5.2 Potential for Systemic Absorption

Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of Vancomycin Hydrochloride Capsules for active *C. difficile* -associated diarrhea. Some patients with inflammatory disorders of the intestinal mucosa also may have significant systemic absorption of vancomycin. These patients may be at risk for the development of adverse reactions associated with higher doses of Vancomycin Hydrochloride Capsules; therefore, monitoring of serum concentrations of vancomycin may be appropriate in some instances, e.g., in patients with renal insufficiency and/or colitis or in those receiving concomitant therapy with an aminoglycoside antibiotic.

5.3 Nephrotoxicity

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) has occurred following oral Vancomycin Hydrochloride Capsules therapy in randomized controlled clinical studies, and can occur either during or after completion of therapy.

The risk of nephrotoxicity is increased in patients >65 years of age [see *ADVERSE REACTIONS, Clinical Trial Experience (6.1) and USE IN SPECIFIC POPULATIONS, Geriatric Use (8.5)*].

In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with Vancomycin Hydrochloride Capsules to detect potential vancomycin induced nephrotoxicity.

5.4 Ototoxicity

Ototoxicity has occurred in patients receiving vancomycin. It may be transient or permanent. It has been reported mostly in patients who have been given excessive intravenous doses, who have an underlying hearing loss, or who are receiving concomitant therapy with another ototoxic agent, such as an aminoglycoside. Serial tests of auditory function may be helpful in order to minimize the risk of ototoxicity [see *ADVERSE REACTIONS, Postmarketing Experience (6.2)*].

5.5 Severe Dermatologic Reactions

Severe dermatologic reactions such as toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP), and

linear IgA bullous dermatosis (LABD) have been reported in association with the use of vancomycin. Cutaneous signs or symptoms reported include skin rashes, mucosal lesions, and blisters.

Discontinue Vancomycin Hydrochloride Capsules at the first appearance of signs and symptoms of TEN, SJS, DRESS, AGEP, or LABD.

5.6 Development of Drug-Resistant Bacteria

Prescribing Vancomycin Hydrochloride Capsules in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria.

5.7 Risk of Allergic Reactions due to Tartrazine

The 125 mg strength capsules contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure to vancomycin hydrochloride in 260 adult subjects in two Phase 3 clinical trials for the treatment of diarrhea associated with *C. difficile*. In both trials, subjects received vancomycin hydrochloride capsules 125 mg orally four times daily. The mean duration of treatment was 9.4 days. The median age of patients was 67, ranging between 19 and 96 years of age. Patients were predominantly Caucasian (93%) and 52% were male.

Adverse reactions occurring in $\geq 5\%$ of vancomycin hydrochloride-treated subjects are shown in Table 1. The most common adverse reactions associated with vancomycin hydrochloride ($\geq 10\%$) were nausea, abdominal pain, and hypokalemia.

Table 1: Common ($\geq 5\%$) Adverse Reactions for Vancomycin Hydrochloride Reported in Clinical Trials for Treatment of Diarrhea Associated with *C. difficile*

System/Organ Class	Adverse Reaction	VancomycinHydrochloride% (N=260)
Gastrointestinal disorders	Nausea	17
	Abdominal pain	15
	Vomiting	9
	Diarrhea	9
	Flatulence	8
General disorders and administration site conditions	Pyrexia	9
	Edema peripheral	6
	Fatigue	5
Infections and infestations	Urinary tract infection	8
Metabolism and nutrition disorders	Hypokalemia	13
Musculoskeletal and connective tissue disorders	Back pain	6
Nervous system disorders	Headache	7

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) occurred in 5% of subjects treated with vancomycin hydrochloride. Nephrotoxicity following vancomycin hydrochloride typically first occurred within one week after completion of treatment (median day of onset was Day 16). Nephrotoxicity following vancomycin hydrochloride occurred in 6% of subjects >65 years of age and 3% of subjects ≤ 65 years of age [see **WARNINGS AND PRECAUTIONS, Nephrotoxicity (5.3)**].

The incidences of hypokalemia, urinary tract infection, peripheral edema, insomnia, constipation, anemia, depression, vomiting, and hypotension were higher among subjects >65 years of age than in subjects ≤65 years of age [see *USE IN SPECIFIC POPULATIONS, Geriatric Use (8.5)*].

Discontinuation of study drug due to adverse events occurred in 7% of subjects treated with vancomycin hydrochloride. The most common adverse events leading to discontinuation of vancomycin hydrochloride were *C. difficile* colitis (<1%), nausea (<1%), and vomiting (<1%).

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of vancomycin hydrochloride. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Ototoxicity: Cases of hearing loss associated with intravenously administered vancomycin have been reported. Most of these patients had kidney dysfunction or a preexisting hearing loss or were receiving concomitant treatment with an ototoxic drug [see *WARNINGS AND PRECAUTIONS, Ototoxicity (5.4)*]. Vertigo, dizziness, and tinnitus have been reported.

Skin and Subcutaneous Tissue Disorders: Severe dermatologic reactions such as toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalized exanthematous pustulosis (AGEP), and linear IgA bullous dermatosis (LABD)

[see *Warnings and Precautions (5.5)*], rashes (including exfoliative dermatitis).

Hematopoietic: Reversible neutropenia, usually starting 1 week or more after onset of intravenous therapy with vancomycin or after a total dose of more than 25 g, has been reported for several dozen patients. Neutropenia appears to be promptly reversible when vancomycin is discontinued. Thrombocytopenia has been reported.

Miscellaneous: Patients have been reported to have had anaphylaxis, drug fever, chills, nausea, eosinophilia, and cases of vasculitis in association with the administration of vancomycin.

A condition has been reported that is similar to the IV-induced syndrome with symptoms consistent with anaphylactoid reactions, including hypotension, wheezing, dyspnea, urticaria, pruritus, flushing of the upper body ("Red Man Syndrome"), pain and muscle spasm of the chest and back. These reactions usually resolve within 20 minutes but may persist for several hours.

7 DRUG INTERACTIONS

No drug interaction studies have been conducted.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Systemic absorption of vancomycin is low following oral administration of Vancomycin Hydrochloride Capsules; however, absorption may vary depending on various factors [see Clinical Pharmacology (12.3)]. There are no available data

on vancomycin use in pregnant women to assess a risk of major birth defects or miscarriage. Available published data on intravenous vancomycin use in pregnancy during the second and third trimesters have not shown an association with adverse maternal or fetal outcomes (see Data).

Vancomycin did not show adverse developmental effects when administered intravenously to pregnant rats and rabbits during organogenesis at doses less than or equal to the recommended maximum human dose (see Data).

Data

Human Data

There are no available data on first trimester use of vancomycin in pregnant women to assess a risk of major birth defects or miscarriage.

A published study evaluated hearing loss and nephrotoxicity in infants of 10 pregnant intravenous drug users treated with intravenous vancomycin for suspected or documented methicillin-resistant *Staphylococcus aureus* in the second or third trimester. The comparison groups were 10 uninfected non-intravenous drug-dependent patients who received no treatment and 10 uninfected untreated intravenous drug-dependent patients. No infant in the vancomycin exposed group had abnormal sensorineural hearing at 3 months of age or nephrotoxicity.

A published prospective study assessed outcomes in 55 pregnant women with a positive Group B streptococcus culture and a high-risk penicillin allergy with resistance to clindamycin or unknown sensitivity who were administered intravenous vancomycin at the time of delivery. Vancomycin dosing ranged from the standard dose of 1 g intravenously every 12 hours to a dose of 20 mg/kg intravenously every 8 hours (maximum individual dose 2 g). No major adverse reactions were recorded either in the mothers or their newborns. None of the newborns had sensorineural hearing loss. Neonatal renal function was not examined, but all of the newborns were discharged in good condition.

Animal Data

Vancomycin did not cause fetal malformation when administered intravenously during organogenesis to pregnant rats (gestation days 6 to 15) and rabbits (gestation days 6 to 18) at the equivalent recommended maximum human dose of 200 mg/kg/day to rats or 120 mg/kg/day to rabbits. No effects on fetal weight or development were seen in rats at the highest dose tested or in rabbits given 80 mg/kg/day (approximately 1 and 0.8 times the recommended maximum human dose based on body surface area). Maternal toxicity was observed in rats (at doses 120 mg/kg and above) and rabbits (at 80 mg/kg and above).

8.2 Lactation

Risk Summary

There are no data on the presence of vancomycin in human milk, the effects on the

breastfed infant, or the effect on milk production following oral administration. Systemic absorption of vancomycin is low following oral administration of Vancomycin Hydrochloride Capsules [see Clinical Pharmacology (12.3)]; therefore, it is unlikely to result in clinically relevant exposure in breastfeeding infants. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Vancomycin Hydrochloride Capsules and any potential adverse effects on the breastfed infant from Vancomycin Hydrochloride

Capsules or from the underlying maternal condition.

8.4 Pediatric Use

Vancomycin Hydrochloride Capsules are indicated in pediatric patients less than 18 years of age for the treatment of *C. difficile* associated diarrhea and enterocolitis caused by *S. aureus* (including methicillin-resistant strains) [see Indications and Usage (1) and Dosage and Administration (2.2)].

8.5 Geriatric Use

In clinical trials, 54% of vancomycin hydrochloride-treated subjects were >65 years of age. Of these, 40% were between the ages of >65 and 75, and 60% were >75 years of age.

Clinical studies with vancomycin hydrochloride in diarrhea associated with *Clostridium difficile* have demonstrated that geriatric subjects are at increased risk of developing nephrotoxicity following treatment with oral vancomycin hydrochloride, which may occur during or after completion of therapy. In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with vancomycin hydrochloride to detect potential vancomycin induced nephrotoxicity [see **WARNINGS AND PRECAUTIONS, Nephrotoxicity (5.3)**; **ADVERSE REACTIONS, Clinical Trials Experience (6.1)** and **CLINICAL STUDIES, Diarrhea Associated with *Clostridium difficile* (14.1)**].

Patients >65 years of age may take longer to respond to therapy compared to patients ≤65 years of age [see **CLINICAL STUDIES, Diarrhea Associated with *Clostridium difficile* (14.1)**]. Clinicians should be aware of the importance of appropriate duration of vancomycin hydrochloride treatment in patients >65 years of age and not discontinue or switch to alternative treatment prematurely.

10 OVERDOSAGE

Supportive care is advised, with maintenance of glomerular filtration. Vancomycin is poorly removed by dialysis. Hemofiltration and hemoperfusion with polysulfone resin have been reported to result in increased vancomycin clearance.

To obtain current information about the treatment of overdose, contact a certified Poison Control Center (1-800-222-1222 or www.poisson.org). In managing overdose, consider the possibility of multiple drug overdoses, interaction among drugs, and unusual drug kinetics.

11 DESCRIPTION

and, in the presence of renal impairment, the possibility of accumulation exists. It should be noted that the total systemic and renal clearances of vancomycin are reduced in the elderly [see *USE IN SPECIFIC POPULATIONS (8.5)*].

12.4 Microbiology

Mechanism of Action

The bactericidal action of vancomycin against *Staphylococcus aureus* and the vegetative cells of *Clostridium difficile* results primarily from inhibition of cell-wall biosynthesis.

In addition, vancomycin alters bacterial-cell-membrane permeability and RNA synthesis.

Mechanism of resistance

Staphylococcus aureus:

S. aureus isolates with vancomycin minimal inhibitory concentrations (MICs) as high as 1024 mcg/mL have been reported.

The exact mechanism of this resistance is not clear but is believed to be due to cell wall thickening and potentially the transfer of genetic material.

Clostridium difficile: Isolates of *C. difficile* generally have vancomycin MICs of <1 mcg/mL, however vancomycin MICs ranging from 4 mcg/mL to 16 mcg/mL have been reported. The mechanism which mediates *C. difficile*'s decreased susceptibility to vancomycin has not been fully elucidated.

Vancomycin has been shown to be active against most isolates of the following microorganisms, both in vitro and in clinical infections [see Indications and Usage (1)].

Gram-positive bacteria

Staphylococcus aureus (including methicillin-resistant isolates) associated with enterocolitis

Anaerobic Gram-positive bacteria

Clostridium difficile isolates associated with *C. difficile* associated diarrhea.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term carcinogenesis studies in animals have been conducted.

At concentrations up to 1000 mcg/mL, vancomycin had no mutagenic effect *in vitro* in the mouse lymphoma forward mutation assay or the primary rat hepatocyte unscheduled DNA synthesis assay. The concentrations tested *in vitro* were above the peak plasma vancomycin concentrations of 20 to 40 mcg/mL usually achieved in humans after slow infusion of the maximum recommended dose of 1 g. Vancomycin had no mutagenic effect *in vivo* in the Chinese hamster sister chromatid exchange assay (400 mg/kg IP) or the mouse micronucleus assay (800 mg/kg IP).

No definitive fertility studies have been conducted.

14 CLINICAL STUDIES

14.1 Diarrhea Associated With *Clostridium Difficile*

In two trials, Vancomycin Hydrochloride Capsules 125 mg orally four times daily for 10 days was evaluated in 266 adult subjects with *C. difficile* -associated diarrhea (CDAD). Enrolled subjects were 18 years of age or older and received no more than 48 hours of treatment with oral vancomycin hydrochloride or oral/intravenous metronidazole in the 5 days preceding enrollment. CDAD was defined as ≥ 3 loose or watery bowel movements within the 24 hours preceding enrollment, and the presence of either *C. difficile* toxin A or B, or pseudomembranes on endoscopy within the 72 hours preceding enrollment. Subjects with fulminant *C. difficile* disease, sepsis with hypotension, ileus, peritoneal signs or severe hepatic disease were excluded.

Efficacy analyses were performed on the Full Analysis Set (FAS), which included randomized subjects who received at least one dose of vancomycin hydrochloride and had any post-dosing investigator evaluation data (N=259; 134 in Trial 1 and 125 in Trial 2).

The demographic profile and baseline CDAD characteristics of enrolled subjects were similar in the two trials. Vancomycin Hydrochloride -treated subjects had a median age of 67 years, were mainly white (93%), and male (52%). CDAD was classified as severe (defined as 10 or more unformed bowel movements per day or $WBC \geq 15000/mm^3$) in 25% of subjects, and 47% were previously treated for CDAD.

Efficacy was assessed by using clinical success, defined as diarrhea resolution and the absence of severe abdominal discomfort due to CDAD, on Day 10. An additional efficacy endpoint was the time to resolution of diarrhea, defined as the beginning of diarrhea resolution that was sustained through the end of the prescribed active treatment period.

The results for clinical success for vancomycin hydrochloride -treated subjects in both trials are shown in Table 2.

Table 2: Clinical Success Rates (Full Analysis Set)

	Clinical Success Rate	95% Confidence Interval
	Vancomycin Hydrochloride% (N)	
Trial 1	81.3 (134)	(74.4, 88.3)
Trial 2	80.8 (125)	(73.5, 88.1)

The median time to resolution of diarrhea was 5 days and 4 days in Trial 1 and Trial 2, respectively. For subjects older than 65 years of age, the median time to resolution was 6 days and 4 days in Trial 1 and Trial 2, respectively. In subjects with diarrhea resolution at end-of-treatment with Vancomycin Hydrochloride Capsules, recurrence of CDAD during the following four weeks occurred in 25 of 107 (23%) and 18 of 102 (18%) in Trial 1 and Trial 2, respectively.

Restriction Endonuclease Analysis (REA) was used to identify *C. difficile* baseline isolates in the BI group. In Trial 1, the Vancomycin Hydrochloride-treated subjects were classified at baseline as follows 31 (23%) with BI strain, 69 (52%) with non-BI strain, and 34 (25%) with unknown strain. Clinical success rates were 87% for BI strain, 81% for non-BI strain, and 76% for unknown strain. In subjects with diarrhea resolution at end-of

treatment with vancomycin hydrochloride, recurrence of CDAD during the following four weeks occurred in 7 of 26 subjects with BI strain, 12 of 56 subjects with non-BI strain, and 6 of 25 subjects with unknown strain.

16 HOW SUPPLIED/STORAGE AND HANDLING

Vancomycin Hydrochloride Capsules, USP are available in:

The 125 mg* CAPSULES have a blue cap and yellow body imprinted with “OP64” on both cap and body in white ink.

The 250 mg* CAPSULES have a blue cap and dark-yellow body imprinted with “OP65” on both cap and body in white ink.

Strength	Pack	NDC Number
Vancomycin 125 mg	A carton contains 2 blister packs. Each blister pack contains 10 capsules for a total of 20 capsules per carton.	NDC No. 76333-264-26
	Each bottle contains 50 capsules.	NDC No. 76333-264-17
Vancomycin 250 mg	A carton contains 2 blister packs. Each blister pack contains 10 capsules for a total of 20 capsules per carton.	NDC No. 76333-265-26
	Each bottle contains 50 capsules.	NDC No. 76333-265-17

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F).
[See USP Controlled Room Temperature]

*Equivalent to vancomycin.

15 REFERENCES

1. Byrd RA., Gries CL, Buening M.: Developmental Toxicology Studies of Vancomycin Hydrochloride Administered Intravenously to Rats and Rabbits. *Fundam Appl Toxicol* 1994; 23: 590-597.

17 PATIENT COUNSELING INFORMATION

Severe Dermatologic Reactions

Advise patients about the signs and symptoms of serious skin manifestations. Instruct patients to stop taking Vancomycin Hydrochloride Capsules immediately and promptly seek medical attention at the first signs or symptoms of skin rash,

mucosal lesions or blisters [see Warnings and Precautions (5.5)].

Antibacterial Resistance

Patients should be counseled that antibacterial drugs including Vancomycin Hydrochloride should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Vancomycin Hydrochloride is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Vancomycin Hydrochloride or other antibacterial drugs in the future.

Manufactured and Distributed by:

Orient Pharma Co., Ltd.

8 Kehu 1st Road, Huwei Chen, Yunlin, Taiwan 63247

Product of Taiwan

Rx only

To request additional information or if you have questions concerning vancomycin hydrochloride capsules please phone Orient Pharma Co., Ltd. at +886-5-631-1331.

PACKAGE/LABEL DISPLAY PANEL

**Vancomycin Hydrochloride Capsules USP, Equiv. to 125 mg vancomycinNDC
76333-264-26Rx Only 20 Capsules**

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-264-26

Rx Only

125 mg (base)

Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

EXP.: XX/XXXX
LOT: XXXXXXXX

PEEL TO OPEN

*Equivalent to 125 mg vancomycin

125 mg *

Vancomycin Hydrochloride
Capsules, USP

NDC-76333-264-26

NDC-76333-264-26

Vancomycin Hydrochloride
Capsules, USP

125 mg*

*Equivalent to 125 mg vancomycin

20 Capsules (2 x 10 units-dose)
Not a Child Resistant Container

Rx only

 orient PHARMA co.,Ltd.

NDC-76333-264-26

Vancomycin Hydrochloride
Capsules, USP

125 mg *

*Equivalent to 125 mg vancomycin

NDC-76333-264-26

Vancomycin Hydrochloride
Capsules, USP

125 mg *

*Equivalent to 125 mg vancomycin

LOT:
EXP:

*Each Capsule Contains: vancomycin hydrochloride equivalent to 125 mg vancomycin

Usual Total Daily Adult Dose: 500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections. See package insert for other dosage information.

Storage: Store at Controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]

Contains color additives including FD&C Yellow No. 5 (tartrazine). Unit Dose Medication

Barcode

Manufactured and Distributed by: Orient Pharma Co., Ltd.
8 Kehu 1st Rd., Huwei Chen, Yunlin 63247, Taiwan


Vancomycin Hydrochloride Capsules USP, Equiv. to 250 mg vancomycin NDC 76333-265-26 Rx Only 20 Capsules

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan


PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

**Vancomycin Hydrochloride
Capsule, USP**

NDC-76333-265-26

Rx Only

250 mg (base)



Bar code

Mfg. by: Orient Pharma, Co., Ltd.
Made in Taiwan

PEEL TO OPEN
LOT: XXXXXXX
EXP.: XX/XXXX

***Each Capsule Contains:** vancomycin hydrochloride equivalent to 250 mg vancomycin

Usual Total Daily Adult Dose: 500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections. See package insert for other dosage information.

Storage: Store at Controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]

Unit Dose Medication



Manufactured and Distributed by: Orient Pharma Co., Ltd.
8 Kehu 1st Rd., Huwei Chen, Yunlin 63247, Taiwan

LOT:
EXP:

NDC-76333-265-26
Vancomycin Hydrochloride
Capsules, USP
250 mg *
*Equivalent to 250 mg vancomycin

NDC-76333-265-26
Vancomycin Hydrochloride
Capsules, USP
250 mg *
*Equivalent to 250 mg vancomycin
20 Capsules (2 x 10 units-dose)
Not a Child Resistant Container
Rx only
orient PHARMA CO., Ltd.

NDC-76333-265-26
Vancomycin Hydrochloride
Capsules, USP
250 mg *
*Equivalent to 250 mg vancomycin

NDC-76333-265-26
Vancomycin Hydrochloride
Capsules, USP
250 mg *
*Equivalent to 250 mg vancomycin

Vancomycin Hydrochloride Capsules USP, Equiv. to 125 mg vancomycin NDC 76333-264-17 Rx Only 50 Capsules

Each Capsule Contains:

Vancomycin hydrochloride USP equivalent to 125 mg vancomycin.

Usual Total Daily Adult Dose:

500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections. See Package insert for other dosage information.

Storage:

Store at controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]

NDC 76333-264-17

Vancomycin Hydrochloride Capsules, USP

125 mg*

* Equivalent to 125 mg vancomycin

Rx only
50 Capsules

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Contains color additives including FD&C Yellow No. 5 (tartrazine)

Manufactured and Distributed by:
Orient Pharma Co., Ltd.
8 Kehu 1st Road, Huwei Chen, Yunlin 63247, Taiwan

Barcode

Vancomycin Hydrochloride Capsules USP, Equiv. to 250 mg vancomycin NDC 76333-265-17 Rx Only 50 Capsules

Each Capsule Contains:

Vancomycin hydrochloride USP equivalent to 250 mg vancomycin.

Usual Total Daily Adult Dose:

500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections. See Package insert for other dosage information.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

NDC 76333-265-17

Vancomycin Hydrochloride Capsules, USP

250 mg*

* Equivalent to 250 mg vancomycin

Rx only
50 Capsules

Storage:

Store at controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]

Manufactured and Distributed by:
Orient Pharma Co., Ltd.
8 Kehu 1st Road, Huwei Chen, Yunlin 63247, Taiwan

Barcode

VANCOMYCIN HYDROCHLORIDE

vancomycin hydrochloride capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76333-264
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of	Strength
-----------------	----------	----------

Ingredient Name	Strength	Strength
VANCOMYCIN HYDROCHLORIDE (UNII: 71WO621TJD) (VANCOMYCIN - UNII:6Q205EH1VU)	VANCOMYCIN	125 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
GELATIN (UNII: 2G86QN327L)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SHELLAC (UNII: 46N107B71O)	
GLYCERYL MONOOLEATE (UNII: C4YAD5F5G6)	

Product Characteristics

Color	blue (Blue Cap) , yellow (Yellow Body)	Score	no score
Shape	CAPSULE (Capsule)	Size	18mm
Flavor		Imprint Code	OP;64
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76333-264-01	2 in 1 CARTON	07/25/2019	
1	NDC:76333-264-26	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:76333-264-17	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210729	07/25/2019	

VANCOMYCIN HYDROCHLORIDE

vancomycin hydrochloride capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76333-265
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VANCOMYCIN HYDROCHLORIDE (UNII: 71WO621TJD) (VANCOMYCIN - UNII:6Q205EH1VU)	VANCOMYCIN	250 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SHELLAC (UNII: 46N107B71O)	
GLYCERYL MONOOLEATE (UNII: C4YAD5F5G6)	

Product Characteristics

Color	blue (Blue Cap) , yellow (Dark-Yellow Body)	Score	no score
Shape	CAPSULE (Capsule)	Size	22mm
Flavor		Imprint Code	OP;65
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76333-265-01	2 in 1 CARTON	07/25/2019	
1	NDC:76333-265-26	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:76333-265-17	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210729	07/25/2019	

Labeler - Orient Pharma Co.,Ltd Yunlin Plant (658849810)**Registrant** - Orient Pharma Co.,Ltd Yunlin Plant (658849810)

Establishment

Name	Address	ID/FEI	Business Operations
Orient Pharma Co.,Ltd		658849810	manufacture(76333-264, 76333-265)

Revised: 6/2022

Orient Pharma Co.,Ltd Yunlin Plant