

**ACID REDUCER - omeprazole tablet, delayed release**  
**Family Dollar Stores, LLC**

-----

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**Drug Facts**

**Active ingredient (in each tablet)**

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium USP)

**Purpose**

Acid reducer

**Use**

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:**

- Do not use if you are allergic to omeprazole.
- Omeprazole may cause severe skin reactions. Symptoms may include:
  - skin reddening
  - blisters
  - rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**.

These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day

- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew or crush tablets.**

### **Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

### **Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77° F) and protect from moisture

### **Inactive ingredients**

crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid and ethyl acrylate copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide.

### **Questions?**

Call **1-855-274-4122**

### **DISTRIBUTED BY:**

MIDWOOD BRANDS, LLC  
500 VOLVO PKWY  
CHESAPEAKE, VA 23320 USA

**MADE IN INDIA**

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablets Container Label)**

### **FAMILY**

**Wellness™**

**See Current Drug Facts**

### **OMEPRAZOLE**

**Delayed-Release Tablets, 20 mg  
Acid Reducer**

**24 HOUR**

**Treats Frequent Heartburn!**

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT  
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

NDC 55319-037-05

**FAMILY Wellness** See Current Drug Facts

**OMEPRAZOLE**   
Delayed-Release Tablets, 20 mg  
Acid Reducer  
Treats Frequent Heartburn

**14 TABLETS**  
ONE 14-DAY COURSE OF TREATMENT  
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT NDC 55319-037-05

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

**TAMPER EVIDENT - Do not use if seal over bottle opening is broken or missing.**

DISTRIBUTED BY:  
MIDWOOD BRANDS, LLC  
500 VOLVO PKWY  
CHESAPEAKE, VA 23320 USA  
MADE IN INDIA

P1433449 LM-5448

Lift Here

No Varnish

Top Ply  
(Page #1)

**Active ingredient (in each tablet)**  
Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium USP)

**Purpose**  
Acid reducer

**Use** ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings** ■ rash. If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you

P1433449 LM-5448

**HINGE**

Back of Top Ply  
(Page #2)

Bottom Ply

**HINGE**

develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours **14-Day Course of Treatment** ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. **Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. **Other information** ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) and protect from moisture **Inactive ingredients** crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid and ethyl acrylate copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide. **Questions?** Call 1-855-274-4122

P1433449 LM-5448

---

**Base  
(Page #3)  
Bottom Ply**

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablets Container  
Carton Label)**

**FAMILY**

**wellness™**

**See Current Drug Facts**

**COMPARE TO**

**PRILOSEC OTC®**

**ACTIVE INGREDIENT\***

**OMEPRAZOLE**

**Delayed-Release**

**Tablets, 20 mg**

**Treats Frequent Heartburn!**

**24 HOUR**

**14 TABLETS**

ACTUAL SIZE

**ONE 14-DAY COURSE OF TREATMENT**

**MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT**

**NDC 55319-037-05**



## ACID REDUCER

omeprazole tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-037
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OMEPRAZOLE MAGNESIUM</b> (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSPROVIDONE (35 .MU.M)</b> (UNII: 40UAA97IT9)	
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	
<b>HYDROXYPROPYL CELLULOSE (90000 WAMW)</b> (UNII: UKE75GEA7F)	
<b>HYPROMELLOSE 2910 (5 MPA.S)</b> (UNII: R75537T0T4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A</b> (UNII: NX76LV5T8J)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>SODIUM STEARYL FUMARATE</b> (UNII: 7CV7WJK4UI)	
<b>DEXTROSE, UNSPECIFIED FORM</b> (UNII: IY9XDZ35W2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>HYDROXYPROPYL CELLULOSE (45000 WAMW)</b> (UNII: 8VAB711C5E)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	RECTANGLE (Oblong)	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	Z;69
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-037-05	1 in 1 CARTON	06/17/2024	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55319-037-62	3 in 1 CARTON	06/17/2024	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206877	06/17/2024	

**Labeler** - Family Dollar Stores, LLC (024472631)

**Registrant** - Aurohealth LLC (078728447)

**Establishment**

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(55319-037) , MANUFACTURE(55319-037)

Revised: 6/2024

Family Dollar Stores, LLC