

**ASPIRIN ADULT LOW DOSE- acetylsalicylic acid tablet, delayed release**  
**Medline Industries, LP**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**985 Aspirin 81mg EC tablets**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

- temporarily relieves minor aches and pains. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- if you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or loss of hearing occurs.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and older: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless otherwise directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 25°C(77°F) excursions permitted between 15°-30°C (59°-68°F)
- use by expiration date on package

**Inactive ingredients**

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Manufacturing Information**

Manufactured by Medline Industries, LP  
Three Lakes Drive, Northfield, IL 60093 USA

Made in the USA with U.S. and Foreign Components

[www.medline.com](http://www.medline.com)

1-800-MEDLINE (633-5463)


REF: OTCM00003

V2 RH22HND

**Package Label**



NDC: 53329-985-67

Adult Low Dose  
**Aspirin**Pain Reliever  
(NSAID)\*\*Compare to the active  
ingredient in Bayer®  
Low Dose**81  
mg****120** ENTERIC  
COATED TABLETS**TAMPER EVIDENT: DO NOT USE IF PRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.****Drug Facts****Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin**Warnings****Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach**REF/OTC M00003 (CONTINUED ON BACK OF LABEL)**

\*This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Low Dose.

www.medline.com Made in the USA with U.S. and Foreign Components. Manufactured for Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093, USA. 1-800-MEDLINE V2-RH24HD

**88277 69599 6**LOT  
EXP

Coating Free

**PEEL  
HERE****Drug Facts (continued)**

ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma**Ask a doctor or pharmacist before use if you are taking a prescription drug for:** ■ gout ■ diabetes ■ arthritis**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs These could be signs of a serious condition.**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.****Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and older: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless otherwise directed by a doctor ■ children under 12 years: consult a doctor**Other information** ■ store at 25°C (77°F) excursions permitted between 15°–30°C (59°–86°F) ■ use by expiration date on package**Inactive ingredients** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate**ASPIRIN ADULT LOW DOSE**

acetylsalicylic acid tablet, delayed release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53329-985
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A</b> (UNII: NX76LV5T8J)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	score with uneven pieces
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	embossed;heart
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-985-67	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/25/2018	

**Labeler** - Medline Industries, LP (025460908)

**Registrant** - Medline Industries, LP (025460908)