

BANDAGE- benzocaine and benzethonium chloride aerosol, spray

HART Health

Spray Bandage

□ Active Ingredients

Benzocaine 10% w/w
Benzethonium Chloride 0.2% w/w

Purpose

Topical Anesthetic
Topical Antiseptic

□ Uses

First aid to help protect against infection in

- minor cuts
- scrapes
- burns

Warnings

For External Use Only

Flammable

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120°F

Do not use

- in or near the eyes or other mucous membranes
- over large portions of the body
- longer than 7 days unless directed by a doctor
- on deep or puncture wounds, raw surfaces, blistered areas, animal bites, or serious burns

Stop use and ask doctor if

- redness, swelling, or irritation occurs
- infection occurs
- condition persists or worsens
- symptoms persist for more than 7 days

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions

Adults and children 12 years of age and over:

- clean the affected area
- shake can well before using
- hold 4 to 6 inches from surface and spray until area is covered with film
- when dry, may be covered with a sterile bandage

Children under 12 years of age: ask a doctor

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive Ingredients

copovidone K35-31, ethyl alcohol, isobutane, n-butane, propane

Principal Display

6.75" x 4"

PMS 321C

Drug Facts

Active Ingredients
Benzocaine 0.2% w/w Topical Anesthetic
Benzocaine 3.2% w/w Topical Anesthetic

Uses
First aid to help protect against infection in
 ■ minor cuts ■ scrapes ■ burns

Warnings For external use only
Flammable ■ keep away from fire or flame ■ contents under pressure
 ■ do not puncture or incinerate container ■ do not expose to temperatures above 120°F

Do not use
 ■ in or near the eyes or other mucous membranes
 ■ over large portions of the body ■ longer than 7 days unless directed by a doctor
 ■ on deep or puncture wounds, raw surfaces, blistered areas, animal bites, or serious burns

Stop use and ask a doctor if ■ redness, swelling or irritation occurs ■ infection occurs
 ■ condition persists or worsens ■ symptoms persist for more than 7 days

Keep out of reach of children.
 If swallowed, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions
 Adults and children 12 years of age and over:
 ■ clean the affected area ■ shake can well before using
 ■ hold can 4 to 6 inches away from injury and spray until area is covered with film
 ■ when dry, may be covered with a sterile bandage
 Children under 12 years of age: ask a doctor

Other Information
 ■ avoid inhaling ■ use only as directed
 ■ intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive Ingredients copovidone K35-3T, ethyl alcohol, isobutane, n-butane, propane

Manufactured for:
HART Health
 Seattle, WA 98124 www.harthealth.com

12-295-21-27
 8 50332-0218 1 7

NDC 50332-0218-1 Item 2765

Spray Bandage
Antiseptic Dressing

Protects cuts and abrasions
 Helps prevent infection

Contains no fluorocarbons
 environmentally safe
 3 oz. (85 gm)



6.75" x 4"

PMS 321C

L2-295-21-278
NDC 50332-0218-1 Item 2765

Drug Facts

Active Ingredients
Benzethonium Chloride 0.2% w/w Topical Anesthetic
Benzocaine 3.2% w/w Topical Anesthetic

Uses
First aid to help protect against infection in

- minor cuts
- scrapes
- burns

Warnings For external use only

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120°F

Do not use

- in or near the eyes or other mucous membranes
- longer than 7 days unless directed by a doctor
- on deep or puncture wounds, raw surfaces, blistered areas, animal bites, or serious burns

Stop use and ask a doctor if

- redness, swelling or irritation occurs
- infection occurs
- condition persists or worsens
- symptoms persist for more than 7 days

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions
Adults and children 12 years of age and over:

- shake can well before using
- clean the affected area
- hold can 4 to 6 inches away from injury and spray until area is covered with film
- when dry, may be covered with a sterile bandage

Other Information
Children under 12 years of age: ask a doctor

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive Ingredients copovidone K35-3T, ethyl alcohol, isobutane, n-butane, propane



Spray Bandage
Antiseptic Dressing

Protects cuts and abrasions
Helps prevent infection

Contains no fluorocarbons environmentally safe
3 oz. (85 gm)



Manufactured for
HART Health
Seattle, WA 98124 www.harthealth.com



12-295-21-278
N 50332-0218-1
8

BANDAGE

benzocaine and benzethonium chloride aerosol, spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50332-0218
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	2720 mg in 85 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

COPOVIDONE K25-31 (UNII: D9C330MD8B)

BUTANE (UNII: 6LV4FOR43R)

PROPANE (UNII: T75W9911L6)

ISOBUTANE (UNII: BXR49TP611)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50332-0218-1	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/01/1974	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M	04/01/1974	

Labeler - HART Health (069560969)

Revised: 1/2024

HART Health