# **DIMENHYDRINATE-** dimenhydrinate tablet CVS Pharmacy, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### 1006-CVS

## **Drug Facts**

## Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Antiemetic

#### Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

**Do not use** for children under 2 years of age unless directed by a doctor.

## Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to an enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

## When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
- adults and children 12 years of age and over: 1 to 2 tablets every 4-6 hours; not to exceed 8 tablets in 24 hours, or as directed by a doctor

- children 6 to under 12 years of age: 1/2 to 1 tablet every 6-8 hours; not to exceed 3 tablets in 24 hours, or as directed by a doctor
- children 2 to under 6 years of age: 1/2 tablet every 6-8 hours; not to exceed 1 1/2 tablets in 24 hours, or as directed by a doctor

#### Other Informations

- each tablet contains: calcium 30 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture

croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose

#### **Questions or comments?**

1-800-231-4670

#### KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

## TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

\* This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine®

## Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2021 CVS/pha rmacy

CVS.com® 1-800-SHOP CVS

V-30287

#### **CVSHealth**

Compare to the active ingredient in Dramamine® Original Formula\*

#### **Motion Sickness**

#### **DIMENHYDRINATE TABLETS**

Antiemetic

Original formula

- Fast acting motion sickness relief
- For children & adults

## 36 TABLETS (50 mg EACH)



## Fast-acting relief of motion sickness





**Original Formula** 

# **Motion Sickness**

**DIMENHYDRINATE TABLETS Antiemetic** 

Original formula

- · Fast-acting motion sickness relief
- For children & adults



Compare to the active ingredient in Dramamine®Original Formula\*

**Original Formula** 

# **Motion Sickness**

DIMENHYDRINATE TABLETS Antiemetic

Original formula

- Fast-acting motion sickness relief
- For children & adults



**Actual Size** 

36 TABLETS (50 mg EACH)

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

\*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine®

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2021 CVS/pharmacy CVS.com® 1-800-SHOP CVS





#### 1-800-231-4670 Chestions or comments?

stearate, microcrystalline cellulose sodium, dicalcium phosphate, magnesium Inactive ingredients croscarmellose

profect from moisture

Detween 15°-30°C (59°-86°F) ■ store at 25°C (77°F); excursions permitted m esch tablet contains: calcium 30 mg

**NAME OF THE PROPERTY OF THE P** 

ponts, or as directed by a doctor every 6-8 hours; not to exceed 1 1/2 tablets in 24 ■ children 2 to under 6 years of age: 1/2 tablet

24 hours, or as directed by a doctor tablet every 6-8 hours; not to exceed 3 tablets in ■ children 6 to under 12 years of age: 1/2 to 1 COCCOL

exceed 8 tablets in 24 hours, or as directed by a over: 1 to 2 tablets every 4-6 hours; not to m adults and children 12 years of age and to prevent or treat motion sickness, see below: starting activity

> should be taken one-half to one hour before ■ to prevent motion sickness, the first dose Directions

> > Drug Facts (continued)

(1-800-222-1222) right away. get medical help or contact a Poison Control Center keep out of reach of children, in case of overdose, professional before use.

If pregnant or breast-feeding, ask a health

operating machinery

use caution when driving a motor vehide or drowsiness

alcohol, sedatives, and tranquilizers may increase

svoid alcoholic drinks

warked drowsiness may occur

When using this product

taking sedatives or tranquilizers.

Ask a doctor or pharmacist before use if you are

■ trouble uninating due to an enlarged prostate

chronic bronchitis

a breathing problem such as emphysema or 🔳 địa ncoma

Ask a doctor before use if you have

directed by a doctor.

Do not use for children under 2 years of age unless **SprimsW** 

ssaujzzib 🔳 gaitimov 🖿 sasusna 🖿

symptoms associated with motion sickness: USG for prevention and treatment of these

Dimenhydrinate 50 mg ..... olfemelfinA.

Active ingredient (in each tablet) Purpose

Story burd

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION



## DIMENHYDRINATE

dimenhydrinate tablet

## **Product Information**

**Product Type HUMAN OTC DRUG** Item Code (Source) NDC:69842-206

**Route of Administration ORAL** 

## **Active Ingredient/Active Moiety**

**Ingredient Name Basis of Strength Strength** 

DIMENHYDRINATE (UNII: JB937PER5C) (CHLORTHEOPHYLLINE - UNII:GE2UA340FM) DIMENHYDRINATE 50 mg

## **Inactive Ingredients**

Ingredient Name	Strength
<b>y</b>	

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)

CALCIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: O7TSZ97GEP)

MAGNESIUM STEARATE (UNII: 70097M6I30)

Product Characteristics						
Color	white	Score	2 pieces			
Shape	ROUND	Size	9mm			
Flavor		Imprint Code	1006;1006			
Contains						

P	ackaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:69842- 206-12	1 in 1 CARTON	07/03/2018		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:69842- 206-36	3 in 1 CARTON	07/03/2018		
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part336	07/03/2018			

## Labeler - CVS Pharmacy, Inc. (062312574)

Revised: 11/2021 CVS Pharmacy, Inc.