

DUAL ACTION COMPLETE- famotidine, calcium carbonate and magnesium hydroxide tablet, chewable
Meijer Distribution Inc

Meijer Distribution, Inc. Dual Action Complete Drug Facts

Active ingredients (in each chewable tablet)

Famotidine 10 mg

Calcium carbonate 800 mg

Magnesium hydroxide 165 mg

Purposes

Acid reducer

Antacid

Use

relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- **do not swallow tablet whole: chew completely**
- to relieve symptoms, **chew** 1 tablet before swallowing
- do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** calcium 330 mg; magnesium 80 mg
- Phenylketonurics: Contains Phenylalanine 2 mg per tablet
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20-25°C (68-77°F)
- protect from moisture

Inactive ingredients

anhydrous lactose, artificial berry flavor, aspartame, D&C red no. 7 calcium lake, dextrans, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyacrylate dispersion, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, talc

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Pepcid Complete® active ingredients
 dual action complete
 Famotidine 10 mg | Calcium Carbonate 800 mg
 Magnesium Hydroxide 165 mg Tablets (Chewable)
 Acid Reducer + Antacid
 SEE NEW WARNINGS
 HEARTBURN RELIEF | Just One Tablet!
 Relieves Heartburn Due to Acid Indigestion
 50 Chewable Tablets | actual size
 Berry Flavor

Drug Facts DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each chewable tablet)

Famotidine 10 mg	Acid reducer
Calcium carbonate 800 mg	Antacid
Magnesium hydroxide 165 mg	Antacid

Purposes

Use relieves heartburn associated with acid indigestion and sour stomach

Warnings
 Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- unexplained weight loss
- stomach pain
- frequent wheezing, particularly with heartburn
- nausea or vomiting
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - do not swallow tablet whole; chew completely
 - to relieve symptoms, chew 1 tablet before swallowing
 - do not use more than 2 chewable tablets in 24 hours
- children under 12 years; ask a doctor

Other information

- each tablet contains: calcium 330 mg; magnesium 80 mg
- Phenylketonurics: Contains Phenylalanine 2 mg per tablet
- read the directions and warnings before use
- store at 20-25°C (68-77°F)
- protect from moisture

Drug Facts (continued)

Questions or comments? 1-800-7-19-92-60

Questions or comments?

www.meijer.com

GRAND RAPIDS, MI

DIST. BY ELDER DISTRIBUTION, INC.

U.S.P. No. D 801, 175

Cap or seal is not broken or other tampering is not evident upon opening.

© 2017 Meijer

1 22173 6E F 4

DUAL ACTION COMPLETE			
famotidine, calcium carbonate and magnesium hydroxide tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-284
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	165 mg
CALCIUM CARBONATE (UNII: H0G9379FGK) (CARBONATE ION - UNII:7UJQ5OPE7D, CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CARBONATE	800 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
ASPARTAME (UNII: Z0H242BBR1)	
D&C RED NO. 7 (UNII: ECW0LZ41X8)	
DEXTRATES (UNII: G263MI44RU)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	PINK (mottled)	Score	no score
Shape	ROUND (bi-layered)	Size	17mm
Flavor	BERRY	Imprint Code	L321
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-284-71	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077355	01/26/2016	

Labeler - Meijer Distribution Inc (006959555)

