

VICKS NYQUIL SEVERE COLD AND FLU- acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled
The Procter & Gamble Manufacturing Company

VICKS[®] NyQuil[™] Severe COLD & FLU Liquicaps

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks

- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed • do not exceed 8 LiquiCaps per 24 hrs

| | |
|---------------------------------|------------------------------------|
| adults & children 12 yrs & over | 2 LiquiCaps with water every 4 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- do not exceed 25°C

Inactive ingredients

FD&C Blue No. 1, gelatin, glycerin, pharmaceutical ink*, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *May contain this ingredient

Questions?

1-800-362-1683

TAMPER EVIDENT: This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

Made in Canada

**DIST. BY
PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 24 LiquiCap Carton

MAX

STRENGTH

VICKS®

NyQuil™

SEVERE

COLD & FLU

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains
Nasal Congestion, Sinus Pressure
Sneezing, Runny Nose
Cough

Nighttime Relief

24 LIQUICAPS™



Drug Facts (continued)
Inactive ingredients FD&C Blue No. 1, gelatin, glycerin, pharmaceutical ink*, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitan solution
*May contain this ingredient.
Questions? 1-800-362-1683

Directions
• take only as directed
• do not exceed 8 LiquiCaps per 24 hrs
adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use
Other information
• store at no greater than 25°C

MAX STRENGTH
VICKS®
NyQuil™
SEVERE
COLD & FLU
Acetaminophen, Phenylephrine HCl,
Doxylamine Succinate, Dextromethorphan HBr
Headache, Fever, Sore Throat,
Minor Aches & Pains
Nasal Congestion, Sinus Pressure
Sneezing, Runny Nose
Cough
24 LIQUICAPS™
Nighttime Relief

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

TAMPER EVIDENT:
This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

Made in Canada
DISTR. BY
PROCTER & GAMBLE,
CINCINNATI OH 45202

P&G
www.pg.com
www.vicks.com
Patents:
www.pg.com/patents
90760628

♻️
COATED PAPER/MULTI-LAYER
BOX TRAY
♻️
100% RECYCLED
INK
PRINTED IN U.S.A.



| Drug Facts | |
|---------------------------------------|-----------------------------|
| Active ingredients (in each LiquiCap) | Purpose |
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg | Cough suppressant |
| Doxylamine succinate 6.25 mg | Antihistamine |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure
• cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains
• headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages
• temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage

Warnings
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
• with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure
• thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus)
• a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
• trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers
• taking the blood thinning drug warfarin

When using this product • do not use more than directed • excitability may occur, especially in children
• marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery
• alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if • you get nervous, dizzy or sleepless
• pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days
• redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ▶



VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:37000-518 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE (UNII: FZ989GH94E) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | green | Score | no score |
| Shape | BULLET | Size | 16mm |
| Flavor | | Imprint Code | NS |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:37000-518-24 | 2 in 1 CARTON | 07/10/2018 | |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:37000-518-08 | 1 in 1 CARTON | 07/01/2019 | |
| 3 | | 8 in 1 BLISTER PACK; Type 0: Not a Combination | | |

| | | | | |
|------------------------------|---------------------------|---|-----------------------------|---------------------------|
| 4 | | Product | | |
| 3 | NDC:37000-518-02 | 2 in 1 POUCH; Type 0: Not a Combination Product | 01/01/2021 | |
| Marketing Information | | | | |
| | Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| | OTC Monograph Drug | M012 | 07/10/2018 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 11/2023

The Procter & Gamble Manufacturing Company