

STOMACH RELIEF ULTRA- bismuth subsalicylate liquid
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)

Drug Facts

Active ingredient (in each 15 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
 - 15 mL (1 dose) every 1/2 or 30 mL (2 doses) every hour as needed
diarrhea/traveler's diarrhea
 - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

- **each 15 mL contains:** sodium 5 mg
- **each 15 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

Compare to the active ingredient of Pepto-Bismol® Ultra*

ULTRA

Stomach Relief

Bismuth Subsalicylate 525 mg per 15 mL

Upset Stomach Reliever / Antidiarrheal

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset stomach
- Diarrhea

2x Strength Per Ouncet

Alcohol Free

Sugar Free

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE,

GOODLETTSVILLE, TN 37072

Package Label

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than 8 doses (120 mL) in 24 hours
 ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ mL = milliliter
 ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ children under 12 years of age: ask a doctor

Other information ■ each 15 mL contains: sodium 5 mg
 ■ each 15 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call 1-888-309-9030

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

†vs. Original Pepto-Bismol®

DISTRIBUTED BY OLD EAST MAIN CO.
 100 MISSION RIDGE
 GOODLETTSVILLE, TN 37072



PLD-B494A LB008195 3

DG health

Compare to the active ingredient of Pepto-Bismol® Ultra*

Ultra Stomach Relief

Bismuth Subsalicylate, 525 mg
 Upset Stomach Reliever/
 Antidiarrheal

- Relieves:
- Heartburn
 - Indigestion
 - Nausea
 - Upset stomach
 - Diarrhea

2x strength per ounce†

Alcohol Free
 Sugar Free

8 FL OZ (237 mL)

**Drug Facts**

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 15 mL dose)

Bismuth subsalicylate 525 mg.....Upset stomach reliver/Antidiarrheal

Purpose

Uses relieves

- travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn
- indigestion ■ nausea ■ gas ■ fullness ■ belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have

- fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are

- taking any drug for ■ diabetes ■ gout ■ arthritis
- anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

DOLLAR GENERAL HEALTH Ultra Stomach Relief

STOMACH RELIEF ULTRA

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-837
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-837-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/19/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008		03/19/2020	

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 12/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)