

**DR MEDECEAN RISING SUN COVER- octinoxate, titanium dioxide, octisalate cream
INNOGENE CO.,LTD**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ACTIVE INGREDIENT

Active ingredients: Ethylhexyl Methoxycinnamate 8.0%, Titanium Dioxide 3.0%, Ethylhexyl Salicylate 2.0%

INACTIVE INGREDIENT

Inactive ingredients: Water, Butylene Glycol, Dibutyl Adipate, Diethylamino Hydroxybenzoyl Hexyl Benzoate, Pentylene Glycol, Glycerin, Cyclopentasiloxane, Bis-Ethylhexyloxyphenol Methoxyphenyl Triazine, Cyclohexasiloxane, Dimethicone/Vinyl Dimethicone Crosspolymer, Arachidyl Alcohol, Polysorbate 60, Stearic Acid, Glyceryl Stearate, PEG-100 Stearate, Allantoin, Oleth-10, Behenyl Alcohol, Aluminum Hydroxide, Triethoxycaprylylsilane, Mannitol, Ammonium Acryloyldimethyltaurate/VP Copolymer, Arachidyl Glucoside, 1,2-Hexanediol, Chlorphenesin, Acrylates/C10-30 Alkyl Acrylate Crosspolymer, Ammonium Glycyrrhizate, Triethanolamine, Cocoglucoside, Citric acid, Adenosine, Disodium EDTA, Laureth-4, Caffeine, Zinc Gluconate, BHT, Sorbic Acid, Aesculus Hippocastanum (Horse Chestnut) Seed Extract, Acetyl Tetrapeptide-15

PURPOSE

Purpose: Sunscreen

WARNINGS

Warnings: For external use only


1. If the Red swellings such as erythema, itchiness, or irritations because of directly sunlight occur, you must stop using immediately and Consult your dermatologist.
2. Do not apply on skin conditions such as wounds.
3. Storage and Handling. a. Keep out of the reach of children. b. Avoid direct sunlight.

DESCRIPTION

Uses: Dr.MEDECEAN Rising Sun Cover Cream will protect your sensitive skin from the Sun.

Directions: Apply an appropriate amount to the exposed skin when you need.

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL





주름 기능성 화장품

닥터 메디션 라이징 선 커버 크림 Dr.MEDECEAN Rising SUN cover cream

Dr.메디션은 자외선 차단제 판매로 인한
기업의 이익보다 건강한 피부를 먼저 생각하고
정확한 정보에 대한 고객의 알 권리를 우선합니다.

Drug Facts

Active ingredients :

Ethylhexyl Methoxycinnamate 8.0%

Titanium Dioxide 3.0%

Ethylhexyl Salicylate 2.0%

[Inactive ingredients]

Water, Butylene Glycol, Dibutyl Adipate, Diethylamino Hydr
oxybenzoyl Hexyl Benzoate, Pentylene Glycol, Glycerin, Cycl
opentasiloxane, Bis-Ethylhexyloxyphenol Methoxyphenyl
Triazine, Cyclohexasiloxane, Dimethicone/Vinyl Dimethicone
Crosspolymer, Arachidyl Alcohol, Polysorbate 60, Stearic Acid,
Glyceryl Stearate, PEG-100 Stearate, Allantoin, Oleth-10, Behe
nyl Alcohol, Aluminum Hydroxide, Triethoxycaprylylsilane, M
annitol, Ammonium Acryloyldimethyltaurate/VP Copolymer,
Arachidyl Glucoside, 1,2-Hexanediol, Chlorphenesin, Acrylat
es/C10-30 Alkyl Acrylate Crosspolymer, Ammonium Glycyrh
izate, Triethanolamine, Coco-glucoside, Citric acid, Adenosin
e, Disodium EDTA, Laureth-4, Caffeine, Zinc Gluconate, BHT,
Sorbic Acid, Aesculus Hippocastanum (Horse Chestnut) See
d Extract, Acetyl Tetrapeptide-15

Purpose : Sunscreen

본 제품에 이상이 있을 경우 공정거래위원회 고시 소비자분쟁
해결기준에 의해 교환 또는 보상해 드립니다.

소비자 상담실: 1899-4298

www.medicen.co.kr



8 809405 064370 >

[Directions]

Apply an appropriate
amount to the exposed skin
when you need.

[Warnings]

For external use only

1. If the Red swellings such as
erythema, itchiness, or
irritations because of directly
sunlight occur, you must stop
using immediately and Consult
your dermatologist.

2. Do not apply on skin
conditions such as wounds.

3. Storage and Handling.

a. Keep out of the reach of
children.

b. Avoid direct sunlight.

Manufactured for INNOGENE

Innogene Anyang Factory
Co., Ltd.
(Digital Empire Building A,
B105-B106) 387 Simin-daero,
Dongan-gu Anyang-si,
Gyeonggi-do, Korea

중량 : 50 g

제조번호 및 사용기한 : 별도표시



Dr.MEDECEAN
Rising SUN cover cream



닥터 메디션 라이징 선 커버 크림

Dr.MEDECEAN Rising Sun cover cream
will protect your sensitive skin from the Sun.



DR MEDECEAN RISING SUN COVER

octinoxate, titanium dioxide, octisalate cream

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:70540-040

Route of Administration

TOPICAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|---------------|
| Octinoxate (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51) | Octinoxate | 4.0 g in 50 g |
| Titanium Dioxide (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) | Titanium Dioxide | 1.5 g in 50 g |
| Octisalate (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W) | Octisalate | 1.0 g in 50 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| WATER (UNII: 059QF0K00R) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:70540-040-02 | 1 in 1 CARTON | 09/01/2017 | |
| 1 | NDC:70540-040-01 | 50 g in 1 CONTAINER; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part352 | 09/01/2017 | |

Labeler - INNOGENE CO.,LTD (688201056)**Registrant** - INNOGENE CO.,LTD (688201056)**Establishment**

| Name | Address | ID/FEI | Business Operations |
|------------------|---------|-----------|------------------------|
| INNOGENE CO.,LTD | | 688201056 | manufacture(70540-040) |

Revised: 9/2017

INNOGENE CO.,LTD