

FOSTER AND THRIVE ANTI DIARRHEAL- loperamide hcl tablet, film coated
Strategic Sourcing Services LLC

McKesson Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20° -25° C (68° -77° F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Principal Display Panel

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

Foster & Thrive™

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls The Symptoms Of Diarrhea

Anti-Diarrheal

Gluten Free

ACTUAL SIZE

12 CAPLETS*

*CAPSULE-SHAPED TABLETS

SEE NEW WARNINGS

000105 / 41915/1

OPEN OTHER END

22453 KM C1

Drug Facts (continued)

Inactive ingredients anhydrous lactose, camphor wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Drug Facts (continued)

Questions or comments?
Call 1-833-358-6431 Monday to Friday 9:00 am to 7:00 pm EST

Drug Facts

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. Loperamide may interact with certain prescription drugs. When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if symptoms get worse or diarrhea lasts for more than 2 days. If pregnant or breast-feeding, ask a health professional before use. These may be signs of a serious condition. You get abdominal swelling or bloating.

Directions
Adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours. Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours. Children 6-8 years (48-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours. Children 2-5 years (24-47 lbs): ask a doctor. Children under 2 years (up to 33 lbs) do not use.

Other information
Store at 20°-25°C (68°-77°F) and see end panel for lot number and expiration date.

Active ingredient (in each caplet)
Loperamide HCl 2 mg An 1-diarrheal

Purpose
Loperamide HCl 2 mg An 1-diarrheal

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
Drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Find night dose on chart, if possible, use weight to dose; otherwise, use age.

Adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours.

Children 9-11 years (60-95 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours.

Children 6-8 years (48-59 lbs): 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours.

Children 2-5 years (24-47 lbs): ask a doctor.

Children under 2 years (up to 33 lbs): do not use.

SAFETY SEALED IN BLISTER UNIT FOR YOUR PROTECTION. IF BROKEN OR TORN RETURN TO STORE.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

NDC 70677-1106-1

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT**

Foster & Thrive™

Loperamide Hydrochloride Tablets, 2 mg


Anti-Diarrheal



©2023 McKesson Corporation
Distributed by: McKesson Corp.,
via Strategic Sourcing Services LLC,
Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com

Rev. 06/23





FOSTER AND THRIVE ANTI DIARRHEAL

loperamide hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1106
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1106-1	12 in 1 CARTON	11/08/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:70677-1106-2	24 in 1 CARTON	11/14/2023	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	11/08/2023	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 12/2023

Strategic Sourcing Services LLC