

FAMOTIDINE- famotidine tablet, film coated
DISCOUNT DRUG MART

Famotidine

Drug Facts

Active ingredient (in each tablet)

Famotidine, USP 10 mg

Famotidine, USP 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

Famotidine USP, 10 mg

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Famotidine USP, 20 mg

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor.
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Famotidine USP, 10 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Famotidine USP, 20 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20° to 25° C (68° to 77° F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide red (only for Famotidine USP, 10 mg), magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Questions?

call **1-800-406-7984**

**DISTRIBUTED BY DRUG MART-FOOD FAIR
MEDINA, OHIO 44256**

PRINCIPAL DISPLAY PANEL - 10 mg Tablet Blister Pack Carton

DISCOUNT
drug
mart
FOOD FAIR

MAY CONTAIN
ANTI-THEFT
DEVICE

Original Strength
ACID CONTROLLER
Famotidine Tablets, USP 10 mg

Acid Reducer
Just One Tablet
Prevents & Relieves Heartburn
Due to Acid Indigestion

*COMPARE TO THE ACTIVE INGREDIENT OF ORIGINAL STRENGTH PEPCID AC®
30 TABLETS

Drug Facts

Active Ingredient (in each tablet)
Famotidine, USP 10 mg Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).



Original Strength

ACID CONTROLLER

Famotidine Tablets, USP 10 mg

Acid Reducer

30 TABLETS



Original Strength

ACID CONTROLLER

Famotidine Tablets, USP 10 mg

Acid Reducer

Just One Tablet
Prevents & Relieves Heartburn
Due to Acid Indigestion

COMPARE TO THE ACTIVE INGREDIENT OF ORIGINAL STRENGTH PEPCID AC®



30 TABLETS

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not eat before bedtime
 - Do not eat rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
 - Do not wear tight-fitting clothing around the stomach
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
 - Quit smoking
 - Overweight, loss weight
 - Quit smoking

Expiration Date:

Batch No.

Non Varnish Area

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.

See end panel for batch number and expiration date.


*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Pepcid AC®.

DISTRIBUTED BY DRUG MART-FOOD FAIR
MEDINA, OHIO 44256 54
www.discount-drugmart.com



SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND CARTON PACKAGE TO: DISCOUNT DRUG MART 2711 COMMERCIAL DRIVE, MEDINA, OHIO 44259

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.
Questions? call 1-800-406-7984



PRINCIPAL DISPLAY PANEL - 20 mg Tablet Bottle Carton

DISCOUNT

drug

mart

FOOD FAIR

MAY CONTAIN

ANTI-THEFT

DEVICE

Maximum Strength

ACID CONTROLLER

Famotidine Tablets, USP 20 mg

Acid Reducer

Just One Tablet

Prevents & Relieves Heartburn

Due to Acid Indigestion

*COMPARE TO THE ACTIVE INGREDIENT OF MAXIMUM STRENGTH PEPCID AC®

25 TABLETS



Maximum Strength

ACID CONTROLLER

Famotidine Tablets, USP 20 mg

Acid Reducer

25 TABLETS



MAY CONTAIN
ANTI-THEFT
DEVICE

Maximum Strength

ACID CONTROLLER

Famotidine Tablets, USP 20 mg

Acid Reducer

Just One Tablet
Prevents & Relieves Heartburn
Due to Acid Indigestion

COMPARE TO THE ACTIVE INGREDIENT OF MAXIMUM STRENGTH PEPCID AC®



25 TABLETS

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Expiration Date:

NON VARNISH

Batch No.:

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.

See end panel for batch number and expiration date. All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Pepcid AC®.

DISTRIBUTED BY DRUG MART-FOOD FAIR

MEDINA, OHIO 44256 54

www.discount-drugmart.com

ROS13



0 19335 112802 8

SATISFACTION GUARANTEED
IF UNSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE
FOR FULL REFUND. IF PACKAGE IS OPENED, RETURN TO STORE FOR
CREDIT. PLEASE DO NOT RETURN TO STORE, SEND RETURN
DISCOUNT DRUG MART, 211 COLUMBUS DRIVE, MEDINA, OHIO 44256

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Drug Facts (continued)

Other Information

- store at 20° to 25° C (68° to 77° F)
- protect from moisture
- read the directions and warnings before use
- Keep the carton. It contains important information.

Inactive Ingredients

colloidal silicon dioxide, microcrystalline cellulose, hydroxypropyl cellulose, hypromellose, magnesium stearate, titanium dioxide, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Questions? call 1-800-406-7984

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- serious condition. See your doctor.
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

Directions

adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.

to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn

do not use more than 2 tablets in 24 hours

children under 12 years: ask a doctor

Stop use and ask a doctor if


- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).



5103161



1913015

Drug Facts	Active Ingredient (in each tablet) Famotidine, USP 20 mg Acid reducer
Drug Facts (continued)	Purpose <ul style="list-style-type: none"> ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition.

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-036
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1200000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-036-25	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA090283	06/25/2010	
------	------------	------------	--

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-037
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYDROXYPROPYL CELLULOSE (1200000 MW) (UNII: RFW2ET671P)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	036
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-037-31	25 in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	07/23/2010	

Labeler - DISCOUNT DRUG MART (047741335)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(53943-036, 53943-037)

Revised: 9/2018

DISCOUNT DRUG MART