

**CEFUROXIME AXETIL- cefuroxime axetil tablet, film coated**  
**Wockhardt Limited**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use cefuroxime axetil tablets safely and effectively. See full prescribing information for cefuroxime axetil tablets.

**CEFUROXIME AXETIL tablets, USP, for oral use**

**Initial U.S. Approval: 1987**

**INDICATIONS AND USAGE**

Cefuroxime axetil is a cephalosporin antibacterial drug indicated for the treatment of the following infections due to susceptible bacteria: (1)

- Pharyngitis/tonsillitis (adults and pediatric patients) (1.1)
- Acute bacterial otitis media (pediatric patients) (1.2)
- Acute bacterial maxillary sinusitis (adults and pediatric patients) (1.3)
- Acute bacterial exacerbations of chronic bronchitis and secondary bacterial infections of acute bronchitis (adults and pediatric patients 13 years and older) (1.4)
- Uncomplicated skin and skin-structure infections (adults and pediatric patients 13 years and older) (1.5)
- Uncomplicated urinary tract infections (adults and pediatric patients 13 years and older) (1.6)
- Uncomplicated gonorrhea (adults and pediatric patients 13 years and older) (1.7)
- Early Lyme disease (adults and pediatric patients 13 years and older) (1.8)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefuroxime axetil tablets and other antibacterial drugs, cefuroxime axetil tablets should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

**DOSAGE AND ADMINISTRATION**

- Tablets and oral suspension are not bioequivalent and are therefore not substitutable on a milligram-per-milligram basis. (2.1)
- Administer tablets with or without food. (2.2)
- Administer cefuroxime axetil tablets as described in the dosage guidelines. (2.2)
- Dosage adjustment is required for patients with impaired renal function. (2.5)

**Adult Patients and Pediatric Patients Dosage Guidelines for Cefuroxime Axetil Tablets**

| Infection   | Dosage                       | Duration (Days) |
|---|------------------------------|-----------------|
| <b>Adults and Adolescents (13 years and older)</b>                              |                              |                 |
| Pharyngitis/tonsillitis (mild to moderate)                                      | 250 mg every 12 hours        | 10              |
| Acute bacterial maxillary sinusitis (mild to moderate)                          | 250 mg every 12 hours        | 10              |
| Acute bacterial exacerbations of chronic bronchitis (mild to moderate)          | 250 or 500 mg every 12 hours | 10              |
| Secondary bacterial infections of acute bronchitis                              | 250 or 500 mg every 12 hours | 5 to 10         |
| Uncomplicated skin and skin-structure infections                                | 250 or 500 mg every 12 hours | 10              |
| Uncomplicated urinary tract infections  | 250 mg every 12 hours        | 7 to 10         |
| Uncomplicated gonorrhea   | 1,000 mg                     | single dose     |
| Early Lyme disease  | 500 mg every 12 hours        | 20              |
| <b>Pediatric Patients younger than 13 years (who can swallow tablets whole)</b> |                              |                 |
| Acute bacterial otitis media  | 250 mg every 12 hours        | 10              |
| Acute bacterial maxillary sinusitis   | 250 mg every 12 hours        | 10              |

**DOSAGE FORMS AND STRENGTHS**

- Tablets: 125 mg, 250 mg and 500 mg (3)

**CONTRAINDICATIONS**

Known hypersensitivity (e.g., anaphylaxis) to cefuroxime axetil tablets or to other  $\beta$ -lactams (e.g., penicillins and cephalosporins). (4)

**WARNINGS AND PRECAUTIONS**

- Serious hypersensitivity (anaphylactic) reactions: In the event of a serious reaction, discontinue cefuroxime axetil tablets and institute appropriate therapy. (5.1)
- *Clostridium difficile*- associated diarrhea (CDAD): If diarrhea occurs, evaluate patients for CDAD. (5.2)

**ADVERSE REACTIONS**

The most common adverse reactions ( $\geq 3\%$ ) for cefuroxime axetil tablets are diarrhea, nausea/vomiting, Jarisch-Herxheimer reaction and vaginitis (early Lyme disease). (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Wockhardt USA LLC. at 1-800-346-6854 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**DRUG INTERACTIONS**

- Oral Contraceptives: Effects on gut flora may lower estrogen reabsorption and reduce efficacy of oral contraceptives. (7.1)
- Drugs that reduce gastric acidity may lower the bioavailability of cefuroxime axetil tablets. (7.2)
- Co-administration with probenecid increases systemic exposure to cefuroxime axetil tablets and is therefore not recommended. (7.3)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2015

**FULL PRESCRIBING INFORMATION: CONTENTS\***

- 1 INDICATIONS AND USAGE**
  - 1.1 Pharyngitis/Tonsillitis
  - 1.2 Acute Bacterial Otitis Media

- 1.3 Acute Bacterial Maxillary Sinusitis
  - 1.4 Acute Bacterial Exacerbations of Chronic Bronchitis and Secondary Bacterial Infections of Acute Bronchitis
  - 1.5 Uncomplicated Skin and Skin-structure Infections
  - 1.6 Uncomplicated Urinary Tract Infections
  - 1.7 Uncomplicated Gonorrhea
  - 1.8 Early Lyme Disease (erythema migrans)
  - 1.10 Usage
  - 2 DOSAGE AND ADMINISTRATION**
    - 2.1 Important Administration Instructions
    - 2.2 Dosage for cefuroxime Axetil Tablets, USP
    - 2.5 Dosage in Patients with Impaired Renal Function
  - 3 DOSAGE FORMS AND STRENGTHS**
  - 4 CONTRAINDICATIONS**
  - 5 WARNINGS AND PRECAUTIONS**
    - 5.1 Anaphylactic Reactions
    - 5.2 Clostridium difficile-associated Diarrhea
    - 5.3 Potential for Microbial Overgrowth
    - 5.4 Development of Drug-resistant Bacteria
    - 5.6 Interference with Glucose Tests
  - 6 ADVERSE REACTIONS**
    - 6.1 Clinical Trials Experience
    - 6.2 Postmarketing Experience
  - 7 DRUG INTERACTIONS**
    - 7.1 Oral Contraceptives
    - 7.2 Drugs that Reduce Gastric Acidity
    - 7.3 Probenecid
    - 7.4 Drug/Laboratory Test Interactions
  - 8 USE IN SPECIFIC POPULATIONS**
    - 8.1 Pregnancy
    - 8.3 Nursing Mothers
    - 8.4 Pediatric Use
    - 8.5 Geriatric Use
    - 8.6 Renal Impairment
  - 10 OVERDOSAGE**
  - 11 DESCRIPTION**
  - 12 CLINICAL PHARMACOLOGY**
    - 12.1 Mechanism of Action
    - 12.3 Pharmacokinetics
    - 12.4 Microbiology
  - 13 NONCLINICAL TOXICOLOGY**
    - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
  - 14 CLINICAL STUDIES**
    - 14.1 Acute Bacterial Maxillary Sinusitis
    - 14.2 Early Lyme Disease
    - 14.3 Secondary Bacterial Infections of Acute Bronchitis
  - 15 REFERENCES**
  - 16 HOW SUPPLIED/STORAGE AND HANDLING**
  - 17 PATIENT COUNSELING INFORMATION**
- \* Sections or subsections omitted from the full prescribing information are not listed.
- 

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 Pharyngitis/Tonsillitis

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (13 years and older) with mild-to-moderate pharyngitis/tonsillitis caused by susceptible strains of *Streptococcus pyogenes*.

##### Limitations of Use

- The efficacy of cefuroxime axetil tablets, USP in the prevention of rheumatic fever was not established in clinical trials.
- The efficacy of cefuroxime axetil tablets, USP in the treatment of penicillin-resistant strains of *Streptococcus pyogenes* has not been demonstrated in clinical trials.

#### 1.2 Acute Bacterial Otitis Media

Cefuroxime axetil tablets, USP are indicated for the treatment of pediatric patients (who can swallow tablets whole) with acute bacterial otitis media caused by susceptible strains of *Streptococcus pneumoniae*, *Haemophilus influenzae* (including  $\beta$ -lactamase-producing strains), *Moraxella catarrhalis* (including  $\beta$ -lactamase-producing strains), or *Streptococcus pyogenes*.

#### 1.3 Acute Bacterial Maxillary Sinusitis

Cefuroxime axetil tablets, USP are indicated for the treatment of adult and pediatric patients (13 years and older) with mild-to-moderate acute bacterial maxillary sinusitis caused by susceptible strains of *Streptococcus pneumoniae* or *Haemophilus influenzae* (non- $\beta$ -lactamase-producing strains only).

##### Limitations of Use

The effectiveness of cefuroxime axetil tablets, USP for sinus infections caused by  $\beta$ -lactamase-producing *Haemophilus influenzae* or *Moraxella catarrhalis* in patients with acute bacterial maxillary

sinusitis was not established due to insufficient numbers of these isolates in the clinical trials [see *Clinical Studies (14.1)*].

#### 1.4 Acute Bacterial Exacerbations of Chronic Bronchitis and Secondary Bacterial Infections of Acute Bronchitis

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (aged 13 and older) with mild-to-moderate acute bacterial exacerbations of chronic bronchitis and secondary bacterial infections of acute bronchitis caused by susceptible strains of *Streptococcus pneumoniae*, *Haemophilus influenzae* ( $\beta$ -lactamase-negative strains), or *Haemophilus parainfluenzae* ( $\beta$ -lactamase-negative strains).

#### 1.5 Uncomplicated Skin and Skin-structure Infections

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (aged 13 and older) with uncomplicated skin and skin-structure infections caused by susceptible strains of *Staphylococcus aureus* (including  $\beta$ -lactamase-producing strains) or *Streptococcus pyogenes*.

#### 1.6 Uncomplicated Urinary Tract Infections

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (aged 13 and older) with uncomplicated urinary tract infections caused by susceptible strains of *Escherichia coli* or *Klebsiella pneumoniae*.

#### 1.7 Uncomplicated Gonorrhea

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (aged 13 and older) with uncomplicated gonorrhea, urethral and endocervical, caused by penicillinase-producing and non-penicillinase-producing susceptible strains of *Neisseria gonorrhoeae* and uncomplicated gonorrhea, rectal, in females, caused by non-penicillinase-producing susceptible strains of *Neisseria gonorrhoeae*.

#### 1.8 Early Lyme Disease (erythema migrans)

Cefuroxime axetil tablets, USP are indicated for the treatment of adult patients and pediatric patients (aged 13 and older) with early Lyme disease (erythema migrans) caused by susceptible strains of *Borrelia burgdorferi*.

#### 1.10 Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefuroxime axetil tablets, USP and other antibacterial drugs, cefuroxime axetil tablets, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Important Administration Instructions

- Cefuroxime axetil tablets, USP and cefuroxime axetil for oral suspension are not bioequivalent and are therefore not substitutable on a milligram-per-milligram basis [see *Clinical Pharmacology (12.3)*].
- Administer cefuroxime axetil tablets, USP as described in the appropriate dosage guidelines [see *Dosage and Administration (2.2)*].
- Administer cefuroxime axetil tablets, USP with or without food.
- Pediatric patients (aged 13 years and older) who cannot swallow the cefuroxime axetil tablets, USP whole should receive cefuroxime axetil for oral suspension because the tablet has a strong, persistent bitter taste when crushed [see *Dosage and Administration (2.2)*].

### 2.2 Dosage for cefuroxime Axetil Tablets, USP

Administer cefuroxime axetil tablets, USP as described in the dosage guidelines table below with or without food.

**Table 1. Adult Patients and Pediatric Patients Dosage Guidelines for Cefuroxime Axetil Tablets, USP**

| Infection   | Dosage                       | Duration (Days) |
|---|------------------------------|-----------------|
| <b>Adults and Adolescents (13 years and older)</b>  |                              |                 |
| Pharyngitis/tonsillitis (mild to moderate)  | 250 mg every 12 hours        | 10              |
| Acute bacterial maxillary sinusitis (mild to moderate)                                      | 250 mg every 12 hours        | 10              |
| Acute bacterial exacerbations of chronic bronchitis (mild to moderate)                      | 250 or 500 mg every 12 hours | 10 <sup>a</sup> |
| Secondary bacterial infections of acute bronchitis  | 250 or 500 mg every 12 hours | 5 to 10         |
| Uncomplicated skin and skin-structure infections  | 250 or 500 mg every 12 hours | 10              |
| Uncomplicated urinary tract infections  | 250 mg every 12 hours        | 7 to 10         |
| Uncomplicated gonorrhea   | 1,000 mg                     | single dose     |
| Early Lyme disease  | 500 mg every 12 hours        | 20              |
| <b>Pediatric Patients younger than 13 years (who can swallow tablets whole)<sup>b</sup></b> |                              |                 |
| Acute bacterial otitis media  | 250 mg                       | 10              |

|                                     |                          |    |
|-------------------------------------|--------------------------|----|
|                                     | every 12 hours           |    |
| Acute bacterial maxillary sinusitis | 250 mg<br>every 12 hours | 10 |

<sup>a</sup> The safety and effectiveness of cefuroxime axetil tablets, USP administered for less than 10 days in patients with acute exacerbations of chronic bronchitis have not been established.

<sup>b</sup> When crushed, the tablet has a strong, persistent bitter taste. Therefore, patients who cannot swallow the tablet whole should receive the oral suspension.

## 2.5 Dosage in Patients with Impaired Renal Function

A dosage interval adjustment is required for patients whose creatinine clearance is <30 mL/min, as listed in Table 4 below, because cefuroxime is eliminated primarily by the kidney [see *Clinical Pharmacology* (12.3)].

**Table 4. Dosing in Adults with Renal Impairment**

| Creatinine Clearance (mL/min) | Recommended Dosage  |
|-------------------------------|---|
| ≥30                           | No dosage adjustment  |
| 10 to <30                     | Standard individual dose given every 24 hours                                 |
| <10<br>(without hemodialysis) | Standard individual dose given every 48 hours                                 |
| Hemodialysis                  | A single additional standard dose should be given at the end of each dialysis |

## 3 DOSAGE FORMS AND STRENGTHS

Cefuroxime axetil tablets are white film-coated round-shaped biconvex unscored tablets available in the following strengths:

- 125 mg of cefuroxime as cefuroxime axetil embossed with W920 on one side and other side plain.
- 250 mg of cefuroxime as cefuroxime axetil embossed with W921 on one side and other side plain.
- 500 mg of cefuroxime as cefuroxime axetil embossed with W922 on one side and other side plain.

## 4 CONTRAINDICATIONS

Cefuroxime axetil tablets are contraindicated in patients with a known hypersensitivity (e.g., anaphylaxis) to cefuroxime axetil or to other β-lactam antibacterial drugs (e.g., penicillins and cephalosporins).

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Anaphylactic Reactions

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients on β-lactam antibacterials. These reactions are more likely to occur in individuals with a history of β-lactam hypersensitivity and/or a history of sensitivity to multiple allergens. There have been reports of individuals with a history of penicillin hypersensitivity who have experienced severe reactions when treated with cephalosporins. Cefuroxime axetil tablets are contraindicated in patients with a known hypersensitivity to cefuroxime axetil or other β-lactam antibacterial drugs [see *Contraindications* (4)]. Before initiating therapy with cefuroxime axetil tablets, inquire about previous hypersensitivity reactions to penicillins, cephalosporins, or other allergens. If an allergic reaction occurs, discontinue cefuroxime axetil tablets and institute appropriate therapy.

### 5.2 Clostridium difficile-associated Diarrhea

*Clostridium difficile*-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including cefuroxime axetil tablets, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin-producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

### 5.3 Potential for Microbial Overgrowth

The possibility of superinfections with fungal or bacterial pathogens should be considered during therapy.

### 5.4 Development of Drug-resistant Bacteria

Prescribing cefuroxime axetil tablets either in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

### 5.6 Interference with Glucose Tests

A false-positive result for glucose in the urine may occur with copper reduction tests, and a false-negative result for blood/plasma glucose may occur with ferricyanide tests in subjects receiving cefuroxime axetil tablets [see *Drug Interactions* (7.4)].

## 6 ADVERSE REACTIONS

The following serious and otherwise important adverse reaction is described in greater detail in the Warnings and Precautions section of the label:

Anaphylactic Reactions [see Warnings and Precautions (5.1)]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### Tablets

**Multiple-dose Dosing Regimens with 7 to 10 Days' Duration:** In multiple-dose clinical trials, 912 subjects were treated with cefuroxime axetil tablets (125 to 500 mg twice daily). It is noted that 125 mg twice daily is not an approved dosage. Twenty (2.2%) subjects discontinued medication due to adverse reactions. Seventeen (85%) of the 20 subjects who discontinued therapy did so because of gastrointestinal disturbances, including diarrhea, nausea, vomiting, and abdominal pain. The percentage of subjects treated with cefuroxime axetil tablets who discontinued study drug because of adverse reactions was similar at daily doses of 1,000, 500, and 250 mg (2.3%, 2.1%, and 2.2%, respectively). However, the incidence of gastrointestinal adverse reactions increased with the higher recommended doses.

The adverse reactions in Table 5 are for subjects (n = 912) treated with cefuroxime axetil tablets in multiple-dose clinical trials.

**Table 5. Adverse Reactions (≥1%) after Multiple-dose Regimens with Cefuroxime Axetil Tablets**

| Adverse Reaction                            | Cefuroxime Axetil Tablets<br>(n = 912) |
|---|--|
| <b>Blood and lymphatic system disorders</b> |  |
| Eosinophilia                                | 1%                                     |
| <b>Gastrointestinal disorders</b>           |  |
| Diarrhea                                    | 4%                                     |
| Nausea/Vomiting                             | 3%                                     |
| <b>Investigations</b>                       |  |
| Transient elevation in AST                  | 2%                                     |
| Transient elevation in ALT                  | 2%                                     |
| Transient elevation in LDH                  | 1%                                     |

The following adverse reactions occurred in less than 1% but greater than 0.1% of subjects (n = 912) treated with cefuroxime axetil tablets in multiple-dose clinical trials.

*Immune System Disorders:* Hives, swollen tongue.

*Metabolism and Nutrition Disorders:* Anorexia.

*Nervous System Disorders:* Headache.

*Cardiac Disorders:* Chest pain.

*Respiratory Disorders:* Shortness of breath.

*Gastrointestinal Disorders:* Abdominal pain, abdominal cramps, flatulence, indigestion, mouth ulcers.

*Skin and Subcutaneous Tissue Disorders:* Rash, itch.

*Renal and Urinary Disorders:* Dysuria.

*Reproductive System and Breast Disorders:* Vaginitis, vulvar itch.

*General Disorders and Administration Site Conditions:* Chills, sleepiness, thirst. *Investigations:* Positive Coombs' test.

**5-Day Regimen:** In clinical trials using cefuroxime axetil tablets 250 mg twice daily in the treatment of secondary bacterial infections of acute bronchitis, 399 subjects were treated for 5 days and 402 subjects were treated for 10 days. No difference in the occurrence of adverse reactions was found between the 2 regimens.

**Early Lyme Disease with 20-Day Regimen:** Two multicenter trials assessed cefuroxime axetil tablets 500 mg twice daily for 20 days. The most common drug-related adverse experiences were diarrhea (10.6%), Jarisch-Herxheimer reaction (5.6%), and vaginitis (5.4%). Other adverse experiences occurred with frequencies comparable to those reported with 7 to 10 days' dosing.

**Single-dose Regimen for Uncomplicated Gonorrhea:** In clinical trials using a single 1,000-mg dose of cefuroxime axetil tablets, 1,061 subjects were treated for uncomplicated gonorrhea.

The adverse reactions in Table 6 were for subjects treated with a single dose of 1,000 mg cefuroxime axetil tablets in US clinical trials.

**Table 6. Adverse Reactions (≥1%) after Single-dose Regimen with 1,000-mg Cefuroxime Axetil Tablets for Uncomplicated Gonorrhea**

| Adverse Reaction                  | Cefuroxime Axetil Tablets<br>(n = 1,061) |
|-----------------------------------|--|
| <b>Gastrointestinal disorders</b> |  |
| Nausea/Vomiting                   | 7%                                       |
| Diarrhea                          | 4%                                       |

The following adverse reactions occurred in less than 1% but greater than 0.1% of subjects (n = 1,061) treated with a single dose of cefuroxime axetil tablets 1,000 mg for uncomplicated gonorrhea in US clinical trials.

*Infections and Infestations:* Vaginal candidiasis.

*Nervous System Disorders:* Headache, dizziness, somnolence.

*Cardiac Disorders:* Tightness/pain in chest, tachycardia.

*Gastrointestinal Disorders:* Abdominal pain, dyspepsia.

*Skin and Subcutaneous Tissue Disorders:* Erythema, rash, pruritus.

*Musculoskeletal and Connective Tissue Disorders:* Muscle cramps, muscle stiffness, muscle spasm of neck, lockjaw-type reaction.

*Renal and Urinary Disorders:* Bleeding/pain in urethra, kidney pain.

*Reproductive System and Breast Disorders:* Vaginal itch, vaginal discharge.

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of cefuroxime axetil tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### Blood and Lymphatic System Disorders

Hemolytic anemia, leukopenia, pancytopenia, thrombocytopenia.

### Gastrointestinal Disorders

Pseudomembranous colitis [see *Warnings and Precautions (5.2)*].

### Hepatobiliary Disorders

Hepatic impairment including hepatitis and cholestasis, jaundice.

### Immune System Disorders

Anaphylaxis, serum sickness-like reaction.

### Investigations

Increased prothrombin time.

### Nervous System Disorders

Seizure, encephalopathy.

### Renal and Urinary Disorders

Renal dysfunction.

### Skin and Subcutaneous Tissue Disorders

Angioedema, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, urticaria.

## 7 DRUG INTERACTIONS

### 7.1 Oral Contraceptives

Cefuroxime axetil may affect the gut flora, leading to lower estrogen reabsorption and reduced efficacy of combined oral estrogen/progesterone contraceptives. Counsel patients to consider alternate supplementary (non-hormonal) contraceptive measures during treatment.

### 7.2 Drugs that Reduce Gastric Acidity

Drugs that reduce gastric acidity may result in a lower bioavailability of cefuroxime axetil tablets compared with administration in the fasting state. Administration of drugs that reduce gastric acidity may negate the food effect of increased absorption of cefuroxime axetil tablets when administered in the postprandial state. Administer cefuroxime axetil tablets at least 1 hour before or 2 hours after administration of short-acting antacids. Histamine-2 (H<sub>2</sub>) antagonists and proton pump inhibitors should be avoided.

### 7.3 Probenecid

Concomitant administration of probenecid with cefuroxime axetil tablets increases serum concentrations of cefuroxime [see *Clinical Pharmacology (12.3)*]. Co-administration of probenecid with cefuroxime axetil is not recommended.

### 7.4 Drug/Laboratory Test Interactions

A false-positive reaction for glucose in the urine may occur with copper reduction tests (e.g., Benedict's or Fehling's solution), but not with enzyme-based tests for glycosuria. As a false-negative result may occur in the ferricyanide test, it is recommended that either the glucose oxidase or hexokinase method be used to determine blood/plasma glucose levels in patients receiving cefuroxime axetil. The presence of cefuroxime does not interfere with the assay of serum and urine creatinine by the alkaline picrate method.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B. There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, cefuroxime axetil tablets should be used during pregnancy only if clearly needed.

Reproduction studies have been performed in mice at doses up to 3,200 mg/kg/day (14 times the recommended maximum human dose based on body surface area) and in rats at doses up to 1,000 mg/kg/day (9 times the recommended maximum human dose based on body surface area) and have revealed no evidence of impaired fertility or harm to the fetus due to cefuroxime axetil.

### 8.3 Nursing Mothers

Because cefuroxime is excreted in human milk, caution should be exercised when cefuroxime axetil is administered to a nursing woman.

### 8.4 Pediatric Use

The safety and effectiveness of cefuroxime axetil tablets have been established for pediatric patients aged 3 months to 12 years for acute bacterial maxillary sinusitis based upon its approval in adults. Use of cefuroxime axetil tablets in pediatric patients is supported by pharmacokinetic and safety data in adults and pediatric patients, and by clinical and microbiological data from adequate and well-controlled trials of the treatment of acute bacterial maxillary sinusitis in adults and of acute otitis media with effusion in pediatric patients. It is also supported by postmarketing adverse events surveillance. [See *Indications and Usage (1), Dosage and Administration (2), Adverse Reactions (6), Clinical Pharmacology (12.3).*]

### 8.5 Geriatric Use

Of the total number of subjects who received cefuroxime axetil tablets in 20 clinical trials, 375 were aged 65 and older while 151 were aged 75 and older. No overall differences in safety or effectiveness were observed between these subjects and younger adult subjects. Reported clinical experience has not identified differences in responses between the elderly and younger adult patients, but greater sensitivity of some older individuals cannot be ruled out.

Cefuroxime is substantially excreted by the kidney, and the risk of adverse reactions may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

### 8.6 Renal Impairment

Reducing the dosage of cefuroxime axetil tablets is recommended for adult patients with severe renal impairment (creatinine clearance <30 mL/min) [see *Dosage and Administration (2.5), Clinical Pharmacology (12.3)*].

## 10 OVERDOSAGE


Overdosage of cephalosporins can cause cerebral irritation leading to convulsions or encephalopathy. Serum levels of cefuroxime can be reduced by hemodialysis and peritoneal dialysis.

## 11 DESCRIPTION

Cefuroxime axetil tablets, USP contain cefuroxime as cefuroxime axetil. Cefuroxime axetil is a semisynthetic, cephalosporin antibacterial drug for oral administration.

The chemical name of cefuroxime axetil (1-(acetyloxy) ethyl ester of cefuroxime) is (RS)-1-hydroxyethyl (6R,7R)-7-[2-(2-furyl)glyoxyl-amido]-3-(hydroxymethyl)-8-oxo-5-thia-1-azabicyclo[4.2.0]-oct-2-ene-2-carboxylate, 7<sup>2</sup>-(Z)-(O-methyl-oxime), 1-acetate 3-carbamate. Its molecular formula is C<sub>20</sub>H<sub>22</sub>N<sub>4</sub>O<sub>10</sub>S, and it has a molecular weight of 510.48.

Cefuroxime axetil is in the amorphous form and has the following structural formula:



Cefuroxime axetil tablets, USP are film-coated and contain the equivalent of 125, 250, or 500 mg of cefuroxime as cefuroxime axetil. Cefuroxime axetil tablets, USP contain the inactive ingredients colloidal silicon dioxide, prosolv SMCC 50, sodium lauryl sulfate, croscarmellose sodium, hydrogenated castor oil, hypromellose, propylene glycol, titanium dioxide, dewaxed shellac powder, dibutyl sebacate and polysorbate 80.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Cefuroxime axetil is an antibacterial drug [see *Clinical Pharmacology (12.4)*].

### 12.3 Pharmacokinetics

#### Absorption

After oral administration, cefuroxime axetil is absorbed from the gastrointestinal tract and rapidly hydrolyzed by nonspecific esterases in the intestinal mucosa and blood to cefuroxime. Serum pharmacokinetic parameters for cefuroxime following administration of cefuroxime axetil tablets to adults are shown in Table 8.

Table 8. Pharmacokinetics of Cefuroxime Administered in the Postprandial State as Cefuroxime Axetil Tablets to Adults<sup>a</sup>

| Dose <sup>b</sup> (Cefuroxime Equivalent) | Peak Plasma Concentration (mcg/mL) | Time of Peak Plasma Concentration (h) | Mean Elimination Half-life (h) | AUC (mcg(h/mL)) |
|---|------------------------------------|---------------------------------------|--------------------------------|-----------------|
| 125 mg                                    | 2.1                                | 2.2                                   | 1.2                            | 6.7             |
| 250 mg                                    | 4.1                                | 2.5                                   | 1.2                            | 12.9            |
| 500 mg                                    | 7                                  | 3                                     | 1.2                            | 27.4            |
| 1,000 mg                                  | 13.6                               | 2.5                                   | 1.3                            | 50              |

<sup>a</sup> Mean values of 12 healthy adult volunteers.

<sup>b</sup> Drug administered immediately after a meal.

**Food Effect:** Absorption of the tablet is greater when taken after food (absolute bioavailability increases from 37% to 52%). Despite this difference in absorption, the clinical and bacteriologic responses of subjects were independent of food intake at the time of tablet administration in 2 trials where this was assessed.

All pharmacokinetic and clinical effectiveness and safety trials in pediatric subjects using the suspension formulation were conducted in the fed state. No data are available on the absorption kinetics of the suspension formulation when administered to fasted pediatric subjects.

**Lack of Bioequivalence:** Oral suspension was not bioequivalent to tablets when tested in healthy adults. The tablet and oral suspension formulations are NOT substitutable on a milligram-per-milligram basis. The area under the curve for the suspension averaged 91% of that for the tablet, and the peak plasma concentration for the suspension averaged 71% of the peak plasma concentration of the tablets. Therefore, the safety and effectiveness of both the tablet and oral suspension formulations were established in separate clinical trials.

#### Distribution

Cefuroxime is distributed throughout the extracellular fluids. Approximately 50% of serum cefuroxime is bound to protein.

#### Metabolism

The axetil moiety is metabolized to acetaldehyde and acetic acid.

#### Excretion

Cefuroxime is excreted unchanged in the urine; in adults, approximately 50% of the administered dose is recovered in the urine within 12 hours. The pharmacokinetics of cefuroxime in pediatric subjects have not been studied. Until further data are available, the renal elimination of cefuroxime axetil established in adults should not be extrapolated to pediatric subjects.

#### Specific Populations

**Renal Impairment:** In a trial of 28 adults with normal renal function or severe renal impairment (creatinine clearance <30 mL/min), the elimination half-life was prolonged in relation to severity of renal impairment. Prolongation of the dosage interval is recommended in adult patients with creatinine clearance <30 mL/min [see *Dosage and Administration (2.5)*].

**Geriatric Patients:** In a trial of 20 elderly subjects (mean age = 83.9 years) having a mean creatinine clearance of 34.9 mL/min, the mean serum elimination half-life was prolonged to 3.5 hours; however, despite the lower elimination of cefuroxime in geriatric patients, dosage adjustment based on age is not necessary [see *Use in Specific Populations (8.5)*].

#### Drug Interactions

Concomitant administration of probenecid with cefuroxime axetil tablets increases the cefuroxime area under the serum concentration versus time curve and maximum serum concentration by 50% and 21%, respectively.

## **12.4 Microbiology**

### Mechanism of Action

Cefuroxime axetil is a bactericidal agent that acts by inhibition of bacterial cell wall synthesis. Cefuroxime axetil has activity in the presence of some  $\beta$ -lactamases, both penicillinases and cephalosporinases, of gram-negative and gram-positive bacteria.

### Mechanism of Resistance

Resistance to cefuroxime axetil is primarily through hydrolysis by  $\beta$ -lactamase, alteration of penicillin-binding proteins (PBPs), decreased permeability, and the presence of bacterial efflux pumps.

Susceptibility to cefuroxime axetil will vary with geography and time; local susceptibility data should be consulted, if available. Beta-lactamase-negative, ampicillin-resistant (BLNAR) isolates of *H. influenzae* should be considered resistant to cefuroxime axetil.

Cefuroxime axetil has been shown to be active against most isolates of the following bacteria, both *in vitro* and in clinical infections [see *Indications and Usage (1)*]:

- Gram-positive bacteria

*Staphylococcus aureus* (methicillin-susceptible isolates only)

*Streptococcus pneumoniae*

*Streptococcus pyogenes*

- Gram-negative bacteria

*Escherichia coli*<sup>a</sup>

*Klebsiella pneumoniae*<sup>a</sup>

*Haemophilus influenzae*

*Haemophilus parainfluenzae*

*Moraxella catarrhalis*

*Neisseria gonorrhoeae*

<sup>a</sup> Most extended spectrum  $\beta$ -lactamase (ESBL)-producing and carbapenemase-producing isolates are resistant to cefuroxime axetil.

- Spirochetes

*Borrelia burgdorferi*

The following *in vitro* data are available, but their clinical significance is unknown. At least 90 percent

of the following microorganisms exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for cefuroxime axetil of 1 mcg/mL. However, the efficacy of cefuroxime axetil in treating clinical infections due to these microorganisms has not been established in adequate and well-controlled clinical trials.

- Gram-positive bacteria

*Staphylococcus epidermidis* (methicillin-susceptible isolates only)

*Staphylococcus saprophyticus* (methicillin-susceptible isolates only)

*Streptococcus agalactiae*

- Gram-negative bacteria

*Morganella morganii*

*Proteus inconstans*

*Proteus mirabilis*

*Providencia rettgeri*

- Anaerobic bacteria

*Peptococcus niger*

#### Susceptibility Test Methods

When available, the clinical microbiology laboratory should provide the results of *in vitro* susceptibility tests for antimicrobial drug products used in local hospitals and practice areas to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting an antibacterial drug product for treatment.

**Dilution Techniques:** Quantitative methods are used to determine antimicrobial MICs. These MICs provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized test method (broth or agar).<sup>1,2</sup> The MIC values should be interpreted according to criteria provided in Table 10.<sup>2,3</sup>

**Diffusion Techniques:** Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. The zone size provides an estimate of the susceptibility of bacteria to antimicrobial compounds. The zone size should be determined using a standardized test method.<sup>4</sup> This procedure uses paper disks impregnated with 30 mcg cefuroxime axetil to test the susceptibility of microorganisms to cefuroxime axetil. The disk diffusion interpretive criteria are provided in Table 10.<sup>3</sup>

**Table 10. Susceptibility Test Interpretive Criteria for Cefuroxime Axetil**

| Pathogen                                  | Minimum Inhibitory Concentrations (mcg/mL) |              |           | Disk Diffusion Zone Diameters (mm) |              |           |
|---|--|--------------|-----------|------------------------------------|--------------|-----------|
|   | (S)  | (I)          | (R)       | (S)                                | (I)          | (R)       |
|   | Susceptible                                | Intermediate | Resistant | Susceptible                        | Intermediate | Resistant |
| <i>Enterobacteriaceae</i> <sup>a</sup>    | ≤4   | 8 to 16      | ≥32       | ≥23                                | 15 to 22     | ≤14       |
| <i>Haemophilus</i> spp. <sup>a,b</sup>    | ≤4   | 8            | ≥16       | ≥20                                | 17 to 19     | ≤16       |
| <i>Moraxella catarrhalis</i> <sup>a</sup> | ≤4   | 8            | ≥16       | -                                  | -            | -         |
| <i>Streptococcus pneumoniae</i>           | ≤1   | 2            | ≥4        | -                                  | -            | -         |

<sup>a</sup> For *Enterobacteriaceae*, *Haemophilus* spp., and *Moraxella catarrhalis*, susceptibility interpretive criteria are based on a dose of 500 mg every 12 hours in patients with normal renal function.

<sup>b</sup> *Haemophilus* spp. includes only isolates of *H. influenzae* and *H. parainfluenzae*.

Susceptibility of staphylococci to cefuroxime may be deduced from testing only penicillin and either ceftiofur or oxacillin.

Susceptibility of *Streptococcus pyogenes* may be deduced from testing penicillin.<sup>3</sup>

A report of "Susceptible" indicates that the antimicrobial drug is likely to inhibit growth of the pathogen if the antimicrobial drug reaches the concentration usually achievable at the site of infection. A report of "Intermediate" indicates that the result should be considered equivocal, and if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where a high dosage of drug can be used. This category also provides a buffer zone that prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that the antimicrobial drug is not likely to inhibit growth of the pathogen if the antimicrobial drug reaches the concentrations usually achievable at the infection site; other therapy should be selected.

**Quality Control:** Standardized susceptibility test procedures require the use of laboratory controls to monitor and ensure the accuracy and precision of supplies and reagents used in the assay, and the techniques of the individual performing the test.<sup>1,2,4</sup> The QC ranges for MIC and disk diffusion testing using the 30-mcg disk are provided in Table 11.<sup>3</sup>

**Table 11. Acceptable Quality Control (QC) Ranges for Cefuroxime Axetil**

| QC Strain                                  | Minimum Inhibitory Concentrations (mcg/mL) | Disk Diffusion Zone Diameters (mm) |
|--|--|------------------------------------|
| <i>Escherichia coli</i> ATCC 25922         | 2 to 8                                     | 20 to 26                           |
| <i>Staphylococcus aureus</i> ATCC 25923    | -  | 27 to 35                           |
| <i>Staphylococcus aureus</i> ATCC 29213    | 0.5 to 2                                   | -                                  |
| <i>Streptococcus pneumoniae</i> ATCC 49619 | 0.25 to 1                                  | -                                  |
| <i>Haemophilus influenzae</i> ATCC 49766   | 0.25 to 1                                  | 28 to 36                           |
| <i>Neisseria gonorrhoeae</i> ATCC 49226    | 0.25 to 1                                  | 33 to 41                           |

### 13 NONCLINICAL TOXICOLOGY

#### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Although lifetime studies in animals have not been performed to evaluate carcinogenic potential, no mutagenic activity was found for cefuroxime axetil in a battery of bacterial mutation tests. Positive results were obtained in an *in vitro* chromosome aberration assay; however, negative results were found in an *in vivo* micronucleus test at doses up to 1.5 g/kg. Reproduction studies in rats at doses up to 1,000 mg/kg/day (9 times the recommended maximum human dose based on body surface area) have revealed no impairment of fertility.

### 14 CLINICAL STUDIES

#### 14.1 Acute Bacterial Maxillary Sinusitis

One adequate and well-controlled trial was performed in subjects with acute bacterial maxillary sinusitis. In this trial, each subject had a maxillary sinus aspirate collected by sinus puncture before treatment was initiated for presumptive acute bacterial sinusitis. All subjects had radiographic and clinical evidence of acute maxillary sinusitis. In the trial, the clinical effectiveness of cefuroxime axetil tablets in treating acute maxillary sinusitis was comparable to an oral antimicrobial agent containing a specific  $\beta$ -lactamase inhibitor. However, microbiology data demonstrated cefuroxime axetil tablets to be effective in treating acute bacterial maxillary sinusitis due only to *Streptococcus pneumoniae* or non- $\beta$ -lactamase-producing *Haemophilus influenzae*. Insufficient numbers of  $\beta$ -lactamase-producing *Haemophilus influenzae* and *Moraxella catarrhalis* isolates were obtained in this trial to adequately evaluate the effectiveness of cefuroxime axetil tablets in treating acute bacterial maxillary sinusitis due to these 2 organisms.

This trial randomized 317 adult subjects, 132 subjects in the United States and 185 subjects in South America. Table 12 shows the results of the intent-to-treat analysis.

**Table 12. Clinical Effectiveness of Cefuroxime Axetil Tablets in the Treatment of Acute Bacterial Maxillary Sinusitis**

|  | US Subjects <sup>a</sup>                                       |                                  | South American Subjects <sup>b</sup>                           |                                  |
|--|--|----------------------------------|--|----------------------------------|
|  | Cefuroxime Axetil Tablets<br>250 mg<br>Twice Daily<br>(n = 49) | Control <sup>c</sup><br>(n = 43) | Cefuroxime Axetil Tablets<br>250 mg<br>Twice Daily<br>(n = 49) | Control <sup>c</sup><br>(n = 43) |
| Clinical success<br>(cure + improvement) | 65%  | 53%                              | 77%  | 74%                              |
| Clinical cure                            | 53%  | 44%                              | 72%  | 64%                              |
| Clinical improvement                     | 12%  | 9%                               | 5%   | 10%                              |

<sup>a</sup> 95% confidence interval around the success difference [-0.08, +0.32].

<sup>b</sup> 95% confidence interval around the success difference [-0.10, +0.16].

<sup>c</sup> Control was an antibacterial drug containing a  $\beta$ -lactamase inhibitor.

In this trial and in a supporting maxillary puncture trial, 15 evaluable subjects had non- $\beta$ -lactamase-producing *Haemophilus influenzae* as the identified pathogen. Of these, 67% (10/15) had this pathogen eradicated. Eighteen (18) evaluable subjects had *Streptococcus pneumoniae* as the identified pathogen. Of these, 83% (15/18) had this pathogen eradicated.

#### 14.2 Early Lyme Disease

Two adequate and well-controlled trials were performed in subjects with early Lyme disease. All subjects presented with physician-documented erythema migrans, with or without systemic manifestations of infection. Subjects were assessed at 1 month posttreatment for success in treating early Lyme disease (Part I) and at 1 year posttreatment for success in preventing the progression to the sequelae of late Lyme disease (Part II).

A total of 355 adult subjects (181 treated with cefuroxime axetil and 174 treated with doxycycline) were randomized in the 2 trials, with diagnosis of early Lyme disease confirmed in 79% (281/355). The clinical diagnosis of early Lyme disease in these subjects was validated by 1) blinded expert reading of photographs, when available, of the pretreatment erythema migrans skin lesion, and 2) serologic confirmation (using enzyme-linked immunosorbent assay [ELISA] and immunoblot assay ["Western" blot]) of the presence of antibodies specific to *Borrelia burgdorferi*, the etiologic agent of Lyme disease. The efficacy data in Table 14 are specific to this "validated" patient subset, while the safety data below reflect the entire patient population for the 2 trials. Clinical data for evaluable subjects in the "validated" patient subset are shown in Table 13.

**Table 13. Clinical Effectiveness of Cefuroxime Axetil Tablets Compared with Doxycycline in the Treatment of Early Lyme Disease**

|  | Part I<br>(1 Month after 20 Days<br>of Treatment) <sup>a</sup>  |   | Part II<br>(1 Year after 20 Days<br>of Treatment) <sup>b</sup>                |  |
|--|---|---|---|--|
|  | Cefuroxime Axetil Tablets<br>500 mg<br>Twice Daily<br>(n = 125) | Doxycycline<br>100 mg<br>3 Times Daily<br>(n = 108) | Cefuroxime Axetil Tablets<br>500 mg<br>Twice Daily<br>(n = 105 <sup>c</sup> ) | Doxycycline<br>100 mg<br>3 Times Daily<br>(n = 83 <sup>c</sup> ) |
| Satisfactory clinical outcome <sup>d</sup> | 91%   | 93%   | 84%   | 87%  |

|                       |     |     |     |     |
|-----------------------|-----|-----|-----|-----|
| Clinical cure/success | 72% | 73% | 73% | 73% |
| Clinical improvement  | 19% | 19% | 10% | 13% |

<sup>a</sup>95% confidence interval around the satisfactory difference for Part I (-0.08, +0.05).

<sup>b</sup>95% confidence interval around the satisfactory difference for Part II (-0.13, +0.07).

<sup>c</sup>n's include subjects assessed as unsatisfactory clinical outcomes (failure + recurrence) in Part I (cefuroxime axetil tablets - 11 [5 failure, 6 recurrence]; doxycycline - 8 [6 failure, 2 recurrence]).

<sup>d</sup>Satisfactory clinical outcome includes cure + improvement (Part I) and success + improvement (Part II).

Cefuroxime axetil tablets and doxycycline were effective in prevention of the development of sequelae of late Lyme disease.

While the incidence of drug-related gastrointestinal adverse reactions was similar in the 2 treatment groups (cefuroxime axetil - 13%; doxycycline - 11%), the incidence of drug-related diarrhea was higher in the cefuroxime axetil arm versus the doxycycline arm (11% versus 3%, respectively).

### 14.3 Secondary Bacterial Infections of Acute Bronchitis

Four randomized, controlled clinical trials were performed comparing 5 days versus 10 days of cefuroxime axetil tablets for the treatment of subjects with secondary bacterial infections of acute bronchitis. These trials enrolled a total of 1,253 subjects (Study 1 n = 360; Study 2 n = 177; Study 3 n = 362; Study 4 n = 354). The protocols for Study 1 and Study 2 were identical and compared cefuroxime axetil tablets 250 mg twice daily for 5 days, cefuroxime axetil tablets 250 mg twice daily for 10 days, and AUGMENTIN® (amoxicillin/clavulanate potassium) 500 mg 3 times daily for 10 days. These 2 trials were conducted simultaneously. Study 3 and Study 4 compared cefuroxime axetil tablets 250 mg twice daily for 5 days, cefuroxime axetil tablets 250 mg twice daily for 10 days, and CECLOR® (cefaclor) 250 mg 3 times daily for 10 days. They were otherwise identical to Study 1 and Study 2 and were conducted over the following 2 years. Subjects were required to have polymorphonuclear cells present on the Gram stain of their screening sputum specimen, but isolation of a bacterial pathogen from the sputum culture was not required for inclusion. Table 14 demonstrates the results of the clinical outcome analysis of the pooled trials Study 1/Study 2 and Study 3/Study 4, respectively.

**Table 14. Clinical Effectiveness of Cefuroxime Axetil Tablets 250 mg Twice Daily in Secondary Bacterial Infections of Acute Bronchitis: Comparison of 5 versus 10 Days' Treatment Duration**

|  | Study 1 and Study 2 <sup>a</sup> |                     | Study 3 and Study 4 <sup>b</sup> |                     |
|--|----------------------------------|---------------------|----------------------------------|---------------------|
|  | 5 Day<br>(n = 127)               | 10 Day<br>(n = 139) | 5 Day<br>(n = 173)               | 10 Day<br>(n = 192) |
| Clinical success<br>(cure + improvement) | 80%                              | 87%                 | 84%                              | 82%                 |
| Clinical cure                            | 61%                              | 70%                 | 73%                              | 72%                 |
| Clinical improvement                     | 19%                              | 17%                 | 11%                              | 10%                 |

<sup>a</sup> 95% confidence interval around the success difference [-0.164, +0.029].

<sup>b</sup> 95% confidence interval around the success difference [-0.061, +0.103].

The response rates for subjects who were both clinically and bacteriologically evaluable were consistent with those reported for the clinically evaluable subjects.

## 15 REFERENCES

1. Clinical and Laboratory Standards Institute (CLSI). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard - Tenth Edition. 2015. CLSI document M07-A10, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA.
2. Clinical and Laboratory Standards Institute (CLSI). Methods for Antimicrobial Dilution and Disk Susceptibility Testing for Infrequently Isolated or Fastidious Bacteria: Approved Guidelines - Second Edition. 2010. CLSI document M45-A2, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA.
3. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-fifth Informational Supplement. 2015. CLSI document M100-S25, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA.
4. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Disk Diffusion Susceptibility Tests; Approved Standard – Twelfth Edition. 2015. CLSI document M02-A12, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Cefuroxime axetil tablets USP, 125 mg of cefuroxime as cefuroxime axetil, are white film-coated, round-shaped biconvex unscored tablets embossed with W920 on one side and plain on other side as follows:

|                    |                  |
|--------------------|------------------|
| 20 Tablets/ Bottle | NDC 64679-920-01 |
| 60 Tablets/ Bottle | NDC 64679-920-02 |

Cefuroxime axetil tablets USP, 250 mg of cefuroxime as cefuroxime axetil, are white film-coated, round-shaped biconvex unscored tablets embossed with W921 on one side and plain on other side as follows:

|                    |                  |
|--------------------|------------------|
| 20 Tablets/ Bottle | NDC 64679-921-01 |
| 60 Tablets/ Bottle | NDC 64679-921-02 |

Cefuroxime axetil tablets USP, 500 mg of cefuroxime as cefuroxime axetil, are white film-coated, round-shaped biconvex unscored tablets embossed with W922 on one side and plain on other side as follows:

20 Tablets/ Bottle NDC 64679-922-01

60 Tablets/ Bottle NDC 64679-922-02

**Store at 20° - 25°C (68° - 77°F). Replace cap securely after each opening.**

## **17 PATIENT COUNSELING INFORMATION**

### Allergic Reactions

Inform patients that cefuroxime axetil tablets is a cephalosporin that can cause allergic reactions in some individuals [see *Warnings and Precautions (5.1)*].

### Clostridium difficile-associated Diarrhea

Inform patients that diarrhea is a common problem caused by antibacterials, and it usually ends when the antibacterial is discontinued. Sometimes after starting treatment with antibacterials, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as 2 or more months after having taken their last dose of the antibacterial. If this occurs, advise patients to contact their physician as soon as possible.

### Crushing Tablets

Instruct patients to swallow the tablet whole, without crushing the tablet. Patients who cannot swallow the tablet whole should receive the oral suspension.

### Drug Resistance

Inform patients that antibacterial drugs, including cefuroxime axetil tablets, should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When cefuroxime axetil tablets are prescribed to treat a bacterial infection, inform patients that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by cefuroxime axetil tablets or other antibacterial drugs in the future.

Trademarks are the property of their respective owners.

### **Manufactured by:**

Wockhardt Limited

Plot No.B-15/2, M.I.D.C. Area, Waluj,

Aurangabad, Maharashtra,

India.

### **Distributed by:**

Wockhardt USA LLC.

20 Waterview Blvd.

Parsippany, NJ 07054

USA.

Rev.020915

## **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

DRUG: Cefuroxime Axetil

GENERIC: Cefuroxime Axetil

DOSAGE: Film-coated Tablets

ADMINSTRATION: Oral

NDC: 64679-921-01

STRENGTH: 250 mg

COLOR: White

SHAPE: Round

SCORE: no score

SIZE: 11 mm

IMPRINT: W;921

QTY: 20 Tablets

NDC 64679-921-01

# Cefuroxime Axetil Tablets, USP

**250 mg**

Rx only

20 Tablets



DRUG: Cefuroxime Axetil

GENERIC: Cefuroxime Axetil

DOSAGE: Film-coated Tablets

ADMINSTRATION: Oral

NDC: 64679-921-02

STRENGTH: 250 mg

COLOR: White

SHAPE: Round

SCORE: no score

SIZE: 11 mm

IMPRINT: W;921

QTY: 60 Tablets

NDC 64679-921-02

# Cefuroxime Axetil Tablets, USP

**250 mg**

Rx only

60 Tablets



DRUG: Cefuroxime Axetil

GENERIC: Cefuroxime Axetil

DOSAGE: Film-coated Tablets

ADMINSTRATION: Oral

NDC: 64679-922-01

STRENGTH: 500 mg

COLOR: White

SHAPE: Capsule

SCORE: no score

SIZE: 20 mm

IMPRINT: W;922

QTY: 20 Tablets

NDC 64679-922-01

# Cefuroxime Axetil Tablets, USP

500 mg

Rx only

20 Tablets

## CEFUROXIME AXETIL

cefuroxime axetil tablet, film coated

### Product Information

|                         |                         |                    |               |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:55648-920 |
| Route of Administration | ORAL                    |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| CEFUROXIME AXETIL (UNII: Z49QDT0J8Z) (CEFUROXIME - UNII:O1R9FJ93ED) | CEFUROXIME        | 125 mg   |

### Inactive Ingredients

| Ingredient Name                            | Strength |
|--|----------|
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48)    |          |
| DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)        |          |
| HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY) |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)           |          |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H)          |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)        |          |
| SHELLAC (UNII: 46N107B71O)                 |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)         |          |
| SMCC (UNII: B357PIG1IF)                    |          |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J)   |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)        |          |

### Product Characteristics

|          |               |              |          |
|----------|---------------|--------------|----------|
| Color    | WHITE (white) | Score        | no score |
| Shape    | ROUND (round) | Size         | 9mm      |
| Flavor   |               | Imprint Code | W;920    |
| Contains |               |              |          |

### Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:55648-920-01 | 20 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:55648-920-02 | 60 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA065166                               | 07/29/2005           |                    |

## CEFUROXIME AXETIL

cefuroxime axetil tablet, film coated

### Product Information

|                         |                         |                    |               |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:55648-921 |
| Route of Administration | ORAL                    |                    |               |

| Active Ingredient/Active Moiety                                     |  |                   |          |
|---|--|-------------------|----------|
| Ingredient Name   |  | Basis of Strength | Strength |
| CEFUROXIME AXETIL (UNII: Z49QDT0J8Z) (CEFUROXIME - UNII:O1R9FJ93ED) |  | CEFUROXIME        | 250 mg   |

| Inactive Ingredients                       |          |
|--|----------|
| Ingredient Name                            | Strength |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48)    |          |
| DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)        |          |
| HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY) |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)           |          |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H)          |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)        |          |
| SHELLAC (UNII: 46N107B71O)                 |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)         |          |
| SMCC (UNII: B357PIG1IF)                    |          |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J)   |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)        |          |

| Product Characteristics |               |              |          |
|-------------------------|---------------|--------------|----------|
| Color                   | WHITE (white) | Score        | no score |
| Shape                   | ROUND (round) | Size         | 11mm     |
| Flavor                  |               | Imprint Code | W;921    |
| Contains                |               |              |          |

| Packaging |                  |   |                      |                    |
|-----------|------------------|---|----------------------|--------------------|
| #         | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
| 1         | NDC:55648-921-01 | 20 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |
| 2         | NDC:55648-921-02 | 60 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |

| Marketing Information |  |                      |                    |
|-----------------------|--|----------------------|--------------------|
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA                  | ANDA065166                               | 07/29/2005           |                    |

| CEFUROXIME AXETIL                     |  |  |  |
|---------------------------------------|--|--|--|
| cefuroxime axetil tablet, film coated |  |  |  |

| Product Information     |                         |                    |               |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:55648-922 |
| Route of Administration | ORAL                    |                    |               |

| Active Ingredient/Active Moiety                                     |  |                   |          |
|---|--|-------------------|----------|
| Ingredient Name   |  | Basis of Strength | Strength |
| CEFUROXIME AXETIL (UNII: Z49QDT0J8Z) (CEFUROXIME - UNII:O1R9FJ93ED) |  | CEFUROXIME        | 500 mg   |

| Inactive Ingredients                       |          |
|--|----------|
| Ingredient Name                            | Strength |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48)    |          |
| DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)        |          |
| HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY) |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)           |          |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H)          |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)        |          |
| SHELLAC (UNII: 46N107B71O)                 |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)         |          |
| SMCC (UNII: B357PIG1IF)                    |          |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J)   |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)        |          |

| Product Characteristics |                   |              |          |
|-------------------------|-------------------|--------------|----------|
| Color                   | WHITE (white)     | Score        | no score |
| Shape                   | CAPSULE (capsule) | Size         | 20mm     |
| Flavor                  |                   | Imprint Code | W;922    |
| Contains                |                   |              |          |

| Packaging |           |                     |                      |                    |
|-----------|-----------|---------------------|----------------------|--------------------|
| #         | Item Code | Package Description | Marketing Start Date | Marketing End Date |

|   |                  |   |  |  |
|---|------------------|---|--|--|
| 1 | NDC:55648-922-01 | 20 in 1 BOTTLE; Type 0: Not a Combination Product |  |  |
| 2 | NDC:55648-922-02 | 60 in 1 BOTTLE; Type 0: Not a Combination Product |  |  |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA065166                               | 07/29/2005           |                    |

**Labeler** - Wockhardt Limited (650069115)

**Registrant** - Wockhardt Limited (650069115)

### Establishment

| Name              | Address | ID/FEI    | Business Operations   |
|-------------------|---------|-----------|---|
| Wockhardt Limited |         | 915122332 | analysis(55648-920, 55648-921, 55648-922) , manufacture(55648-920, 55648-921, 55648-922) , label(55648-920, 55648-921, 55648-922) , pack(55648-920, 55648-921, 55648-922) |

Revised: 9/2015

Wockhardt Limited