

AMMONIUM LACTATE- ammonium lactate cream
Padagis Israel Pharmaceuticals Ltd

Ammonium Lactate Cream, 12%*

Rx Only


For Dermatologic Use Only

Not for Ophthalmic, Oral or Intravaginal Use

DESCRIPTION

*Ammonium Lactate Cream, 12% is a formulation of 12% lactic acid neutralized with ammonium hydroxide, as ammonium lactate with a pH of 4.4 - 5.4. Ammonium Lactate Cream, 12% also contains cetyl alcohol, glycerin, glyceryl stearate, laureth-4, light mineral oil, magnesium aluminum silicate, methylcellulose, methylparaben, PEG-100 stearate, polyoxyl 40 stearate, propylene glycol, propylparaben, and purified water.

Lactic acid is a racemic mixture of 2-hydroxypropanoic acid and has the following structural formula:



CLINICAL PHARMACOLOGY

Lactic acid is an alpha-hydroxy acid. It is a normal constituent of tissues and blood. The alpha-hydroxy acids (and their salts) are felt to act as humectants when applied to the skin. This property may influence hydration of the stratum corneum. In addition, lactic acid, when applied to the skin, may act to decrease corneocyte cohesion. The mechanism(s) by which this is accomplished is not yet known.

An *in vitro* study of percutaneous absorption of ammonium lactate cream using human cadaver skin indicates that approximately 6.1% of the material was absorbed after 68 hours.

INDICATIONS AND USAGE

Ammonium Lactate Cream, 12% is indicated for the treatment of ichthyosis vulgaris and xerosis.

CONTRAINDICATIONS

Ammonium Lactate Cream, 12% is contraindicated in those patients with a history of hypersensitivity to any of the label ingredients.

WARNING

Use of this product should be discontinued if hypersensitivity to any of the ingredients is noted. Sun exposure to areas of the skin treated with ammonium lactate cream should be minimized or avoided (see **PRECAUTIONS**).

PRECAUTIONS

General -

For external use only. Stinging or burning may occur when applied to skin with fissures, erosions, or that is otherwise abraded (for example, after shaving the legs). Caution is advised when used on the face because of the potential for irritation. The potential for post-inflammatory hypo- or hyperpigmentation has not been studied.

Information for Patients

Patients using ammonium lactate cream should receive the following information and instructions:

1. This medication is to be used as directed by the physician, and should not be used for any disorder other than for which it was prescribed. It is for external use only. Avoid contact with eyes, lips, or mucous membranes.
2. Patients should minimize or avoid use of this product on areas of the skin that may be exposed to natural or artificial sunlight, including the face. If sun exposure is unavoidable, clothing should be worn to protect the skin.
3. This medication may cause stinging or burning when applied to skin with fissures, erosions, or abrasions (for example, after shaving the legs).
4. If the skin condition worsens with treatment, the medication should be promptly discontinued.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

The topical treatment of CD-1 mice with 12%, 21% or 30% ammonium lactate formulations for two years did not produce a significant increase in dermal or systemic tumors in the absence of increased exposure to ultraviolet radiation. The maximum systemic exposure of the mice in this study was 0.7 times the maximum possible systemic exposure in humans. However, a long-term photocarcinogenicity study in hairless albino mice suggested that topically applied 12% ammonium lactate cream enhanced the rate of ultraviolet light-induced skin tumor formation.

The mutagenic potential of ammonium lactate cream was evaluated in the Ames assay and in the mouse *in vivo* micronucleus assay, both of which were negative.

In dermal Segment I and III studies with ammonium lactate cream, there were no effects observed in fertility or pre- or post-natal development parameters in rats at dose levels of 300 mg/kg/day (1800 mg/m²/day), approximately 0.4 times the human topical dose

Pregnancy: Teratogenic Effects:

Pregnancy Category B - Animal reproduction studies have been performed in rats and rabbits at doses up to 0.7 and 1.5 times the human dose, respectively (600 mg/kg/day, corresponding to 3600 mg/m²/day in the rat and 7200 mg/m²/day in the rabbit) and have revealed no evidence of impaired fertility or harm to the fetus due to ammonium lactate cream. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, ammonium lactate cream should be used during pregnancy only if clearly needed.

Nursing Mothers -

Although lactic acid is a normal constituent of blood and tissues, it is not known to what extent this drug affects normal lactic acid levels in human milk. Because many drugs are excreted in human milk, caution should be exercised when ammonium lactate cream is administered to a nursing woman.

Pediatric Use -

The safety and efficacy of Ammonium Lactate Cream, 12% has been established in patients as young as 2 years old.

Geriatric Use -

Clinical studies of ammonium lactate cream, 12% did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between elderly and younger patients. In general, dose selection for an elderly patient should be cautious.

ADVERSE REACTIONS

In controlled clinical trials of patients with ichthyosis vulgaris, the most frequent adverse reactions in patients treated with ammonium lactate cream were rash (including erythema and irritation) and burning/stinging. Each was reported in approximately 10-15% of patients. In addition, itching was reported in approximately 5% of patients.

In controlled clinical trials of patients with xerosis, the most frequent adverse reactions in patients treated with ammonium lactate cream were transient burning, in about 3% of patients, stinging, dry skin and rash, each reported in approximately 2% of patients.

DOSAGE AND ADMINISTRATION

Apply to the affected areas and rub in thoroughly. Use twice daily or as directed by a physician.

HOW SUPPLIED

Ammonium Lactate Cream, 12% is available as follows:

385 g bottle (NDC 45802-**493**-26)

280 g carton (NDC 45802-**493**-83) containing two 140 g tubes (NDC 45802-**493**-98)

STORAGE

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Manufactured By Padagis
Minneapolis, MN 55427
www.padagis.com
Rev 12-21

Principal Display Panel

NDC 45802-493-26

Rx Only

Ammonium Lactate Cream, 12%

For Dermatologic Use Only

Not for Ophthalmic, Oral or Intravaginal Use

NET WT 385 g

NDC 45802-493-26

Rx Only

Ammonium Lactate Cream,

12%*

NET WT 385 g



CONTAINS: *Ammonium lactate (equivalent to 12% lactic acid), cetyl alcohol, glycerin, glyceryl stearate, laureth-4, light mineral oil, magnesium aluminum silicate, methylcellulose, methylparaben, PEG-100 stearate, polyoxyl 40 stearate, propylene glycol, propylparaben, and purified water.

**For dermatological use only.
Not for ophthalmic use.**

Usual dosage: Apply twice daily; see insert.

Keep out of the reach of children.

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

For lot number and expiration date, see bottle.

Manufactured By Padagis, Minneapolis, MN 55427
www.padagis.com

2204342 Rev 12-21 A 99R97 RC F2



3 45802-493-26 2

GTIN 00345802493262

IMPRINT AREA

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]

Lot [insert product's lot number]

Exp [insert product's expiration date]

AMMONIUM LACTATE

ammonium lactate cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:45802-493
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIUM LACTATE (UNII: 67M901L9NQ) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	12 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
CETYL ALCOHOL (UNII: 936JST6JCN)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PEG-100 STEARATE (UNII: YD01N1999R)	
POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
LAURETH-4 (UNII: 6HQ855798J)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-493-83	2 in 1 CARTON	01/25/2007	03/31/2025
1		140 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:45802-493-26	385 g in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	02/19/2007	
3	NDC:45802-493-98	140 g in 1 TUBE; Type 0: Not a Combination Product	08/16/2006	08/16/2006

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075774	08/16/2006	

Labeler - Padagis Israel Pharmaceuticals Ltd (600093611)

