

**BISACODYL- bisacodyl tablet, delayed release  
Bryant Ranch Prepack**

-----  
**Major 44-327**

***Active ingredient (in each tablet)***

Bisacodyl USP, 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness, and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

***Inactive ingredients***

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

***Questions or comments?***

**1-800-426-9391**

**HOW SUPPLIED**

Bisacodyl USP, 5 mg

NDC: 71335-2010-0: 25 Tablets in a BOTTLE

NDC: 71335-2010-1: 30 Tablets in a BOTTLE

NDC: 71335-2010-2: 2 Tablets in a BOTTLE

NDC: 71335-2010-3: 3 Tablets in a BOTTLE

NDC: 71335-2010-4: 4 Tablets in a BOTTLE

NDC: 71335-2010-5: 10 Tablets in a BOTTLE

NDC: 71335-2010-6: 90 Tablets in a BOTTLE

NDC: 71335-2010-7: 8 Tablets in a BOTTLE

NDC: 71335-2010-8: 100 Tablets in a BOTTLE

NDC: 71335-2010-9: 20 Tablets in a BOTTLE

Repackaged/Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504

**Bisacodyl 5mg Tablet**



GTIN 00371335201012  
Lot 208220  
Exp 7/5/2026  
SN 0123456789

Drug Facts	
<b>Active ingredient (in each tablet)</b> Bisacodyl 5 mg	<b>Purpose</b> Stimulant Laxative
<b>Uses</b> For relief of occasional constipation and irregularity - this product generally produces bowel movement in 6 to 12 hours	
<b>Warnings</b> Do not use if you cannot swallow without chewing. Ask a doctor before use if you have *stomach pain, nausea, vomiting *noticed a sudden change in bowel habits that lasts more than 2 weeks. When using this product *do not chew or crush tablets* *do not use within 1 hour after taking an antacid or milk *it may cause stomach discomfort, faintness and cramps. Stop use and ask a doctor if * you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. *you need to use a laxative for more than 1 week. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Other Information</b> *Tamper Evident: Do not use if imprinted seal under cap is missing or broken *Store at room temperature 15°-30°C (59°-86°F) *Avoid excessive humidity * package not child resistant	
<b>Directions</b> *Take with a glass of water *Adults and children 12 years and older, take 1 to 3 tablets in a single daily dose *Children 12 and under: ask a doctor	
<b>Inactive Ingredients</b> Acacia, ammonium hydroxide, calcium carbonate, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, iron oxide black, lactose, magnesium stearate, methylparaben, PEG, polydextrose, polyvinyl acetate phthalate, propylparaben, propylene glycol, povidone, shellac, simethicone, silica, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate, wax.	

**NDC 71335-2010-1**

**Bisacodyl Tablets, USP**

**5 mg**

**30 Tablets**



**Repackaged by:** Bryant Ranch Prepack, Inc. Burbank, CA 91504 USA  
**Manufactured by:** LNK International, Inc.



**BISACODYL**

bisacodyl tablet, delayed release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71335-2010(NDC:0904-6748)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>AMMONIA</b> (UNII: 5138Q19F1X)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERROSO FERRIC OXIDE</b> (UNII: XMOM87F357)	

<b>ANHYDROUS LACTOSE</b> (UNII: 3S5Y5LH9PMK)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM ALGINATE</b> (UNII: C269C4G2ZQ)
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	5
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71335-2010-0	25 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	
2	NDC:71335-2010-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	03/03/2022	
3	NDC:71335-2010-2	2 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	
4	NDC:71335-2010-3	3 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	
5	NDC:71335-2010-4	4 in 1 BOTTLE; Type 0: Not a Combination Product	01/03/2022	
6	NDC:71335-2010-5	10 in 1 BOTTLE; Type 0: Not a Combination Product	01/18/2022	
7	NDC:71335-2010-6	90 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	
8	NDC:71335-2010-7	8 in 1 BOTTLE; Type 0: Not a Combination Product	01/12/2022	
9	NDC:71335-2010-8	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	
10	NDC:71335-2010-9	20 in 1 BOTTLE; Type 0: Not a Combination Product	04/03/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	12/01/2018	

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

## Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(71335-2010) , RELABEL(71335-2010)

Revised: 7/2024

Bryant Ranch Prepack