

**MUCUS RELIEF DAYTIME SEVERE COLD NIGHTTIME COLD AND FLU MAXIMUM STRENGTH- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl, guaifenesin**  
**TARGET Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients in Daytime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

**Active ingredients in Nighttime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCL 5 mg

**Purpose for Daytime**

**Pain reliever/fever reducer**

Cough suppressant

Expectorant

Nasal decongestant

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

## **DAYTIME**

- temporarily relieves these common cold and flu symptoms
  - headache
  - nasal congestion
  - sore throat
  - cough
  - minor aches and pains
  - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
  - temporarily reduces fever

## **NIGHTTIME**

- temporarily relieves these common cold and flu symptoms
  - cough
  - headache
  - minor aches and pains
  - sore throat
  - nasal congestion
  - runny nose and sneezing
  - controls cough to help you get to sleep
  - temporarily reduces fever

## **Warnings**

### **DAYTIME and NIGHTTIME**

**Liver warning:** These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

### **DAYTIME and NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

#### **DAYTIME**

- liver disease
- diabetes
- high blood pressure
- heart disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

#### **NIGHTTIME**

- liver disease
- diabetes
- high blood pressure
- heart disease
- glaucoma
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

#### **DAYTIME**

taking the blood thinning drug warfarin

#### **NIGHTTIME**

taking the blood thinning drug warfarin

taking sedatives or tranquilizers

### **When using this product,**

#### **DAYTIME**

**do not use more than directed**

#### **NIGHTTIME**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

#### **DAYTIME and NIGHTTIME**

- nervousness, dizziness, or sleeplessness occurs
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

#### **DAYTIME and NIGHTTIME**

ask a health professional before use.

### **Keep out of reach of children.**

#### **DAYTIME and NIGHTTIME**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### **DAYTIME**

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

#### **NIGHTTIME**

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels (Daytime and Nighttime) in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

### **Other information**

#### **DAYTIME and NIGHTTIME**

- store between 20-25°C (68-77°F)

- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

## **Inactive ingredients**

### **DAYTIME**

FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide

### **NIGHTTIME**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide

## **Question or comments?**

Call 1-800-910-6874

## **Principal Display Panel**

### **DAYTIME**

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Day Severe Cold\*\*\*

maximum strength

daytime severe cold

acetaminophen (pain reliever / fever reducer)

dextromethorphan HBr (cough suppressant)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

relieves aches, fever and sore throat

controls cough

relieves nasal and chest congestion

thins and loosens mucus

AGES 12+ YEARS

SOFTGELS\*\* (\*\*LIQUID-FILLED CAPSULES)

### **NIGHTTIME**

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Cold & Flu\*\*\*

maximum strength

nighttime

Cold & Flu

acetaminophen (pain reliever / fever reducer)

dextromethorphan HBr (cough suppressant)

doxylamine succinate (antihistamine)

phenylephrine HCL (nasal decongestant)

relieves aches, fever and sore throat

controls cough

relieves nasal congestion

relieves runny nose and sneezing

AGES 12+ YEARS

SOFTGELS\*\* (\*\*LIQUID-FILLED CAPSULES)

\*\*\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Severe Cold and Maximum Strength Mucinex® Fast-Max® Night Cold & Flu Liquid Gels

**TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by Target Corporation

Minneapolis, MN 55403

TM & ©2019 Target Brands, Inc.

**Product Label**

nighttime cold and flu (continued)

**Drug Facts (continued)**

- cough that occurs with too much phlegm (mucus)
- Ask a doctor or pharmacist before use if you are:
  - taking the blood thinning drug warfarin
  - taking sedatives or tranquilizers

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Directions**

- do not take more than directed (see Overdose warning)
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide

**Questions or comments?**  
Call 1-800-919-8874

daytime severe cold

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg..... Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg..... Cough suppressant  
Guaifenesin 200 mg..... Expectorant  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Purposes**

**Uses**

- temporarily relieves these common cold and flu symptoms
  - headache
  - nasal congestion
  - sore throat
  - cough
  - minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and make coughs more productive
- temporarily reduces fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Directions**

- do not take more than directed (see Overdose warning)
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

**Questions or comments?**  
Call 1-800-919-8874

daytime severe cold (continued)

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Directions**

- do not take more than directed (see Overdose warning)
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

**Questions or comments?**  
Call 1-800-919-8874

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Day Severe Cold Liquid Gels\*\*\*

maximum strength  
**daytime severe cold**

**acetaminophen** (pain reliever/fever reducer)  
**dextromethorphan HBr** (cough suppressant)  
**guaifenesin (expectorant)**  
**phenylephrine HCl** (nasal decongestant)

relieves aches, fever and sore throat  
controls cough  
relieves nasal and chest congestion  
thins and loosens mucus



ACTUAL SIZE

AGES 12+ YEARS

16 SOFTGELS\*\* (\*\*LIQUID-FILLED CAPSULES)  
24 TOTAL SOFTGELS

NDC 11673-906-24

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Cold & Flu Liquid Gels\*\*\*

maximum strength  
**nighttime cold and flu**

**acetaminophen** (pain reliever/fever reducer)  
**dextromethorphan HBr** (cough suppressant)  
**doxylamine succinate** (antihistamine)  
**phenylephrine HCl** (nasal decongestant)

relieves aches, fever and sore throat  
controls cough  
relieves nasal congestion  
relieves runny nose and sneezing



ACTUAL SIZE

AGES 12+ YEARS

8 SOFTGELS\*\* (\*\*LIQUID-FILLED CAPSULES)

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg..... Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg..... Cough suppressant  
Doxylamine succinate 6.25 mg..... Antihistamine  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms
  - cough
  - headache
  - minor aches and pains
  - sore throat
  - nasal congestion
  - runny nose and sneezing
- controls cough to help you get to sleep
- temporarily reduces fever

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Directions**

- do not take more than directed (see Overdose warning)
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

**Inactive ingredients** FD&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

**Questions or comments?**  
Call 1-800-919-8874

094 14 0126 R02 C-000703-01-089

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2019 Target Brands, Inc.



3 59726 51824 6 PLD-C584A F0006280

**PARENTS:**  
Learn about how medicine abuse.org

**TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Lot / Exp:  
Product of:

**TARGET Daytime Maximum Strength Severe Cold Nighttime Maximum Strength Cold and Flu**

**MUCUS RELIEF DAYTIME SEVERE COLD NIGHTTIME COLD AND FLU MAXIMUM STRENGTH**  
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl, guaifenesin kit

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-906

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-906-24	1 in 1 KIT; Type 0: Not a Combination Product	09/30/2019	08/29/2025

Quantity of Parts			
Part #	Package Quantity		Total Product Quantity
Part 1	8 BLISTER PACK		8
Part 2	16 BLISTER PACK		16

# NIGHTTIME COLD AND FLU MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hcl capsule

## Product Information

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	42A
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 CARTON		
-		1 in 1 BLISTER PACK; Type 0: Not a Combination		



Product

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2019	

## Part 2 of 2

### MUCUS RELIEF DAYTIME SEVERE COLD MAXIMUM STRENGTH

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl capsule

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm

<b>Flavor</b>		<b>Imprint Code</b>	12A
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	09/30/2019	08/29/2025	

<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	09/30/2019	08/29/2025	

**Labeler** - TARGET Corporation (006961700)