

TERRAMYCIN 50- oxytetracycline powder
Phibro Animal Health

Terramycin® 50
(oxytetracycline)
TYPE A MEDICATED ARTICLE

Active Drug Ingredient:

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to
oxytetracycline hydrochloride 50 g/lb

CAUTION:

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

CAUTION: For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT OR BELOW 25°C (77°F), EXCURSIONS PERMITTED UP TO 40°C (104°F)
SEE BACK PANEL FOR COMPLETE MIXING DIRECTIONS
USE DIRECTIONS, AND WARNINGS

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

Mixing and Use Directions

Thoroughly mix the amount of this Type A medicated article according to the directions below with at least an equal amount by weight of feed formula ingredients prior to blending into a complete feed.

Indications for Use	Oxytetracycline Amount	lb of Terramycin 50/ton
----------------------------	-------------------------------	--------------------------------

CHICKENS		
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	100-200 g/ton Feed continuously for 7-14 days	2-4
Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline	400 g/ton Feed continuously for 7-14 days	8
Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to oxytetracycline	500 g/ton Feed continuously for 5 days	10
WARNING: At 500 g/ton level, withdraw 24 hours before slaughter. Zero-day withdrawal period for lower use levels. Do not use in feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues. Do not administer to chickens producing eggs for human consumption.		
TURKEYS		
Control of hexamitiasis caused by <i>Hexamitameleagridis</i> susceptible to oxytetracycline	100 g/ton Feed continuously for 7-14 days	2
Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to oxytetracycline	200 g/ton Feed continuously for 7-14 days	4
Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline	25 mg/lb of body weight daily Feed continuously for 7-14 days	16.7 ¹
WARNING: At 200 g/ton use level or higher, withdraw 5 days before slaughter. Zero-day withdrawal period for lower use levels. Do not administer to turkeys producing eggs for human consumption.		
SWINE		
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurellamultocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	10 ²
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for not more than 14 days	10 ²
CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE		
For growing cattle (over 400 lb) for the	75 mg/head/day	1.5 ³

reduction of the incidence of liver abscesses	Feed continuously	
Prevention and treatment of the early stages of shipping fever complex. (Feed 3-5 days before and after arrival in feedlots.)	0.5-2.0 g/head/day	10-40 ³
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	100 ⁴
SHEEP		
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	24 ⁵
WARNING: 5-day withdrawal before slaughter.		
¹ If bird weighs 10 lb, consuming 0.6 lb of complete feed per day ² Pig consuming 4 lb of complete feed per 100 lb body weight per day ³ Include in feed supplement based on consumption of 2 lb of supplement per head per day ⁴ If animal weighs 500 lb, consuming 2 lb of supplement per head per day ⁵ If lamb weighs 60 lb, consuming 1 lb of supplement per head per day		

RESTRICTED DRUG (CALIFORNIA) - USE ONLY AS DIRECTED

NOT FOR HUMAN USE

Terramycin is a registered trademark of Pfizer, Inc., licensed to Phibro Animal Health Corporation, for Oxytetracycline
PHIBRO and PHIBRO logo design are trademarks owned by Phibro Animal Health Corporation or its affiliates
Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.7 kg)

Approved by FDA under NADA # 095-143

8842000

101-9009-10

Principal Display Package



TERRAMYCIN 50

oxytetracycline powder

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-7200
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTETRACYCLINE HYDROCHLORIDE (UNII: 4U7K4N52Z M) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	50 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
RICE BRAN (UNII: R60QEP13IC)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-7200-0	22.7 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA095143	03/31/2010	

Labeler - Phibro Animal Health (006989008)**Registrant** - Phibro Animal Health (006989008)

Revised: 7/2023

Phibro Animal Health