ASPIRIN- as pirin tablet CARDINAL HEALTH

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Aspirin 81mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommeded by your doctor. Because of
 its delayed action, this product will not provide fast relief of headaches or other symptoms
 needing immediate relief.
- ask your doctor about other uses for this product

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- any new symptoms occur
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs

These could be signs of a serious illness.

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

• drink a full glass of water with each dose

adults and children 12 years take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours and over unless directed by a doctor children under 12 years consult a doctor

Other information

- store at room temperature
- save carton for full direction and warnings

Inactive ingredients

croscarmellose sodium, D&C yellow# 10 Lake, FD&C yellow# 6, hypromellose, lactose, methcrylic acid copolymer, microcrystaline cellulose, silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethlyl citrate

Questions or comments?

Adverse drug event call: (866) 562-2756

NDC 49781-097-52

LEADER

[†]Compare to <code>Bayer</code> low dose aspirin active ingredient

LEADER


□#□ASPIRIN REGIMEN

Aspirin 81 mg

Pain Reliever (NSAID)

Low Dose Safety Coated

120 ENTERIC COATED TABLETS



ASPIRIN

aspirin tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:49781-097

Route of Administration ORAL

Active Ingredient/Active Moiety

l	Ingredient Name	Basis of Strength	Strength
ı	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	

Product Characteristics				
Color	yello w	Score	no score	
Shape	ROUND	Size	8mm	
Flavor		Imprint Code	EA8	
Contains				

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:49781-097-52	1 in 1 CARTON	09/20/2014		
1		120 in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:49781-097-58	300 in 1 BOTTLE; Type 0: Not a Combination Product	09/20/2014		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part343	09/20/2014		

Labeler - CARDINAL HEALTH (097537435)

Registrant - Pharbest Pharmaceuticals, Inc. (557054835)

Revised: 11/2017 CARDINAL HEALTH