

NAUZENE - sodium citrate syrup
Alva-Amco Pharmcal Companies, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 15-mL tablespoon)

Dextrose (glucose) 4.35 g.....anti-nausea

Levulose (fructose) 4.17 g.....anti-nausea

Sodium citrate dihydrate, 0.921 g.....antacid

Uses

For the relief of nausea associated with upset stomach, including that due to overindulgence in food or drink.

Warnings

Do not use if you have Hereditary Fructose Intolerance (HFI). This product contains fructose.

Ask a doctor before use if you

- you have diabetes because this product contains sugar
- are on a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking any other medications. Antacids may interact with certain prescription drugs.

When using this product do not take more than 6 does in a 24-hour period.

Stop use and ask a doctor if nausea lasts more than two weeks or recurs frequently.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Read all package directions and warnings before use and use only as directed.
- Adults: 1 - 2 tablespoons (15 - 30 mL).
- Children: Consult a doctor for appropriate dosage.
- Dosage may be repeated after 30 minutes not to exceed 6 doses in a 24-hour period unless advised by a doctor.
- For maximum effectiveness never dilute Nausea or drink fluids of any kind immediately before or after taking Nausea.
- Nausea Liquid is intended for use by normally healthy persons only.
- Persons under 18 years of age should use only as directed by a doctor.

Other information

- Sodium content 193 mg/tablespoon (15 mL).
- Store at room temperature.
- ** Contents sealed: Each bottle of Nauzene Liquid has a tamper evident seal around the cap and a safety seal under the cap. Do not use if either of these seals appears broken. NOTE: Nauzene is not intended as a substitute for a balanced nutritional diet or as an electrolyte replenishment.
- Serious side effects associated with the use of this product may be reported to the phone number provided below.

Inactive ingredients

Corn syrup, FDC Red No. 40 lake, flavor, fructose syrup, glycerin, methylparaben, phosphoric acid, purified water.

Questions? 1-800-792-2582

72-4R
0209

Nauzene®
Liquid

Wild
Cherry
Flavor

Drug Facts

Active ingredients Purpose (in each 15-mL tablespoon)

Dextrose (Glucose) 4.35 g Anti-nausea
Levulose (Fructose) 4.17 g Anti-nausea
Sodium citrate dihydrate, Antacid
0.921 g

Uses For the relief of nausea associated with upset stomach, including that due to overindulgence in food or drink.

Warnings

Do not use if you have Hereditary Fructose Intolerance (HFI). This product contains fructose.

Ask a doctor before use if you ■ have diabetes because this product contains sugar ■ are on a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking any other medications. Antacids may interact with certain prescription drugs.

When using this product do not take more than 6 doses in a 24-hour period.

Stop use and ask a doctor if nausea lasts more than two weeks or recurs frequently.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Read all package directions and warnings before use and use only as directed. ■ Adults: 1 - 2 tablespoons (15 - 30 mL). ■ Children: Consult a doctor for appropriate dosage. ■ Dosage may be repeated after 30 minutes not to exceed 6 doses in a 24-hour period unless advised

Drug Facts (continued)

Directions (continued)

by a doctor. ■ For maximum effectiveness never dilute Nauzene or drink fluids of any kind immediately before or after taking Nauzene. ■ Nauzene Liquid is intended for use by normally healthy persons only. ■ Persons under 18 years of age should use only as directed by a doctor.

Other Information: ■ Sodium content 193 mg/tablespoon (15 mL)
■ Store at room temperature

■ ****Contents sealed:** Each bottle of Nauzene Liquid has a tamper evident seal around the cap and a safety seal under the cap. Do not use if either of these seals appears broken. NOTE: Nauzene is not intended as a substitute for a balanced nutritional diet or as an electrolyte replenishment. ■ Serious side effects associated with the use of this product may be reported to the phone number provided below.

Inactive Ingredients Corn syrup, FD&C Red No. 40 lake, flavor, fructose syrup, glycerin, methylparaben, phosphoric acid, purified water.

Questions? 1-800-792-2582

*ALVA-AMCO trade name for sodium citrate dihydrate is **Electrolyte-S10™**.

KEEP TIGHTLY CLOSED WHEN NOT IN USE. STORE AWAY FROM EXCESSIVE HEAT AND COLD.

www.NAUZENE.com



ALVA-AMCO Pharmaceutical Cos., Inc.
Niles, IL 60714, U.S.A.

©1999 - 2009 ALVA. ®Reg. U.S. Patent Office

4

with **Electrolyte-S10™**

Nauzene®

Liquid

For Nausea

with **Electrolyte-S10™**

Wild Cherry Flavor

Nauzene®

Liquid

For Nausea

For Relief of Upset Stomach Including Nausea

- Prompt Relief
- No Aspirin
- No Salicylates



4 MINUTE FORMULA™

** Contents Sealed
4 fl(oz) LIQUID (118 ml)

with **Electrolyte-S10™**

Wild Cherry Flavor

Nauzene®

Liquid

For Nausea

Great Tasting
FOR THE WHOLE FAMILY

- No chalky taste
- No medicinal flavor
- No bitter aftertaste

† In laboratory tests, **Nauzene** reaches 99% of its neutralizing ability within **4 minutes**.



LOT NO.:

EXP.:

PROD. NO: 72004



0 72959 72004 9

456
09 10 11 12
789ABC
123456
789ABC
123456
789ABC
123456
789ABC

NAUZENE

sodium citrate syrup

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:52389-172

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	TRISODIUM CITRATE DIHYDRATE	.921 g in 15 mL
Dextrose (UNII: IY9XDZ35W2) (Dextrose - UNII:IY9XDZ35W2)	Dextrose	4.35 g in 15 mL
Fructose (UNII: 6YSS42VSEV) (Fructose - UNII:6YSS42VSEV)	Fructose	4.17 g in 15 mL

Inactive Ingredients

Ingredient Name	Strength
corn syrup (UNII: 9G5L16BK6N)	
FD&C Red No. 40 (UNII: WZB9127XOA)	
High fructose corn syrup (UNII: XY6UN3QB6S)	
Glycerin (UNII: PDC6A3C0OX)	
Methylparaben (UNII: A2I8C7HI9T)	
Phosphoric acid (UNII: E4GA8884NN)	
Water (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY (flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52389-172-04	118 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part357	02/15/2000	

Labeler - Alva-Amco Pharmacial Companies, Inc. (042074856)

Revised: 5/2010

Alva-Amco Pharmacial Companies, Inc.