

**FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE -  
fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended  
release**

**PUBLIX SUPER MARKETS INC**

-----  
**Fexofenadine Hydrochloride and Pseudoephedrine Hydrochloride Extended-Release Tablets, 60  
mg/120 mg**

***Drug Facts***

<b><i>Active ingredients (in each extended-release tablet)</i></b>	<b><i>Purpose</i></b>
Fexofenadine HCl, USP 60 mg	Antihistamine
Pseudoephedrine HCl, USP 120 mg	Nasal Decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids

- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

**Other information**

- do not use if carton is opened or if individual blister units are torn or opened
- store between 68° to 77°F (20° to 25°C)
- USP dissolution test is pending.

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.

DISTRIBUTED BY  
 PUBLIX SUPER MARKETS, INC.,  
 3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811

**PRINCIPAL DISPLAY PANEL - 30 Tablet Blister Pack Carton**

NDC 56062-999-30

NON-DROWSY  
 ORIGINAL PRESCRIPTION STRENGTH

**allergyreliefD**

**FEXOFENADINE HCl 60 mg/ANTI-HISTAMINE  
 PSEUDOEPHEDRINE HCl 120 mg/NASAL DECONGESTANT  
 EXTENDED-RELEASE TABLETS, USP**

ALLERGY & CONGESTION • INDOOR & OUTDOOR ALLERGIES

**12-hour relief of:**

- Nasal & sinus congestion due to colds or allergies
- Sneezing; runny nose; itchy, watery eyes & itchy nose or throat due to allergies

ACTUAL SIZE

**30**

EXTENDED-RELEASE  
TABLETS

\*Compare to the active ingredients  
in Allegra-D<sup>®</sup>

**DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN**

### Drug Facts (continued)

#### Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

#### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Drug Facts (continued)

#### Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### Other information

- do not use if carton is opened or if individual blister units are torn or opened
- store between 68° to 77°F (20° to 25°C)
- USP dissolution test is pending.

#### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, ethyl cellulose, ferric oxide yellow, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid.



0519



publix.

COMPLETE SATISFACTION OR YOUR MONEY BACK

PUBLIC GUARANTEE:

1-888-267-3037

3300 PUBLIC CORPORATE PARKWAY, LAKELAND, FL 33811 publix.com

PUBLIC SUPER MARKETS, INC.

DISTRIBUTED BY

\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Allegra-D®.

FEXOFENADINE HCl 60 mg/ANTIHISTAMINE  
PSEUDOEPHEDRINE HCl 120 mg/NASAL DECONGESTANT  
EXTENDED-RELEASE TABLETS, USP



NDC 56062-899-30

NON-DROWSY  
ORIGINAL PRESCRIPTION STRENGTH

**allergyreliefD**

**FEXOFENADINE HCl 60 mg/ANTIHISTAMINE  
PSEUDOEPHEDRINE HCl 120 mg/NASAL DECONGESTANT**

# EXTENDED-RELEASE TABLETS, USP

## ALLERGY & CONGESTION • INDOOR & OUTDOOR ALLERGIES

### 12-hour relief of:

- Nasal & sinus congestion due to colds or allergies
- Sneezing; runny nose; itchy, watery eyes & itchy nose or throat due to allergies



ACTUAL SIZE

# 30

 EXTENDED-RELEASE TABLETS

\*Compare to the active ingredients in Allegra-D®

**DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN**

### Drug Facts

#### Active ingredients (in each extended-release tablet) Purpose

Fexofenadine HCl, USP 60 mg.....Antihistamine  
Pseudoephedrine HCl, USP 120 mg.....Nasal Decongestant

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages


### Drug Facts (continued)


- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

#### Warnings

##### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing





## FEXOFENADINE HYDROCHLORIDE AND PSEUDOEPHEDRINE HYDROCHLORIDE

fexofenadine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:56062-999
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FEXOFENADINE HYDROCHLORIDE</b> (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ETHYLCELLULOSE, UNSPECIFIED</b> (UNII: 7Z8S9VYZ4B)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	WHITE, YELLOW	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (bilayer)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	724
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-999-30	1 in 1 CARTON	04/01/2019	
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090818	04/01/2019	

**Labeler** - PUBLIX SUPER MARKET S INC (006922009)

**Establishment**

Name	Address	ID/FEI	Business Operations
Sun Pharmaceutical Industries Limited		650445203	ANALYSIS(56062-999) , MANUFACTURE(56062-999)

Revised: 5/2019

PUBLIX SUPER MARKET S INC