IODINE TOPICAL SOLUTION - povidone-iodine solution Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

ACTIVE INGREDIENT

Povidone Iodine,10% w/v (equivalent to 1% titratable iodine)

PURPOSE

Antiseptic

USE

First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns

WARNINGS

For external use only.

Do not

- use in the eyes
- use on individuals who are allergic or sensitive to iodine
- apply over large areas of the body

Discontinue use and ask a doctor

- if condition persists or gets worse
- for use longer than 1 week

Ask a doctor in case of

- deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- Clean the affected area
- Apply a small amount of this product to the area 1-3 times daily
- May be covered with a sterile bandage

• If bandaged, let dry first

OTHER INFORMATION

Store at room temperature 15° - 30° C (59° - 86° F)

INACTIVE INGREDIENTS

citric acid, glycerin, nonoxynol-8, purified water, sodium hydroxide

LABEL INFORMATION

NDC 41250-677-94

meijer

Iodine Topical Solution

povidone iodine USP, 10%

TOPICAL ANTISEPTIC
HELPS PREVENT INFECTION IN MINOR CUTS, SCRAPES AND BURNS
WATER SOLUABLE


MEIJER PHARMACIST RECOMMENDED

COMPARE TO BETADINE® active ingredient*

8 FL OZ (236 mL)

DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544

*This product is not manufactured of distributed by Purdue Frederick, owner of the registered trademark, Betadine®.



Drug Facts

Purpose Active ingredient

Povidone lodine USP, 10% w/v (equivalent to

1% titratable iodine) Antiseptic

First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns

Warnings For external use only.

- use in the eyes
 use on individuals who are allergic or sensitive to iodine
- · apply over large areas of the body

Discontinue use and ask a doctor

 if condition persists or gets worse · for use longer than 1 week

Ask a doctor in case of

- · deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children, If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- · Clean the affected area
- · Apply a small amount of this product to the area 1-3 times
- May be covered with a sterile bandage
 • If bandaged, let dry first

Other information

Store at room temperature: 15° - 30° C (59° - 86° F)

Inactive ingredients citric acid, glycerin, nonoxynol-8, purified water, sodium hydroxide

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark, Betadine®.

Rev. A

DIST. BY MEIJER DISTRIBUTION, INC **GRAND RAPIDS, MI 49544**





IODINE TOPICAL SOLUTION

povidone-iodine solution

Product Information

HUMAN OTC DRUG NDC:41250-677 Product Type Item Code (Source)

TOPICAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name **Basis of Strength** Strength povidone-io dine (UNII: 85H0 HZU99M) (povidone-io dine - UNII:85H0 HZU99M) po vido ne-io dine .10 mg in 1 mL

Inactive Ingredients		
Ingredient Name	Strength	
citric acid monohydrate (UNII: 2968PHW8QP)		
glycerin (UNII: PDC6A3C0OX)		

sodium hydroxide (UNII: 55X04QC32I)	
water (UNII: 059QF0KO0R)	

l	Packaging			
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:41250-677-94	236 mL in 1 BOTTLE		

Marketing Inform			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	12/02/2004	

Labeler - Meijer Distribution Inc (006959555)

Revised: 12/2009 Meijer Distribution Inc