

CHERRY THROAT DROPS- benzocaine lozenge
GRANDALL DISTRIBUTING, LLC

Grandall (as PLD) - Benzocaine 15 mg Menthol 3.6 mg Cherry Throat drops (48201-012)

Active ingredients (in each lozenge)

Benzocaine 15 mg

Menthol 3.6 mg

Purpose

Oral anesthetic

Oral anesthetic

Uses temporarily relieves these symptoms:

occasional minor irritation, pain, sore mouth and sore throat

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

Stop use and ask doctor if

- sore throat is severe or irritation, pain or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 5 years and older - allow 1 lozenge to dissolve slowly in mouth. May be repeated every 2 hours as necessary or as directed by a doctor

children under 5 years - do not use

Inactive ingredients: Corn starch, FD&C red no. 40, glucose syrup, glycerin, medium chain triglycerides, natural and artificial flavors, propylene glycol, soybean oil, sucrose and water.

Questions? Or to report an adverse event call 1-800-245-2898. Monday - Friday, 9AM to 4PM EST

Angidol

SORE THROAT

cherry

Menthol/Benzocaine Oral Anesthetic

NDC 48201-012-18

Angidol

Menthol/Benzocaine Oral Anesthetic

- **Temporarily relieves minor sore throat pain and irritation**
- **Fast and long acting**
- **Oral anesthetic**



18 LOZENGES

Gluten-Free

PN-320228



Distributed by:
GRANDALL ORAL CO., INC.
Shawnee, CA 95094-2707
1-800-394-6442
www.grandall.com



**No Ink or Coating
Lot/Exp Info**

No Ink or Coating

No Ink or Coating

Drug Facts (continued)
Inactive ingredients Com starch, FD&C Red #40, glucose syrup, glycerin, medium chain triglycerides, natural & artificial flavors, propylene glycol, soybean oil, sucrose and water.
Questions? Call 1-800-245-2898, Monday through Friday, 9AM - 4PM EST.
 This product is not manufactured or distributed by Beckett Benzocaine owner of the registered trademark Cepacol.
 WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BUSTERS IS OPEN OR DAMAGED.

Drug Facts
Active ingredients (in each lozenge)
 Benzocaine 16 mg
 Menthol 3.6 mg
Purpose
 Oral anesthetic
Uses temporarily relieves these symptoms:
 ■ occasional minor irritation, pain, sore mouth and sore throat
Warnings
 Methemoglobinemia warning: a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis), headache, rapid heart rate, ■ shortness of breath, ■ dizziness or lightheadedness, ■ fatigue or lack of energy.
 Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.
 Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.
 Do not use ■ in children under 2 years of age ■ for teething ■ in children under 2 years of age
 Stop use and ask doctor if ■ sore throat is severe or irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.
Directions adults and children allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor.
 children under 5 years do not use

Other information ■ store at room temperature ■ protect from moisture

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:48201-012
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	3.6 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	17mm
Flavor	CHERRY	Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:48201-012-18	18 in 1 BOX; Type 0: Not a Combination Product	08/07/2018	



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/07/2018	

Labeler - GRANDALL DISTRIBUTING, LLC (044428324)

Revised: 1/2024

GRANDALL DISTRIBUTING, LLC