FIBER LAX- calcium polycarbophil tablet, film coated PD-Rx Pharmaceuticals, Inc.

Drug Facts

Active ingredient (in each captab)

Calcium polycarbophil 625 mg

Purpose

Bulk-Forming Laxative (equivalent to polycarbophil 500 mg)

Uses

- relieves occasional constipation
- generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Do not use

longer than one week

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that lasts over two weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take Fiber-Lax two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding
- you fail to have a bowel movement after using this product.

These may indicate a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Take Fiber-Lax with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.

- do not take more than 4 doses in 24 hours
- adults & children 12 years and over: 2 captabs 1 to 4 times a day
- children under 12 years: ask a doctor

Other information

- each captab contains: calcium 140 mg
- store below 30°C (86°F) do not refrigerate protect from humidity

Inactive ingredients

Caramel, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl sulfate and stearic acid

Questions or comments?

Call 1-800-645-2158

TAMPER EVIDENT: DO NOT USE IF BLISTER SEAL IS BROKEN OR DAMAGED

*Rugby Laboratories, Inc. is not affiliated with the owner of the trademark FiberCon®.

Distributed by: Rugby Laboratories

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152

www. Rugbylaboratories.com

16 HOW SUPPLIED

Fiber-Lax Tablets are available as follows:

500 mg: Off White, oval, scored tablet, debossed "CPC 339". Available in bottles of 30 captabs (NDC 55289-949-30) Fiber-Lax

Drug Facts	Drug Facts (continued)	NDC 55289-949-30		
Active Ingredient Purpose (in each captab) Calcium polycarbophil 625mg (equivalent to polycarbophil 500 mg)	Directions Take Fiber-Lax winth at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough fiquid may cause choking. See choking warning.			
USES • relieves oaccasional constipation • generally produces bowel movement in 12 to 72 hours	do not take more than 4 doses in 24 hours adults & children 12 years and over: 2 captabs 1 to 4 times a day children under 12 years: ask a doctor	Fiber-Lax		
Warning: Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.		Bulk-Forming Laxative		
chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention. Do not use longer than one week	Inactive Ingredients caramé, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl suifate and stearic add	Promotes Regularity Non-Habit Forming 55280 94930		
Ask a doctor before use if you have • abdominal pain, nausea, or vomiting • noticed a sudden change in bowel habits that lasts over two weeks	sodium lauryl sulfate and stearic and Questions or comments? 1-800-645-2158	500 mg Marketed and Packaged By		
Ask a doctor or pharmacist before use if you are taking any other drug. Take Fiber-Lax two or more hours before or after other drugs. Laxatives may affect how other drugs work.	FIL: FOR GTIN: 00355289949305	PD-Rx Pharmaceuticals, Inc Oklahoma City, OK 73127		
Stop use and ask a doctor if • you have rectal bleeding • you fail to have a bowel movement after using this product. These may indicate a serious condition.	SNO: K20C890005	1-405-942-3040 v.8.19.0		
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	EXP: 11/2022 LOT: K20C89	30 Captabs TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL IS BROKEN OR MISSING FROM BOTTLE.		

film coated				
HUMAN OTC DRUG	Item Code (Source)	NDC:55289-949	9(NDC:0	536-4306)
ORAL				
-				
Ingredient Name Basis of Stre				
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL -CALCIUMUNII: W25LM17A4W)POLYCARBOPH			-	625 mg
Ingredient Name				
CARAMEL (UNII: T9D99G2B1R)				
(UNII: 2S7830E561)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
097M6I30)				
IE (UNII: OP1R32D61U)			
87QVP)				
	ORAL Moiety redient Name : 8F049NKY49) (POLY0 Ingredient Na (UNII: 2S7830E561) /3WO) 097M6I30)	HUMAN OTC DRUG Item Code (Source) ORAL Item Code (Source) Moiety Item Code (Source) edient Name Item Code (Source) sF049NKY49) (POLYCARBOPHIL - Item Code (Source) Ingredient Name Item Code (Source) (UNII: 2S7830E561) Item Code (Source) (UNII: 2S7830E561) Item Code (Source) 097M6I30) Item Code (Source) Item Code (Source) Item Code (Source)	HUMAN OTC DRUG Item Code (Source) NDC:55289-94 ORAL Moiety Moiety Basis of Stress redient Name Basis of Stress : 8F049NKY49) (POLYCARBOPHIL - CALCIUM POLYCARBOPHIL Ingredient Name CALCIUM POLYCARBOPHIL (UNII: 2S 7830E561)	HUMAN OTC DRUG ORAL Item Code (Source) NDC:55289-949(NDC:02 Moiety State redient Name Basis of Strength CALCIUM POLYCARBOPHIL - SF049NKY49) (POLYCARBOPHIL - CALCIUM POLYCARBOPHIL Ingredient Name Str Ingredient Name Str (UNII: 257830E561) 5 /3WO) Str (UNII: 257830E561) 5 (JUNII: 257830E561) 5 (JUN

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

SI	LICON DIOXIDE (
SODIUM LAURYL SULFATE (UNII: 368GB5141J)							
STEARIC ACID (UNII: 4ELV7Z65AP)							
Pı	roduct Chara	cteristics					
Co			Score		2 pieces		
Sł	nape	OVAL Size		19mm			
Fla	avor			ode	CPC;339		
Сс	ontains						
-							
Pa	ackaging						
#	ltem Code	Package Descri	ption ^N	Marketing Start Date	Marketing End Date		
1	NDC:55289- 949-30	30 in 1 BOTTLE, PLASTIC; Type 0 Combination Product	: Not a 03	8/27/2021			
Marketing Information							
Μ		Application Number of	r Monograph 🛛 🛚	Marketing Start	Marketing End		
M	Marketing Category	Citation		Date	Date		
		Citation		Date /01/2010	Date		

Labeler - PD-Rx Pharmaceuticals, Inc. (156893695)

Registrant - PD-Rx Pharmaceuticals, Inc. (156893695)

Establishment						
Name	Address	ID/FEI	Business Operations			
PD-Rx Pharmaceuticals, Inc.		156893695	repack(55289-949)			

Revised: 10/2023

PD-Rx Pharmaceuticals, Inc.