BASIC CARE LOPERAMIDE HYDROCHLORIDE- loperamide hcl tablet, film coated

Amazon.com Services LLC

Amazon Loperamide Hydrochloride Tablets Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- · diarrhea lasts for more than 2 days

• you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Anti-Diarrheal

Compare to Imodium® A-D active ingredient

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the symptoms of diarrhea

See New Warnings

actual size

24 CAPLETS*

*CAPSULE-SHAPED TABLETS



22462 BN C OPEN OTHER END

Drug Facts

Act ive ing redient Purpos e (in each caplet)

Loperami de HC12 mg......Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

All ergy alert: Donot use if you have ever had a rashor other allergic reaction to lopera mide HC I Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Aska doctor before use if you have

■ fever ■ mucus in the stool ■a history of liver dis ease

■ a history of abnormal heart rhythm

Aska doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product fredness, drows iress or dizzines smay occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse

■ diarrhea lasts for more than 2 days

you get abdominal swelling or buiging.
These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Pois on Control Center right away (1-800-222-1222)

Directions

■ drink plenty of clear fluids to help preventdehydration caused by diarrhea

■ find right obse on chart. If possible, use weight to dose, otherwise, use age.

adults and children 12 years and o ver	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 ye ars (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 y ears (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 y ears (34 to 47 lbs)	ask adoctor
children under 2 years (up to 33 lbs)	do not use

Other information

"This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A.D. DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

CONVENIENT RECLOSING TAB



BASIC CARE LOPERAMIDE HYDROCHLORIDE

loperamide hcl tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)		
CARNAUBA WAX (UNII: R12CBM0EIZ)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		

Product Characteristics			
Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging		

	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:72288-224- 62	24 in 1 CARTON	02/16/2021	
l	1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	02/16/2021	

Labeler - Amazon.com Services LLC (128990418)

Revised: 2/2021 Amazon.com Services LLC