

## **CHERRY SORE THROAT LOZENGES- benzocaine lozenge**

**Bestco Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Benzocaine 15 mg**

**Menthol 3.6 mg**

### **Active ingredients (in each lozenge)**

Benzocaine 15 mg

Menthol 3.6 mg

### **Purposes**

Oral anesthetic

Oral anesthetic

**Uses** temporarily relieves these symptoms:

occasional minor irritation, pain, sore mouth and sore throat

### **Warnings**

#### **Methemoglobinemia warning:**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

### **Do not use**

- for teething
- in children under 2 years of age

### **Stop use and ask doctor if**

- sore throat is severe or irritation, pain, or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact Poison Control Center right away.

### **Directions**

adults and children 5 years and older - allow 1 lozenge to dissolve slowly in mouth. May be repeated every 2 hours as necessary or as directed by a doctor

children under 5 years - do not use

### **Other information**

- store at room temperature
- protect from moisture

**Inactive ingredients:** Corn starch, FD&C red no. 40, glucose syrup, glycerin, medium chain triglycerides, natural and artificial flavors, propylene glycol, soybean oil, sucrose and water.

**Questions?** Call **1-800-245-2898**, Monday - Friday, 9AM to 4PM EST



# best

cherry 18 lozenges

health sore throat relief 3.6 mg Menthol • 15 mg Benzocaine • Oral Anesthetic

Compare to active ingredient in Cepacol®\*

# best

health sore throat relief

3.6 mg Menthol • 15 mg Benzocaine • Oral Anesthetic

fast, soothing, temporary relief of minor sore throat pain and irritation



cherry 18 lozenges

PN-60001717



Gluten-Free  
Distributed by:  
BestCo LLC,  
Mooresville, NC 28115  
©2021

\*This product is not manufactured or distributed by Reckitt Benckiser owner of the registered trademark Cepacol®. WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BLISTER IS OPEN OR DAMAGED.

**Drug Facts (continued)**  
**Inactive ingredients** Corn starch, corn syrup, FD&C Red #40, glycerin, medium chain triglycerides, artificial flavor, propylene glycol, soybean oil, sucrose and water.  
**Questions?** Call 1-800-245-2898, Monday through Friday, 9AM - 4PM EST

**Drug Facts**  
**Active ingredients (in each lozenge)**  
Benzocaine 15 mg.....Oral anesthetic  
Menthol 3.6 mg.....Oral anesthetic  
**Uses** temporarily relieves these symptoms:  
■ occasional minor irritation, pain, sore mouth and sore throat  
**Warnings**  
**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.  
**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.  
**Do not use**  
■ for teething ■ in children under 2 years of age  
**Stop use and ask doctor if**  
■ sore throat is severe or irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days  
**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center right away.  
**Directions**  
adults and children  
5 years and older  
allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor  
children under 5 years  
do not use  
**Other information** ■ store at room temperature ■ protect from moisture



# CHERRY SORE THROAT LOZENGES

benzocaine lozenge

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52642-036
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg
<b>MENTHOL</b> (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	3.6 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0KO0R)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	17mm
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	B
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:52642-036-18	18 in 1 BLISTER PACK; Type 0: Not a Combination Product	08/15/2018	



### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	08/15/2018	

**Labeler** - Bestco Inc. (002149136)

**Registrant** - Bestco Inc. (002149136)

### Establishment

Name	Address	ID/FEI	Business Operations
Bestco Inc.		002149136	manufacture(52642-036)