

DIPHENHYDRAMINE HYDROCHLORIDE- diphenhydramine hydrochloride injection
HF Acquisition Co. LLC, DBA Health First

DIPHENHYDRAMINE HYDROCHLORIDE- diphenhydramine hydrochloride injection

Description

Diphenhydramine Hydrochloride Injection is a sterile, nonpyrogenic solution for intravenous or deep intramuscular use as an antihistaminic agent. Each mL contains diphenhydramine hydrochloride 50 mg and benzethonium chloride 100 mcg in Water for Injection. pH 4.0-6.5; sodium hydroxide and/or hydrochloric acid added, if needed, for pH adjustment.

The chemical name of diphenhydramine hydrochloride is 2-(Diphenylmethoxy)-N,N-dimethylethylamine hydrochloride. The structural formula is as follows:



C₁₇H₂₁NO • HCl MW 291.82

Diphenhydramine hydrochloride occurs as a white crystalline powder and is freely soluble in water and alcohol.

Clinical Pharmacology

Diphenhydramine hydrochloride is an antihistamine with anticholinergic (drying) and sedative side effects. Antihistamines appear to compete with histamine for cell receptor sites on effector cells.

Diphenhydramine hydrochloride in the injectable form has a rapid onset of action. Diphenhydramine is widely distributed throughout the body, including the CNS. A portion of the drug is excreted unchanged in the urine, while the rest is metabolized via the liver. Detailed information on the pharmacokinetics of Diphenhydramine Hydrochloride Injection is not available.

Indications & Usage

Diphenhydramine Hydrochloride Injection is effective in adults and pediatric patients, other than premature infants and neonates, for the following conditions when the oral form is impractical:

Antihistaminic

For amelioration of allergic reactions to blood or plasma, in anaphylaxis as an adjunct to epinephrine and other standard measures after the acute symptoms have been controlled and for other uncomplicated allergic conditions of the immediate type when oral therapy is impossible or contraindicated.

Motion Sickness

For active treatment of motion sickness.

Antiparkinsonism

For use in parkinsonism, when oral therapy is impossible or contraindicated, as follows: parkinsonism in the elderly who are unable to tolerate more potent agents, mild cases of parkinsonism in other age groups and in other cases of parkinsonism in combination with centrally acting anticholinergic agents.

Contraindications

Use in Neonates or Premature Infants

This drug should not be used in neonates or premature infants.

Use in Nursing Mothers

Because of the higher risk of antihistamines for infants generally, and for neonates and prematures in particular, antihistamine therapy is contraindicated in nursing mothers.

Use as a Local Anesthetic

Because of the risk of local necrosis, this drug should not be used as a local anesthetic.

Antihistamines are also Contraindicated in the Following Conditions

Hypersensitivity to diphenhydramine hydrochloride and other antihistamines of similar chemical structure.

Warnings

Antihistamines should be used with considerable caution in patients with narrow-angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy or bladder-neck obstruction.

Local necrosis has been associated with the use of subcutaneous or intradermal use of intravenous diphenhydramine.

Use in Pediatric Patients

In pediatric patients, especially, antihistamines in overdosage may cause hallucinations, convulsions or death.

As in adults, antihistamines may diminish mental alertness in pediatric patients. In the young pediatric patient, particularly, they may produce excitation.

Use in the Elderly (approximately 60 years or older)

Antihistamines are more likely to cause dizziness, sedation and hypotension in elderly patients.

Precautions

General

Diphenhydramine hydrochloride has an atropine-like action and, therefore, should be used with caution in patients with a history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular disease or hypertension. Use with caution in patients with lower respiratory disease, including asthma.

Information for Patients

Patients taking diphenhydramine hydrochloride should be advised that this drug may cause drowsiness and has an additive effect with alcohol.

Patients should be warned about engaging in activities requiring mental alertness, such as driving a car or operating appliances, machinery, etc.

Drug Interactions

Diphenhydramine hydrochloride has additive effects with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers, etc.)

MAO inhibitors prolong and intensify the anticholinergic (drying) effects of antihistamines.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals to determine mutagenic and carcinogenic potential have not been performed.

Pregnancy

Teratogenic Effects—Pregnancy Category B

Reproduction studies have been performed in rats and rabbits at doses up to 5 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to diphenhydramine hydrochloride. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Pediatric Use

Diphenhydramine should not be used in neonates and premature infants (see Contraindications).

Diphenhydramine may diminish mental alertness, or in the young pediatric patient, cause excitation. Overdosage may cause hallucinations, convulsions or death (see Warnings and Overdosage sections).

See also Dosage and Administration section.

Adverse Reactions

The most frequent adverse reactions are italicized.

General

Urticaria; drug rash; anaphylactic shock; photosensitivity; excessive perspiration; chills; dryness of mouth,

nose and throat.

Cardiovascular System

Hypotension, headache, palpitations, tachycardia, extrasystoles.

Hematologic System

Hemolytic anemia, thrombocytopenia, agranulocytosis.

Nervous System

Sedation, sleepiness, dizziness, disturbed coordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paresthesia, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, neuritis, convulsions.

Gastrointestinal System

Epigastric distress, anorexia, nausea, vomiting, diarrhea, constipation.

Genitourinary System

Urinary frequency, difficult urination, urinary retention, early menses.

Respiratory System

Thickening of bronchial secretions, tightness of chest and wheezing, nasal stuffiness.

Overdosage

Antihistamine overdosage reactions may vary from central nervous system depression to stimulation. Stimulation is particularly likely in pediatric patients. Atropine-like signs and symptoms, dry mouth; fixed, dilated pupils; flushing, and gastrointestinal symptoms may also occur.

Stimulants should not be used.

Vasopressors may be used to treat hypotension.

Dosage and Administration

THIS PRODUCT IS FOR INTRAVENOUS OR INTRAMUSCULAR ADMINISTRATION ONLY.

Diphenhydramine Hydrochloride Injection is indicated when the oral form is impractical.

DOSAGE SHOULD BE INDIVIDUALIZED ACCORDING TO THE NEEDS AND THE RESPONSE OF THE PATIENT.

Pediatric Patients, Other Than Premature Infants and Neonates

5 mg/kg/24 hours or 150 mg/m²/24 hours. Maximum daily dosage is 300 mg. Divide into four doses, administered intravenously at a rate generally not exceeding 25 mg/min, or deep intramuscularly.

Adults

10 to 50 mg intravenously at a rate generally not exceeding 25 mg/min, or deep intramuscularly; 100 mg if required; maximum daily dosage is 400 mg.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

How Supplied

DIPHENHYDRAMINE HCl INJECTION, USP is supplied in the following dosage forms.

NDC 51662-1219-1. DIPHENHYDRAMINE HCl INJECTION, USP 50mg/mL 1mL VIAL

NDC 51662-1219-2 Pouch containing a single DIPHENHYDRAMINE HCl INJECTION, USP 50mg/mL 1mL VIAL

NDC 51662-1219-3 Case of 25 pouches of DIPHENHYDRAMINE HCl INJECTION, USP 50mg/mL 1mL VIAL

HF Acquisition Co LLC, DBA HealthFirst
Mukilteo, WA 98275

Also supplied in the following manufactures dosage forms

Diphenhydramine Hydrochloride Injection, USP 50 mg/mL

1 mL vials packaged in 25s (NDC 0641-0376-25)

Storage

Protect from light. Keep covered in carton until time of use. Store at 20°-25°C (68°-77°F), excursions permitted to 15°-30°C (59°-86°F) [See USP Controlled Room Temperature].

To report SUSPECTED ADVERSE REACTIONS, contact West-Ward Pharmaceutical Corp. at 1-877-845-0689, or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. For Product Inquiry call 1-877-845-0689.

Manufactured by:

[West-Ward logo]

WEST-WARD
PHARMACEUTICALS
Eatontown, NJ 07724 USA

Revised May 2011
462-220-01

Serialized Label

Vial Label

PRINCIPAL DISPLAY LABEL - NDC 51662-1219-2 SERIALIZED POUCH LABEL

 Pouch label

PRINCIPAL DISPLAY LABEL - NDC 51662-1219-3 SERIALIZED CASE LABEL

Case Label

PRINCIPAL DISPLAY LABEL - NDC 51662-1219-3 SERIALIZED RFID LABEL

DIPHENHYDRAMINE HYDROCHLORIDE

diphenhydramine hydrochloride injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:51662-1219(NDC:0641-0376)
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51662-1219-1	1 mL in 1 VIAL; Type 0: Not a Combination Product	08/16/2018	
2	NDC:51662-1219-3	25 in 1 CASE	06/15/2020	
2	NDC:51662-1219-2	1 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080817	08/16/2018	

Labeler - HF Acquisition Co. LLC, DBA Health First (045657305)**Registrant** - HF Acquisition Co. LLC, DBA Health First (045657305)**Establishment**

Name	Address	ID/FEI	Business Operations
HF Acquisition Co. LLC, DBA Health First		045657305	relabel(51662-1219)

Revised: 2/2024

HF Acquisition Co. LLC, DBA Health First