

STOMACH RELIEF- bismuth subsalicylate capsule, liquid filled
Spirit Pharmaceuticals LLC

Stomach Relief

Drug Facts

Active Ingredient (in each tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and anti-diarrheal

Uses

relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate.

Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if you have

• an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have

• fever • mucus in the stool

Ask a doctor or pharmacist before use if you are if you are

you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product

• symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

• swallow with water, do not chew • adults and children 12 years and over: • 2 softgels (1 dose) every 1/2 hour or 4 softgels (2 doses) every hour as needed for diarrhea • 2 soft gels (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) • do not exceed 8 doses (16 softgels) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

• **each softgel contains:** Salicylate 99 mg • very low sodium • store between 15-30°C (59-86°F)

Inactive ingredients

Aerosil, beeswax, FD&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide

Questions or comments?

1-888-333-9792

Principal Display Panel



COMPOSTABLE
REFILL POUCH



CABINET:

Stomach Relief Refill

COMPARE TO THE ACTIVE INGREDIENT IN:


Pepto-Bismol®*

Bismuth Subsalicylate 262 mg

RELIEVES:

NAUSEA · HEARTBURN · INDIGESTION · UPSET STOMACH
DIARRHEA

50 SOFTGELS



TO OPEN:

1. POSITION THUMBS INSIDE THE SHORT FLANGE OVER THE LOCKS.
2. GRIP WHILE PIVOTING THUMBS OUTWARD TO OPEN THE LOCKS.

Tamper evident: do not use if pouch is open

Drug Facts

Active Ingredient (in each tablet)	Purpose
Bismuth subsalicylate 262 mg.....	Upset stomach reliever and anti-diarrheal

Uses relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate.

Do not take if you are

- allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have

- fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions • swallow with water, do not chew • adults and children 12 years and over: • 2 softgels (1 dose) every 1/2 hour or 4 softgels (2 doses) every hour as needed for diarrhea • 2 softgels (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) • do not exceed 8 doses (16 softgels) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea ▶

Drug Facts (continued)

Other Information • each softgel contains: Salicylate 99 mg • very low sodium • store between 15–30°C (59–86°F)

Inactive ingredients Aerosil, beeswax, FD&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide


Questions or comments?

1-888-333-9792

It takes guts to eliminate plastic.

* This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Pepto-Bismol®

† Less than the limit of detection and consistent with gluten-free diet labeling per FDA



NDC: 68210-4170-1

ACC-1-10020
Distributed by: Cabinet Health P.B.C.
Question/Comments? 1-888-333-9792
www.cabinethealth.com

Lot / Expiration ▼



Compostable & Plastic Free



Independently Quality Tested



Gluten Free Verified

STOMACH RELIEF

bismuth subsalicylate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68210-4238
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
SYNTHETIC BEESWAX (UNII: 08MNR5YE2R)	
SILICON (UNII: Z4152N8IU)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
PARAFFIN (UNII: I9O0E3H2ZE)	
WATER (UNII: 059QF0KO0R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
HYDROXYMETHYL CELLULOSE (UNII: 273FM27VK1)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	262
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-4238-5	60 in 1 POUCH; Type 0: Not a Combination Product	11/17/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M008

11/17/2022

Labeler - Spirit Pharmaceuticals LLC (179621011)

Revised: 12/2023

Spirit Pharmaceuticals LLC