

**HARRIS TEETER NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled**  
**Harris Teeter, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Harris Teeter, LLC Night Time Cold & Flu Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed

by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you

do not notice any signs or symptoms.

### **Directions**

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- store at 20-25°C (68-77°F)

### **Inactive ingredients**

D&C yellow no. 10, edible ink\*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution \*may contain this ingredient

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

NightTime

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®

Pain Reliever/Fever Reducer • Acetaminophen

Cough Suppressant • Dextromethorphan HBr

Antihistamine • Doxylamine Succinate

COLD & FLU

Multi-Symptom Relief

Aches, Fever & Sore Throat

Sneezing, Runny Nose

Cough

24 SOFTGELS

Actual Size

**PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org



# NightTime

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®\*\*

**Pain Reliever/Fever Reducer • Acetaminophen  
Cough Suppressant • Dextromethorphan HBr  
Antihistamine • Doxylamine Succinate**

## COLD & FLU

### Multi-Symptom Relief

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose
- Cough

**24** SOFTGELS



Actual Size

**CONVENIENT RECLOSEING TAB**

\*\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu LiquiCaps®.

### Drug Facts

**DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN**

#### Active ingredients (in each softgel)

#### Purpose

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

#### Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- minor aches and pains
- runny nose and sneezing
- sore throat
- headache
- fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

- Ask a doctor before use if you have
- cough that occurs with too much phlegm (mucus)
  - a breathing problem such as emphysema or chronic bronchitis
  - liver disease
  - glaucoma
  - persistent or chronic cough as occurs with smoking, asthma, or emphysema
  - trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

© 2008  
 All rights reserved.  
 "made in Canada"  
 HARRIS TEETER, LLC, MATTHEWS, NC 28055

When using this product ■ excitability may occur, especially in children  
 ■ be careful when driving a motor vehicle or operating machinery ■ avoid alcoholic drinks  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days  
 ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed – see overdose warning ■ do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Drug Facts (continued)  
 Other information ■ store at 20-25°C  
 Inactive ingredients D&C yellow no. 10 glycol, purified water, sorbitol sodium solution  
 Questions or comments? 1-800-719-4

**OPEN OTHER END**

05662 HT C2

## HARRIS TEETER NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69256-056
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

### Product Characteristics

<b>Color</b>	GREEN (clear)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	056
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-056-62	12 in 1 CARTON	11/21/2015	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/21/2015	

**Labeler** - Harris Teeter, LLC (048463103)

Revised: 12/2019

Harris Teeter, LLC