

VET ONE- silver nitrate applicator stick
MWI Veterinary Supply

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

For cauterization of skin or mucous membrane and for removing warts and granulated tissue.

DOSAGE AND ADMINISTRATION:

Moisten the applicator tip with distilled, deionized or purified water and apply the silver nitrate to the affected area by holding, rubbing, or rotating the tip along the affected tissue. The strength of the action is controlled by the dilution with distilled, deionized or purified water. One silver nitrate applicator is generally sufficient for each application. The action of the silver nitrate can be stopped by washing the area with saline solution (0.9% sodium chloride). *Using saline solution to wet the applicator tips, or residual saline from wound flushing/washing will interfere with the action of silver nitrate resulting in cauterization failure.* Only use distilled, deionized or purified water to wet applicator tips. Blot dry wounds that have been flushed/washed with saline prior to applying silver nitrate.

DOSAGE FORMS AND STRENGTHS:

Stick: 50 mg

CONTRAINDICATIONS:

Silver salts stain tissue black due to deposition of reduced silver. The stain gradually disappears within a period of two weeks. Prolonged ingestion or absorption of silver compounds leads to deposition of silver in connective tissues, producing a slate-blue discoloration of the skin known as argyria. This discoloration may also appear on mucous membranes such as the margins of gums. The sclera of the eye is also stained.

WARNINGS AND PRECAUTIONS:

WARNING: KEEP OUT OF REACH OF CHILDREN. The active ingredients are poisonous and may be fatal when ingested in sufficient doses. The symptoms include toxic gastroenteritis, which may lead to coma, convulsion, paralysis and profound alteration of respiration. If poisoning occurs, immediately consult a physician.
WARNING: DO NOT USE ON THE EYES. In case of eye contact, hold eyes open and immediately flush thoroughly with water for at least 15 minutes and consult a physician.
CAUTION: SILVER NITRATE IS A CAUSTIC SUBSTANCE. Chemical burns may result from inappropriate use of product.

- Wear chemical resistant gloves while using this product. Wear other appropriate personal protective equipment as needed.
 - Take care to confine the silver nitrate to the area being treated by using an appropriate physical or chemical barrier to prevent staining or burning of untreated tissue.
 - Skin contact time with applicators should be minimal when used on thin delicate skin or neonates.
 - Avoid prolonged contact with skin or other surfaces since staining may occur.
- CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.**

STORAGE:


Store in the closed package at room temperature in a dry place protected from light. Silver nitrate will oxidize and turn dark brown upon exposure to light, however this does not affect the product’s potency or utility. Exposure to moisture can cause the tip to break or loosen from the applicator. Store away from vaporous chemicals.

DISPOSAL:

Expired or unused applicators may be disposed of according to applicable federal, state and local regulations.

PACKAGE LABEL PRINCIPAL DISPLAY PANEL

<p>NDC 13985-706-10</p>  <p>Silver Nitrate Applicator – 15 cm (6 inches)</p> <p>Active Ingredient: Each Applicator Contains Silver Nitrate..... 75% Petroleum Jelly..... 25%</p> <p>INDICATIONS AND USAGE: For cauterization of skin or mucous membrane and for removing warts and granulated tissue.</p> <p>DOSSAGE AND ADMINISTRATION: Maintain the applicator tip with distilled, deionized or purified water and apply the silver nitrate to the affected area by holding, rubbing, or rotating the tip along the affected tissue. The strength of the action is controlled by the dilution with distilled, deionized or purified water. One silver nitrate applicator is generally sufficient for each application. The action of the silver nitrate can be stopped by washing the area with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator tip, or residual silver from wound re-epithelializing will interfere with the action of silver nitrate resulting in cauterization failure. Only use distilled, deionized or purified water to wet applicator tips. Dried or wounds that have been bathed/washed with saline prior to applying silver nitrate.</p> <p>CONTRAINDICATIONS: Silver salts stain tissue black due to deposition of reduced silver. The stain gradually disappears within a period of two weeks. Prolonged exposure or absorption of silver compounds leads to deposition of silver in connective tissues, producing a slate-blue discoloration of the skin known as argyria. This discoloration may also appear on mucous membranes such as the margins of gums. The sclera of the eye is also stained.</p> <p>POSAGE FORMS AND STRENGTHS: Stick, 50 mg</p> <p>CFI 1000045FW-019</p> <p>W1 510011</p>	<p>NDC 13985-706-10</p> <p>INDICATIONS AND USAGE: For cauterization of skin or mucous membrane and for removing warts and granulated tissue.</p> <p>DOSSAGE AND ADMINISTRATION: Maintain the applicator tip with distilled, deionized or purified water and apply the silver nitrate to the affected area by holding, rubbing, or rotating the tip along the affected tissue. The strength of the action is controlled by the dilution with distilled, deionized or purified water. One silver nitrate applicator is generally sufficient for each application. The action of the silver nitrate can be stopped by washing the area with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator tip, or residual silver from wound re-epithelializing will interfere with the action of silver nitrate resulting in cauterization failure. Only use distilled, deionized or purified water to wet applicator tips. Dried or wounds that have been bathed/washed with saline prior to applying silver nitrate.</p> <p>CONTRAINDICATIONS: Silver salts stain tissue black due to deposition of reduced silver. The stain gradually disappears within a period of two weeks. Prolonged exposure or absorption of silver compounds leads to deposition of silver in connective tissues, producing a slate-blue discoloration of the skin known as argyria. This discoloration may also appear on mucous membranes such as the margins of gums. The sclera of the eye is also stained.</p> <p>POSAGE FORMS AND STRENGTHS: Stick, 50 mg</p> <p>CFI 1000045FW-019</p>
<p>NDC 13985-706-10</p>  <p>Silver Nitrate Applicator – 15 cm (6 inches)</p> <p>Active Ingredient: Each Applicator Contains Silver Nitrate..... 75% Petroleum Jelly..... 25%</p> <p>INDICATIONS AND USAGE: For cauterization of skin or mucous membrane and for removing warts and granulated tissue.</p> <p>DOSSAGE AND ADMINISTRATION: Maintain the applicator tip with distilled, deionized or purified water and apply the silver nitrate to the affected area by holding, rubbing, or rotating the tip along the affected tissue. The strength of the action is controlled by the dilution with distilled, deionized or purified water. One silver nitrate applicator is generally sufficient for each application. The action of the silver nitrate can be stopped by washing the area with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator tip, or residual silver from wound re-epithelializing will interfere with the action of silver nitrate resulting in cauterization failure. Only use distilled, deionized or purified water to wet applicator tips. Dried or wounds that have been bathed/washed with saline prior to applying silver nitrate.</p> <p>CONTRAINDICATIONS: Silver salts stain tissue black due to deposition of reduced silver. The stain gradually disappears within a period of two weeks. Prolonged exposure or absorption of silver compounds leads to deposition of silver in connective tissues, producing a slate-blue discoloration of the skin known as argyria. This discoloration may also appear on mucous membranes such as the margins of gums. The sclera of the eye is also stained.</p> <p>POSAGE FORMS AND STRENGTHS: Stick, 50 mg</p> <p>CFI 1000045FW-019</p>	<p>NDC 13985-706-10</p> <p>INDICATIONS AND USAGE: For cauterization of skin or mucous membrane and for removing warts and granulated tissue.</p> <p>DOSSAGE AND ADMINISTRATION: Maintain the applicator tip with distilled, deionized or purified water and apply the silver nitrate to the affected area by holding, rubbing, or rotating the tip along the affected tissue. The strength of the action is controlled by the dilution with distilled, deionized or purified water. One silver nitrate applicator is generally sufficient for each application. The action of the silver nitrate can be stopped by washing the area with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator tip, or residual silver from wound re-epithelializing will interfere with the action of silver nitrate resulting in cauterization failure. Only use distilled, deionized or purified water to wet applicator tips. Dried or wounds that have been bathed/washed with saline prior to applying silver nitrate.</p> <p>CONTRAINDICATIONS: Silver salts stain tissue black due to deposition of reduced silver. The stain gradually disappears within a period of two weeks. Prolonged exposure or absorption of silver compounds leads to deposition of silver in connective tissues, producing a slate-blue discoloration of the skin known as argyria. This discoloration may also appear on mucous membranes such as the margins of gums. The sclera of the eye is also stained.</p> <p>POSAGE FORMS AND STRENGTHS: Stick, 50 mg</p> <p>CFI 1000045FW-019</p>



VI 510011

Silver Nitrate Applicator – 15 cm (6 inches)

Active Ingredients: Each Applicator Contains
Silver Nitrate..... 75%
Petroleum Jelly..... 25%

INDICATIONS AND USAGE:

For membrane and for removing warts and granulated tissue.

DOSSAGE AND ADMINISTRATION:

1. **Distilled, dechlorinated or purified water** and apply the silver nitrate to the affected area by tip along the affected tissue. The strength of the action is controlled by the dilution with distilled water. Silver nitrate applicator is generally subjected for each application. The action of the silver nitrate applicator with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator. Only applying banding will react with the action of silver nitrate resulting in cauterization of tissue. Only if water to wet applicator. Do not dry wounds that have been treated with silver nitrate prior

CONTRAINDICATIONS AND STRENGTHS:

CONTRAINDICATIONS:

Use to deposition of indurated silver. The stain gradually disappears within a period of two weeks. Use of silver compounds leads to deposition of silver in connective tissues, producing a stain blue in argenta. This discoloration may also appear on mucous membranes such as the margins of the stained

CF 190000469w-019



VI 510011

Silver Nitrate Applicator – 15 cm (6 inches)

Active Ingredients: Each Applicator Contains
Silver Nitrate..... 75%
Petroleum Jelly..... 25%

INDICATIONS AND USAGE:

For membrane and for removing warts and granulated tissue.

DOSSAGE AND ADMINISTRATION:

1. **Distilled, dechlorinated or purified water** and apply the silver nitrate to the affected area by tip along the affected tissue. The strength of the action is controlled by the dilution with distilled water. Silver nitrate applicator is generally subjected for each application. The action of the silver nitrate applicator with saline solution (0.9% sodium chloride). Using saline solution to wet the applicator. Only applying banding will react with the action of silver nitrate resulting in cauterization of tissue. Only if water to wet applicator. Do not dry wounds that have been treated with silver nitrate prior

CONTRAINDICATIONS AND STRENGTHS:

CONTRAINDICATIONS:

Use to deposition of indurated silver. The stain gradually disappears within a period of two weeks. Use of silver compounds leads to deposition of silver in connective tissues, producing a stain blue in argenta. This discoloration may also appear on mucous membranes such as the margins of the stained

CF 190000469w-019



VI 510011

Silver Nitrate Applicator

WARNINGS AND PRECAUTIONS:

WARNING: KEEP OUT OF REACH OF CHILDREN. The active ingredients are a poisonous and may be fatal when ingested in sufficient doses. The symptoms include toxic gastroenteritis, which may lead to coma, convulsion, paralysis and profound alteration of respiration. If poisoning occurs, immediately consult a physician.
WARNING: DO NOT USE ON THE EYES. In case of eye contact, hold eyes open and immediately flush thoroughly with water for at least 15 minutes and consult a physician.

CAUTION: SILVER NITRATE IS A CAUSTIC SUBSTANCE. Chemical burns may result from inappropriate use of product.

- Wear chemical resistant gloves while using the product. Wear other appropriate personal protective equipment as needed.
- Take care to confine the silver nitrate to the area being treated by using an appropriate physical or chemical barrier to prevent staining or burning of untreated tissue.
- Skin contact with applicators should be minimal when used on thin delicate skin or neonates.
- Avoid prolonged contact with skin or other surfaces since staining may occur.

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.

STORAGE:

Store in the closed package at room temperature in a dry place protected from light. Silver nitrate will oxidize and turn dark brown upon exposure to light. However this does not affect the product's potency or safety. Exposure to moisture can cause the tip to break or loosen from the applicator. Store away from vaporous chemicals.

DISPOSAL:

Expired or unused applicators may be disposed of according to applicable federal, state and local regulations.

Distributed by:
VET ONE (Step 1) Inc.
Item: 01205
NEW WICHAMAL
INC. 0119
VI 510011

CF 190000469w-019



VI 510011

Silver Nitrate Applicator

WARNINGS AND PRECAUTIONS:

WARNING: KEEP OUT OF REACH OF CHILDREN. The active ingredients are a poisonous and may be fatal when ingested in sufficient doses. The symptoms include toxic gastroenteritis, which may lead to coma, convulsion, paralysis and profound alteration of respiration. If poisoning occurs, immediately consult a physician.
WARNING: DO NOT USE ON THE EYES. In case of eye contact, hold eyes open and immediately flush thoroughly with water for at least 15 minutes and consult a physician.

CAUTION: SILVER NITRATE IS A CAUSTIC SUBSTANCE. Chemical burns may result from inappropriate use of product.

- Wear chemical resistant gloves while using the product. Wear other appropriate personal protective equipment as needed.
- Take care to confine the silver nitrate to the area being treated by using an appropriate physical or chemical barrier to prevent staining or burning of untreated tissue.
- Skin contact with applicators should be minimal when used on thin delicate skin or neonates.
- Avoid prolonged contact with skin or other surfaces since staining may occur.

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.

STORAGE:

Store in the closed package at room temperature in a dry place protected from light. Silver nitrate will oxidize and turn dark brown upon exposure to light. However this does not affect the product's potency or safety. Exposure to moisture can cause the tip to break or loosen from the applicator. Store away from vaporous chemicals.

DISPOSAL:

Expired or unused applicators may be disposed of according to applicable federal, state and local regulations.

Distributed by:
VET ONE (Step 1) Inc.
Item: 01205
NEW WICHAMAL
INC. 0119
VI 510011

CF 190000469w-019



VI 510011

Silver Nitrate Applicator

WARNINGS AND PRECAUTIONS:

WARNING: KEEP OUT OF REACH OF CHILDREN. The active ingredients are a poisonous and may be fatal when ingested in sufficient doses. The symptoms include toxic gastroenteritis, which may lead to coma, convulsion, paralysis and profound alteration of respiration. If poisoning occurs, immediately consult a physician.
WARNING: DO NOT USE ON THE EYES. In case of eye contact, hold eyes open and immediately flush thoroughly with water for at least 15 minutes and consult a physician.

CAUTION: SILVER NITRATE IS A CAUSTIC SUBSTANCE. Chemical burns may result from inappropriate use of product.

- Wear chemical resistant gloves while using the product. Wear other appropriate personal protective equipment as needed.
- Take care to confine the silver nitrate to the area being treated by using an appropriate physical or chemical barrier to prevent staining or burning of untreated tissue.
- Skin contact with applicators should be minimal when used on thin delicate skin or neonates.
- Avoid prolonged contact with skin or other surfaces since staining may occur.

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.

STORAGE:

Store in the closed package at room temperature in a dry place protected from light. Silver nitrate will oxidize and turn dark brown upon exposure to light. However this does not affect the product's potency or safety. Exposure to moisture can cause the tip to break or loosen from the applicator. Store away from vaporous chemicals.

DISPOSAL:

Expired or unused applicators may be disposed of according to applicable federal, state and local regulations.

Distributed by:
VET ONE (Step 1) Inc.
Item: 01205
NEW WICHAMAL
INC. 0119
VI 510011

CF 190000469w-019



VI 510011

Silver Nitrate Applicator

WARNINGS AND PRECAUTIONS:

WARNING: KEEP OUT OF REACH OF CHILDREN. The active ingredients are a poisonous and may be fatal when ingested in sufficient doses. The symptoms include toxic gastroenteritis, which may lead to coma, convulsion, paralysis and profound alteration of respiration. If poisoning occurs, immediately consult a physician.
WARNING: DO NOT USE ON THE EYES. In case of eye contact, hold eyes open and immediately flush thoroughly with water for at least 15 minutes and consult a physician.

CAUTION: SILVER NITRATE IS A CAUSTIC SUBSTANCE. Chemical burns may result from inappropriate use of product.

- Wear chemical resistant gloves while using the product. Wear other appropriate personal protective equipment as needed.
- Take care to confine the silver nitrate to the area being treated by using an appropriate physical or chemical barrier to prevent staining or burning of untreated tissue.
- Skin contact with applicators should be minimal when used on thin delicate skin or neonates.
- Avoid prolonged contact with skin or other surfaces since staining may occur.

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION.

STORAGE:

Store in the closed package at room temperature in a dry place protected from light. Silver nitrate will oxidize and turn dark brown upon exposure to light. However this does not affect the product's potency or safety. Exposure to moisture can cause the tip to break or loosen from the applicator. Store away from vaporous chemicals.

DISPOSAL:

Expired or unused applicators may be disposed of according to applicable federal, state and local regulations.

Distributed by:
VET ONE (Step 1) Inc.
Item: 01205
NEW WICHAMAL
INC. 0119
VI 510011

CF 190000469w-019

poisonous and may be fatal when
 on may lead to coma, convulsion,
 ily control a physician,
 in and immediately flush thoroughly
 result from inappropriate use of
 personal protective equipment as
 optical, physical or chemical barrier to
 lab or records.

GR1000049Rev019

poisonous and may be fatal when
 on may lead to coma, convulsion,
 ily control a physician,
 in and immediately flush thoroughly
 result from inappropriate use of
 personal protective equipment as
 optical, physical or chemical barrier to
 lab or records.

GR1000049Rev019

VET one Item 510011
100 Applicators/Tube
Silver Nitrate Applicators 6 inch

Each Applicator Contains:
 A Mixture of 75% Silver Nitrate, 25% Potassium Nitrate



NDC 13985-706-10



(01)00313985706103
 (21)T90025560
 (17)221231
 (10)5645

Distributed By:
MWI Veterinary Supply, Inc.
Boise, ID 83705
www.VetOne.net

Made in the USA

Open This End

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. WEAR CHEMICAL RESISTANT GLOVES WHILE USING THIS PRODUCT. STORE IN THE CLOSED PACKAGE AT ROOM TEMPERATURE IN A DRY PLACE PROTECTED FROM LIGHT. SILVER NITRATE WILL OXIDIZE AND TURN DARK BROWN UPON EXPOSURE TO LIGHT, THIS DOES NOT AFFECT THE PRODUCT'S POTENCY OR UTILITY. SEE INSERT FOR COMPLETE PRODUCT INFORMATION.

1

VETONE-PRO-LAB-RevA19

VET one Item 510011
12 Tubes of 100/Box
Silver Nitrate Applicators 6 inch

Each Applicator Contains:
 A Mixture of 75% Silver Nitrate, 25% Potassium Nitrate



NDC 13985-706-10



(01)20313985706107
 (21)B90002514
 (17)221231
 (10)5645

MWI Veterinary Supply, Inc.
Boise, ID 83705
www.VetOne.net

Made in the USA

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. WEAR CHEMICAL RESISTANT GLOVES WHILE USING THIS PRODUCT. STORE IN THE CLOSED PACKAGE AT ROOM TEMPERATURE IN A DRY PLACE PROTECTED FROM LIGHT. SILVER NITRATE WILL OXIDIZE AND TURN DARK BROWN UPON EXPOSURE TO LIGHT, THIS DOES NOT AFFECT THE PRODUCT'S POTENCY OR UTILITY. SEE INSERT FOR COMPLETE PRODUCT INFORMATION.

A

1

VETONE-BOX-LAB-RevA19

VET ONE

silver nitrate applicator stick

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:13985-706
Route of Administration	Topical		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SILVER NITRATE (UNII: 95IT3W8JZE) (SILVER CATION - UNII:57N7B0K90A)	SILVER NITRATE	38.21 mg
POTASSIUM NITRATE (UNII: RU45X2JN0Z) (NITRATE ION - UNII:T93E9Y2844)	POTASSIUM NITRATE	12.74 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROCHLORIC ACID (UNII: QTT17582CB)	0.036 mg
SODIUM CHLORIDE (UNII: 451W47IQ8X)	0.021 mg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-706-10	100 in 1 TUBE; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug	01/01/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/01/2010	

Labeler - MWI Veterinary Supply (019926120)

Registrant - GF Health Products, Inc. (132016168)

Establishment

Name	Address	ID/FEI	Business Operations
GF Health Products, Inc.		798817599	manufacture(13985-706)

Revised: 12/2019

MWI Veterinary Supply