

**ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone liquid**  
**Retail Business Services, LLC.**

-----  
**Careone Regular Strength Antacid**

**ACTIVE INGREDIENTS (in each 10 mL )**

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg  
Magnesium hydroxide 400 mg  
Simethicone 40 mg

**PURPOSE**

Antacid  
Antacid  
Antigas

**USE(S)**

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

**WARNINGS**

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- kidney disease
- a magnesium-restricted diet

**ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**STOP USE AND ASK DOCTOR**

if symptoms last more than two weeks

## **KEEP OUT OF REACH OF CHILDREN**

### **DIRECTIONS**

- **shake well before use**
- mL = milliliter
- adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor

### **OTHER INFORMATION**

- **each 10 mL contains:** magnesium 170 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

### **INACTIVE INGREDIENTS**

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

### **PRINCIPAL DISPLAY PANEL**

NDC 72476-509-05

### **CAREONE REGULAR STRENGTH**

#### **ANTACID LIQUID**

Aluminum hydroxide 400 mg  
Magnesium hydroxide 400 mg  
Simethicone 40 mg


Original Flavor

Relieves:

Acid Indigestion,

Pressure & Bloating (Gas)

**12 FL OZ (355 mL)**Alcohol content 0.2% v/v



**Drug Facts** **DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

**Active ingredients (in each 10 mL) Purpose**  
 Aluminum hydroxide (equiv. to dried gel, USP) 400 mg...Antacid  
 Magnesium hydroxide 400 mg.....Antacid  
 Simethicone 40 mg.....Antigas

**Uses** relieves: ■ acid indigestion ■ heartburn ■ sour stomach  
 ■ upset stomach and gas associated with these symptoms

**Warnings**  
 Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.  
**Ask a doctor before use if you have** ■ kidney disease  
 ■ a magnesium-restricted diet  
**Ask a doctor or pharmacist before use if you are** ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.  
**Stop use and ask a doctor** if symptoms last more than two weeks **Keep out of reach of children.**

**Directions** ■ shake well before use ■ mL = milliliter  
 ■ adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor ■ children under 12 years: consult a doctor

**Other information** ■ each 10 mL contains: magnesium 170 mg, sodium 5 mg ■ store at controlled room temperature 20°C-25°C (68°F-77°F) ■ do not freeze

**Inactive ingredients** ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol.

DISTRIBUTED BY:  
 ADUSA DISTRIBUTION, LLC  
 SALISBURY, NC 28147  
 1-833-992-3872  
 Quality guaranteed or your money back.

MFR# 53041  
 REV 0421

**ANTACID**

aluminum hydroxide, magnesium hydroxide, simethicone liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72476-509
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 10 mL

**Inactive Ingredients**

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%) (UNII: S38J6RZN16)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

### Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-509-05	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/21/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	10/21/2021	

**Labeler** - Retail Business Services, LLC. (967989935)

### Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(72476-509)

Revised: 12/2024

Retail Business Services, LLC.