

**DESGEN DM TABS- dextromethorphan hydrobromide, guaifenesin,
pseudoephedrine hydrochloride tablet
KRAMER NOVIS**

DESGEN DM Tabs

Drug Facts

Active ingredients (in each tablet)

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Pseudoephedrine HCl, 30 mg

Purpose

Antitussive

Expectorant

Nasal Decongestant

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses • temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold or inhaled irritants • helps loosen phlegm and thin bronchial secretions to make coughs more productive • temporarily relieves nasal congestion

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • difficulty in urination due to enlarged prostate gland • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • cough accompanied by excessive phlegm (mucus)

When using this product • do not exceed recommended dosage

Stop use and ask a doctor if • cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition • symptoms do not improve within 7 days or are accompanied by fever • nervousness, dizziness, or sleeplessness occur

If pregnant or breastfeeding ask a health professional before use.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and over	2 tablets every 4 hours, not to exceed 8 tablets in 24 hours or as directed by a doctor
Children 6 to under 12 years of age	1 tablets every 4 hours, not to exceed 4 tablets in 24 hours or as directed by a doctor
Children 2 under 6 years of age	½ tablets every 4 hours, not to exceed 2 tablets in 24 hours or as directed by a doctor
Children under 2 years of age	Consult physician

Inactive ingredients: Magnesium Stearate, microcrystalline cellulose, sodium starch glycolate

Other information

- Store at room temperature 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing

New Formulation

Contains the same active ingredients as Despec® DM Tablets*

• ANTITUSSIVE • EXPECTORANT • NASAL DECONGESTANT

Manufactured in the USA for Kramer Novis. San Juan, PR 00917. www.kramernovis.com / (787) 767-2072

*** Despec® DM Tablets is a registered trademark of International Ethical Laboratories. This product is not manufactured, distributed or marketed by**

Packaging

FRONT PANEL

NDC 52083-648-10

DESGEN[®] DM Tabs

New Formulation

Contains the same active ingredients as Despec[®] DM Tablets*

**• ANTITUSSIVE • EXPECTORANT
• NASAL DECONGESTANT**

Each tablet contains:

Dextromethorphan HBr	10 mg
Guaifenesin	200 mg
Pseudoephedrine HCl	30 mg

Net Contents: 100 Tablets

Manufactured in the USA for
Kramer Novis. San Juan, PR 00917.
www.kramernovis.com / (787) 767-2072


Pharmaceuticals within reach of patients.

Drug Facts

Active Ingredients (in each tablet)

Dextromethorphan HBr 10 mg Antitussive
Guaifenesin 200 mg Expectorant
Pseudoephedrine HCl, 30 mg Nasal Decongestant

Purpose

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold or inhaled irritants ■ helps loosen phlegm and thin bronchial secretions to make coughs more productive ■ temporarily relieves nasal congestion

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes

Peel Here

N 52083-64810 6



DRUG FACTS LABELING

Drug Facts

Active Ingredients (in each tablet)	Purpose
Dextromethorphan HBr 10 mg	Antitussive
Guaifenesin 200 mg	Expectorant
Pseudoephedrine HCl, 30 mg	Nasal Decongestant

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold or inhaled irritants ■ helps loosen phlegm and thin bronchial secretions to make coughs more productive ■ temporarily relieves nasal congestion

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes

Peel Here

Drug Facts (continued)

■ difficulty in urination due to enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by excessive phlegm (mucus)

When using this product ■ **do not exceed recommended dosage**

Stop use and ask a doctor if ■ cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ■ symptoms do not improve within 7 days or are accompanied by fever ■ nervousness, dizziness, or sleeplessness occur

If pregnant or breastfeeding ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Do not exceed recommended dosage.

Adults and children 12 years of age and over	2 tablets every 4 hours, not to exceed 8 tablets in 24 hours or as directed by a doctor
Children 6 to under 12 years of age	1 tablet every 4 hours, not to exceed 4 tablets in 24 hours or as directed by a doctor
Children 2 under 6 years of age	½ tablets every 4 hours, not to exceed 2 tablets in 24 hours or as directed by a doctor
Children under 2 years of age	Consult physician

Other information

■ Store at room temperature 59°-86°F (15°-30°C) (see USP Controlled Room



DESGEN DM TABS

dextromethorphan hydrobromide, guaifenesin, pseudoephedrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52083-648
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	KN648
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52083-648-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/26/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/26/2015	

Labeler - KRAMER NOVIS (090158395)

Registrant - KRAMER NOVIS (090158395)

Revised: 1/2024

KRAMER NOVIS