


ISOSORBIDE DINITRATE- isosorbide dinitrate tablet
NCS HealthCare of KY, LLC dba Vanguard Labs

Isosorbide Dinitrate Tablets

Rx only

DESCRIPTION

Isosorbide dinitrate (ISDN) is 1,4:3,6-dianhydro-D-glucitol 2,5-dinitrate, an organic nitrate whose structural formula is



and whose molecular weight is 236.14. The organic nitrates are vasodilators, active on both arteries and veins.

Isosorbide dinitrate is a white, crystalline powder compound which is stable in air and in solution, has a melting point of 69°C to 72°C and optically active. Isosorbide dinitrate is very soluble in acetone, sparingly soluble in alcohol and very slightly soluble in water.

Each Isosorbide dinitrate tablet contains 5 mg, 10 mg, 20 mg, 30 mg or 40 mg of isosorbide dinitrate. The inactive ingredients in each tablet are microcrystalline cellulose, colloidal silicon dioxide, and magnesium stearate. The 5mg, 10 mg, 20 mg and 30 mg dosage strengths also contain anhydrous lactose and sodium starch glycolate Type A. Additionally 5 mg contains FD & C red 40 Al lake. 10 mg, 20mg, 30 mg and 40 mg contains FD & C Blue 1 Al lake and D & C Yellow 10 Al lake.

FDA approved dissolution test specifications differ from USP.

CLINICAL PHARMACOLOGY

The principal pharmacological action of isosorbide dinitrate is relaxation of vascular smooth muscle and consequent dilatation of peripheral arteries and veins, especially the latter. Dilatation of the veins promotes peripheral pooling of blood and decreases venous return to the heart, thereby reducing left ventricular end-diastolic pressure and pulmonary capillary wedge pressure (preload). Arteriolar relaxation reduces systemic vascular resistance, systolic arterial pressure, and mean arterial pressure (afterload). Dilatation of the coronary arteries also occurs. The relative importance of preload reduction, afterload reduction, and coronary dilatation remains undefined.

Dosing regimens for most chronically used drugs are designed to provide plasma concentrations that are continuously greater than a minimally effective concentration. This strategy is inappropriate for organic nitrates. Several well-controlled clinical trials have used exercise testing to assess the anti-anginal efficacy of continuously-delivered nitrates. In the large majority of these trials, active agents were no more effective than placebo after 24 hours (or less) of continuous therapy. Attempts to overcome nitrate tolerance by dose escalation, even to doses far in excess of those used acutely, have consistently failed. Only after nitrates have been absent from the body for several hours has their anti-anginal efficacy been restored.

Pharmacokinetics

Absorption of isosorbide dinitrate after oral dosing is nearly complete, but bioavailability is highly variable (10% to 90%), with extensive first-pass metabolism in the liver. Serum levels reach their maxima about an hour after ingestion. The average bioavailability of ISDN is about 25%; most studies have observed progressive increases in bioavailability during chronic therapy.

Once absorbed, the volume of distribution of isosorbide dinitrate is 2 to 4 L/kg, and this volume is cleared at the rate of 2 to 4 L/min, so ISDN's half-life in serum is about an hour. Since the clearance exceeds hepatic blood flow, considerable extra hepatic metabolism must also occur. Clearance is affected primarily by denitration to the 2-mononitrate (15 to 25%) and the 5-mononitrate (75 to 85%).

Both metabolites have biological activity, especially the 5-mononitrate. With an overall half-life of about 5 hours, the 5-mononitrate is cleared from the serum by denitration to isosorbide, glucuronidation to the 5-mononitrate glucuronide, and denitration/hydration to sorbitol. The 2-mononitrate has been less well studied, but it appears to participate in the same metabolic pathways, with a half-life of about 2 hours.

The daily dose-free interval sufficient to avoid tolerance to organic nitrates has not been well defined. Studies of nitroglycerin (an organic nitrate with a very short half-life) have shown that daily dose-free intervals of 10 to 12 hours are usually sufficient to minimize tolerance. Daily dose-free intervals that have succeeded in avoiding tolerance during trials of moderate doses (e.g., 30 mg) of immediate-release ISDN have generally been somewhat longer (at least 14 hours), but this is consistent with the longer half-lives of ISDN and its active metabolites.

Few well-controlled clinical trials of organic nitrates have been designed to detect rebound or withdrawal effects. In one such trial, however, subjects receiving nitroglycerin had *less* exercise tolerance at the end of the daily dose-free interval than the parallel group receiving placebo. The incidence, magnitude, and clinical significance of similar phenomena in patients receiving ISDN have not been studied.

Clinical Trials

In clinical trials, immediate-release oral isosorbide dinitrate has been administered in a variety of regimens, with total daily doses ranging from 30 mg to 480 mg. Controlled trials of single oral doses of isosorbide dinitrate have demonstrated effective reductions in exercise-related angina for up to 8 hours. Anti-anginal activity is present about 1 hour after dosing.

Most controlled trials of multiple-dose oral ISDN taken every 12 hours (or more

frequently) for several weeks have shown statistically significant anti-anginal efficacy for only 2 hours after dosing. Once-daily regimens, and regimens with one daily dose-free interval of at least 14 hours (e.g., a regimen providing doses at 0800, 1400, and 1800 hours), have shown efficacy after the first dose of each day that was similar to that shown in the single-dose studies cited above. The effects of the second and later doses have been smaller and shorter-lasting than the effect of the first.

From large, well-controlled studies of other nitrates, it is reasonable to believe that the maximal achievable daily duration of anti-anginal effect from isosorbide dinitrate is about 12 hours. No dosing regimen for isosorbide dinitrate, however, has ever actually been shown to achieve this duration of effect. One study of 8 patients, who were administered a pretitrated dose (average 27.5 mg) of immediate-release ISDN at 0800, 1300, and 1800 hours for 2 weeks, revealed that significant anti-anginal effectiveness was discontinuous and totaled about 6 hours in a 24 hour period.

INDICATIONS AND USAGE

Isosorbide dinitrate tablets are indicated for the prevention of angina pectoris due to coronary artery disease. The onset of action of immediate-release oral isosorbide dinitrate is not sufficiently rapid for this product to be useful in aborting an acute anginal episode.

CONTRAINDICATIONS

Isosorbide dinitrate is contraindicated in patients who are allergic to isosorbide dinitrate or any of its ingredients.

Do not use isosorbide dinitrate in patients who are taking certain drugs for erectile dysfunction (phosphodiesterase inhibitors), such as sildenafil, tadalafil, or vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia.

Do not use isosorbide dinitrate in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension.

WARNINGS

Amplification of the vasodilatory effects of isosorbide dinitrate by sildenafil can result in severe hypotension. The time course and dose dependence of this interaction have not been studied. Appropriate supportive care has not been studied, but it seems reasonable to treat this as a nitrate overdose, with elevation of the extremities and with central volume expansion.

The benefits of immediate-release oral isosorbide dinitrate in patients with acute myocardial infarction or congestive heart failure have not been established. If one elects to use isosorbide dinitrate in these conditions, careful clinical or hemodynamic monitoring must be used to avoid the hazards of hypotension and tachycardia. Because the effects of oral isosorbide dinitrate are so difficult to terminate rapidly, this formulation is not recommended in these settings.

PRECAUTIONS

General

Severe hypotension, particularly with upright posture, may occur with even small doses of isosorbide dinitrate. This drug should therefore be used with caution in patients who may be volume depleted or who, for whatever reason, are already hypotensive.

Hypotension induced by isosorbide dinitrate may be accompanied by paradoxical bradycardia and increased angina pectoris.

Nitrate therapy may aggravate the angina caused by hypertrophic cardiomyopathy.

As tolerance to isosorbide dinitrate develops, the effect of sublingual nitroglycerin on exercise tolerance, although still observable, is somewhat blunted.

Some clinical trials in angina patients have provided nitroglycerin for about 12 continuous hours of every 24-hour day. During the daily dose-free interval in some of these trials, anginal attacks have been more easily provoked than before treatment, and patients have demonstrated hemodynamic rebound and *decreased* exercise tolerance. The importance of these observations to the routine, clinical use of immediate-release oral isosorbide dinitrate is not known.

In industrial workers who have had long-term exposure to unknown (presumably high) doses of organic nitrates, tolerance clearly occurs. Chest pain, acute myocardial infarction, and even sudden death have occurred during temporary withdrawal of nitrates from these workers, demonstrating the existence of true physical dependence.

Information for Patients

Patients should be told that the anti-anginal efficacy of isosorbide dinitrate is strongly related to its dosing regimen, so the prescribed schedule of dosing should be followed carefully. In particular, daily headaches sometimes accompany treatment with isosorbide dinitrate. In patients who get these headaches, the headaches are a marker of the activity of the drug. Patients should resist the temptation to avoid headaches by altering the schedule of their treatment with isosorbide dinitrate, since loss of headache may be associated with simultaneous loss of anti-anginal efficacy. Aspirin and/or acetaminophen, on the other hand, often successfully relieve isosorbide dinitrate-induced headaches with no deleterious effect on isosorbide dinitrate's anti-anginal efficacy.

Treatment with isosorbide dinitrate may be associated with lightheadedness on standing, especially just after rising from a recumbent or seated position. This effect may be more frequent in patients who have also consumed alcohol.

Drug Interactions

The vasodilating effects of isosorbide dinitrate may be additive with those of other vasodilators. Alcohol, in particular, has been found to exhibit additive effects of this variety.

Concomitant use of isosorbide dinitrate with phosphodiesterase inhibitors in any form is contraindicated (see **CONTRAINDICATIONS**).

Concomitant use of isosorbide dinitrate with riociguat, a soluble guanylate cyclase stimulator, is contraindicated (see **CONTRAINDICATIONS**).

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term studies in animals have been performed to evaluate the carcinogenic potential of isosorbide dinitrate. In a modified two-litter reproduction study, there was no remarkable gross pathology and no altered fertility or gestation among rats fed isosorbide dinitrate at 25 or 100 mg/kg/day.

Pregnancy

At oral doses 35 and 150 times the maximum recommended human daily dose, isosorbide dinitrate has been shown to cause a dose-related increase in embryotoxicity (increase in mummified pups) in rabbits. There are no adequate, well-controlled studies in pregnant women. Isosorbide dinitrate should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether isosorbide dinitrate is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when isosorbide dinitrate is administered to a nursing woman.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Clinical studies of isosorbide dinitrate did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

Adverse reactions to isosorbide dinitrate are generally dose-related, and almost all of these reactions are the result of isosorbide dinitrate's activity as a vasodilator. Headache, which may be severe, is the most commonly reported side effect. Headache may be recurrent with each daily dose, especially at higher doses. Transient episodes of lightheadedness, occasionally related to blood pressure changes, may also occur. Hypotension occurs infrequently, but in some patients it may be severe enough to warrant discontinuation of therapy. Syncope, crescendo angina, and rebound hypertension have been reported but are uncommon.

Extremely rarely, ordinary doses of organic nitrates have caused methemoglobinemia in normal-seeming patients. Methemoglobinemia is so infrequent at these doses that further discussion of its diagnosis and treatment is deferred (see **OVERDOSAGE**).

Data are not available to allow estimation of the frequency of adverse reactions during treatment with isosorbide dinitrate tablets.

OVERDOSAGE

Hemodynamic Effects

The ill effects of isosorbide dinitrate overdose are generally the results of isosorbide dinitrate's capacity to induce vasodilatation, venous pooling, reduced cardiac output, and hypotension. These hemodynamic changes may have protean manifestations, including increased intracranial pressure, with any or all of persistent throbbing headache, confusion, and moderate fever; vertigo; palpitations; visual disturbances; nausea and vomiting (possibly with colic and even bloody diarrhea); syncope (especially in the upright posture); air hunger and dyspnea, later followed by reduced ventilatory effort; diaphoresis, with the skin either flushed or cold and clammy; heart block and bradycardia; paralysis; coma; seizures; and death.

Laboratory determinations of serum levels of isosorbide dinitrate and its metabolites are not widely available, and such determinations have, in any event, no established role in the management of isosorbide dinitrate overdose.

There are no data suggesting what dose of isosorbide dinitrate is likely to be life-threatening in humans. In rats, the median acute lethal dose (LD₅₀) was found to be 1100 mg/kg.

No data are available to suggest physiological maneuvers (e.g., maneuvers to change the pH of the urine) that might accelerate elimination of isosorbide dinitrate and its active metabolites. Similarly, it is not known which, if any, of these substances can usefully be removed from the body by hemodialysis.

No specific antagonist to the vasodilator effects of isosorbide dinitrate is known, and no intervention has been subject to controlled studies as a therapy for isosorbide dinitrate overdose. Because the hypotension associated with isosorbide dinitrate overdose is the result of venodilatation and arterial hypovolemia, prudent therapy in this situation should be directed toward increase in central fluid volume. Passive elevation of the patient's legs may be sufficient, but intravenous infusion of normal saline or similar fluid may also be necessary.

The use of epinephrine or other arterial vasoconstrictors in this setting is likely to do more harm than good.

In patients with renal disease or congestive heart failure, therapy resulting in central volume expansion is not without hazard. Treatment of isosorbide dinitrate overdose in these patients may be subtle and difficult, and invasive monitoring may be required.

Methemoglobinemia

Nitrate ions liberated during metabolism of isosorbide dinitrate can oxidize hemoglobin into methemoglobin. Even in patients totally without cytochrome b₅ reductase activity, however, and even assuming that the nitrate moieties of isosorbide dinitrate are quantitatively applied to oxidation of hemoglobin, about 1 mg/kg of isosorbide dinitrate should be required before any of these patients manifests clinically significant ($\geq 10\%$) methemoglobinemia. In patients with normal reductase function, significant production of methemoglobin should require even larger doses of isosorbide dinitrate. In one study in which 36 patients received 2 to 4 weeks of continuous nitroglycerin therapy at 3.1 to 4.4 mg/hr (equivalent, in total administered dose of nitrate ions, to 4.8 to 6.9 mg of bioavailable isosorbide dinitrate per hour), the average methemoglobin level measured

was 0.2%; this was comparable to that observed in parallel patients who received placebo.

Notwithstanding these observations, there are case reports of significant methemoglobinemia in association with moderate overdoses of organic nitrates. None of the affected patients had been thought to be unusually susceptible.

Methemoglobin levels are available from most clinical laboratories. The diagnosis should be suspected in patients who exhibit signs of impaired oxygen delivery despite adequate cardiac output and adequate arterial pO₂. Classically, methemoglobinemic blood is described as chocolate brown, without color change on exposure to air.

When methemoglobinemia is diagnosed, the treatment of choice is methylene blue, 1 to 2 mg/kg intravenously.

DOSAGE AND ADMINISTRATION

As noted under **CLINICAL PHARMACOLOGY**, multiple-dose studies with ISDN and other nitrates have shown that maintenance of continuous 24-hour plasma levels results in refractory tolerance. Every dosing regimen for isosorbide dinitrate tablets must provide a daily dose-free interval to minimize the development of this tolerance. With immediate-release ISDN, it appears that one daily dose-free interval must be at least 14 hours long.

As also noted under **CLINICAL PHARMACOLOGY**, the effects of the second and later doses have been smaller and shorter-lasting than the effects of the first.

Large controlled studies with other nitrates suggest that no dosing regimen with isosorbide dinitrate tablets should be expected to provide more than about 12 hours of continuous anti-anginal efficacy per day.

As with all titratable drugs, it is important to administer the minimum dose which produces the desired clinical effect. The usual starting dose of isosorbide dinitrate is 5 mg to 20 mg, two or three times daily. For maintenance therapy, 10 mg to 40 mg, two or three times daily is recommended. Some patients may require higher doses. A daily dose-free interval of at least 14 hours is advisable to minimize tolerance. The optimal interval will vary with the individual patient, dose and regimen.

HOW SUPPLIED

Isosorbide dinitrate Tablets are available as follows:

5 mg, Light pink to pink round tablets, debossed with, "Λ" on one side and score line on the other side.

21

10 mg, Light green to green mottled round tablets, debossed with, "Λ" on one side and score line on the other side.

22

Blistercards of 30 NDC 0615-8463-39

20 mg, Light green to green mottled round tablets, debossed with, "Λ" on one side and score line on the other side.

23

Blistercards of 30 NDC 0615-8464-39

30 mg, Light green to green mottled round tablets, debossed with, "Λ" on one side and score line on the other side.

24

Blistercards of 30 NDC 0615-8465-39

40 mg, Light green to green mottled round tablets, debossed with, "Λ" on one side and score line on the other side.

45

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Protect from light.

Keep bottles tightly closed.

Dispense in a light-resistant, tight container.

All trademarks are the property of their respective owners.

Manufactured by:

Rubicon Research Private Limited
Ambarnath, Dist: Thane, 421506 India

Distributed by:

Advagen Pharma Ltd
666 Plainsboro Road
Suite 605 Plainsboro, NJ 08536, US

Revision: 06/22

PRINCIPAL DISPLAY PANEL - 10 mg



VLI NDC 0615-8463-39

Isosorbide Dinitrate

798880620139 Tabs USP 10 mg Rx only

LOT 8463-

EXP

8463-AA-39 v00

QTY 30

Mfg By Rubicon for Advagen (NDC 72888-082-01) PKG BY VANGAIRD GLASGOW, KY 42141

(Advagen NDC 72888-082-01)

Isosorbide Dinitrate Tabs USP 10 mg

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1
Start Date	Start Time		

STORE AT 20° - 25°C (68° - 77° F) (SEE USP CONTROLLED ROOM TEMPERATURE)

Dispense in a tight, light-resistant container. Protect from light.

See package insert or label for dosage information FOR INSTITUTIONAL USE ONLY

30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	8463-AA-B-v00 Vangard Labs Glasgow, KY 42141 LOT 8463 - EXP 7288806201
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 10 mg LOT 8463 - EXP 7288806201

24 The overall configuration of this package is a trademark of Omnicare, Inc. 17 9 1

PRINCIPAL DISPLAY PANEL - 20 mg



0615-8464-39

Isosorbide
Dinitrate

Tab USP 20 mg
LOT 8464-
Rx only

EXP

8464-AA-39 v00

QTY
30

Mfg By Rubicon for
Advagen
(NDC 72888-083-01)
PKG BY VANGAIRD
GLASGOW, KY 42141

(Advagen NDC 72888-083-01)
Isosorbide
Dinitrate
Tabs USP 20 mg

		16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
	24	17	9
			1

Omnicare The overall configuration of this package is a trademark of Omnicare, Inc.

Received:

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
	17	9	1
Start Date	Start Time		

STORE AT 20° - 25° C (68° - 77° F)
(SEE USP CONTROLLED ROOM TEMPERATURE)

Dispense in a tight, light-resistant container.
Protect from light.

See package insert or label for dosage information
FOR INSTITUTIONAL USE ONLY

<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>8464-AA-B-v00 Vangard Labs Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>
<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>
<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>
<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>
<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>
<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>	<p>Pkg by Vangard, Glasgow, KY 42141</p> <p>Isosorbide Dinitrate Tab USP 20 mg LOT 8464 - EXP 7288806301</p>

PRINCIPAL DISPLAY PANEL - 30 mg



ULI NDC 0615-8465-39

Isosorbide
Dinitrate

Tab USP 30 mg
LOT 8465-

Rx only

EXP

8465-AA-39 v00

QTY
30

Mfg By Rubicon for
Advagen
(NDC 72888-084-01)
PKG BY VANGAIED
GLASGOW, KY 42141

(Advagen NDC 72888-084-01)

Isosorbide
Dinitrate
Tab USP 30 mg

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
	17	9	1

Start Date _____ Start Time _____

STORE AT 20° - 25°C (68° - 77° F)
(SEE USP CONTROLLED ROOM TEMPERATURE)

Dispense in a tight, light-resistant container.
Protect from light.

See package insert or label for dosage information
FOR INSTITUTIONAL USE ONLY

30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	8465-AA-B-v00 Vangard Labs Glasgow, KY 42141 LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP
Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP 	Pkg by Vangard, Glasgow, KY 42141 Isosorbide Dinitrate Tab USP 30 mg LOT 8465 - EXP



The overall configuration of this package is a trademark of Omnicare, Inc.

ISOSORBIDE DINITRATE

isosorbide dinitrate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8463(NDC:72888-082)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOSORBIDE DINITRATE (UNII: IA7306519N) (ISOSORBIDE DINITRATE - UNII:IA7306519N)	ISOSORBIDE DINITRATE	10 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

Product Characteristics

Color	green (Light green to green)	Score	2 pieces
Shape	ROUND	Size	6mm
Flavor		Imprint Code	UpArrowhead22
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8463-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/02/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215723	07/08/2022	

ISOSORBIDE DINITRATE

isosorbide dinitrate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8464(NDC:72888-083)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOSORBIDE DINITRATE (UNII: IA7306519N) (ISOSORBIDE DINITRATE - UNII:IA7306519N)	ISOSORBIDE DINITRATE	20 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

Product Characteristics

Color	green (Light green to green)	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	UpArrowhead23
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8464-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/15/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215723	07/08/2022	

ISOSORBIDE DINITRATE

isosorbide dinitrate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8465(NDC:72888-084)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOSORBIDE DINITRATE (UNII: IA7306519N) (ISOSORBIDE DINITRATE - UNII:IA7306519N)	ISOSORBIDE DINITRATE	30 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

Product Characteristics

Color	green (Light green to green)	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	UpArrowhead24
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8465-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215723	07/08/2022	

Labeler - NCS HealthCare of KY, LLC dba Vanguard Labs (050052943)

Establishment

Name	Address	ID/FEI	Business Operations
NCS HealthCare of KY, LLC dba Vanguard Labs		050052943	repack(0615-8463, 0615-8464, 0615-8465)

Revised: 5/2023

NCS HealthCare of KY, LLC dba Vanguard Labs