

**CANDESARTAN CILEXETIL AND HYDROCHLOROTHIAZIDE - candesartan cilexetil and hydrochlorothiazide tablet**  
**Macleods Pharmaceuticals Limited**

-----  
**Candesartan Cilexetil and Hydrochlorothiazide Tablets, for oral use**  
**Rx only**

**WARNING: FETAL TOXICITY**


- **When pregnancy is detected, discontinue candesartan cilexetil and hydrochlorothiazide tablets as soon as possible.**
- **Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus. See Warnings: Fetal Toxicity**

**DESCRIPTION**

Candesartan cilexetil and hydrochlorothiazide tablets combines an angiotensin II receptor (type AT<sub>1</sub>) antagonist and a diuretic, hydrochlorothiazide.

Candesartan cilexetil, USP a nonpeptide, is chemically described as (±)-1-Hydroxyethyl 2-ethoxy-1-[p-(*o*-1*H*-tetrazol-5-ylphenyl)benzyl]-7-benzimidazolecarboxylate, cyclohexyl carbonate (ester).


Its empirical formula is C<sub>33</sub>H<sub>34</sub>N<sub>6</sub>O<sub>6</sub>, and its structural formula is



↓ site of ester hydrolysis.

Candesartan cilexetil USP is a white to off-white powder with a molecular weight of 610.67. It is practically insoluble in water and sparingly soluble in methanol. Candesartan cilexetil USP is a racemic mixture containing one chiral center at the cyclohexyloxycarbonyloxy ethyl ester group. Following oral administration, candesartan cilexetil USP undergoes hydrolysis at the ester link to form the active drug, candesartan, which is achiral.

Hydrochlorothiazide USP is 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide. Its empirical formula is  $C_7H_8ClN_3O_4S_2$  and its structural formula is



Hydrochlorothiazide USP is a white, or practically white, crystalline powder with a molecular weight of 297.72, which is slightly soluble in water, but freely soluble in sodium hydroxide solution.

Candesartan cilexetil and hydrochlorothiazide tablets are available for oral administration in three tablet strengths of candesartan cilexetil USP and hydrochlorothiazide USP.

Candesartan cilexetil and hydrochlorothiazide tablets 16 mg/12.5 mg contain 16 mg of candesartan cilexetil USP and 12.5 mg of hydrochlorothiazide USP.

Candesartan cilexetil and hydrochlorothiazide tablets 32 mg/12.5 mg contain 32 mg of candesartan cilexetil USP and 12.5 mg of hydrochlorothiazide USP.

Candesartan cilexetil and hydrochlorothiazide tablets 32 mg/25 mg contain 32 mg of candesartan cilexetil USP and 25 mg of hydrochlorothiazide USP.

The inactive ingredients of the tablets are carboxymethylcellulose calcium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, corn starch, glycerin, and ferric oxide (yellow). Ferric oxide (red) is also added to the 16 mg/12.5 mg and 32 mg/25 mg tablets as colorant.

## CLINICAL PHARMACOLOGY

### Mechanism of Action

Angiotensin II is formed from angiotensin I in a reaction catalyzed by angiotensin-converting enzyme (ACE, kininase II).

Angiotensin II is the principal pressor agent of the renin-angiotensin system, with effects that include vasoconstriction, stimulation of synthesis and release of aldosterone, cardiac stimulation, and renal reabsorption of sodium. Candesartan blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT<sub>1</sub> receptor in many tissues, such as vascular smooth muscle and the adrenal gland. Its action is, therefore, independent of the pathways for angiotensin II synthesis.

There is also an AT<sub>2</sub> receptor found in many tissues, but AT<sub>2</sub> is not known to be associated with cardiovascular homeostasis. Candesartan has much greater affinity (>10,000- fold) for the AT<sub>1</sub> receptor than for the AT<sub>2</sub> receptor.

Blockade of the renin-angiotensin system with ACE inhibitors, which inhibit the biosynthesis of angiotensin II from angiotensin I, is widely used in the treatment of hypertension. ACE inhibitors also inhibit the degradation of bradykinin, a reaction also catalyzed by ACE. Because candesartan does not inhibit ACE (kininase II), it does not affect the response to bradykinin. Whether this difference has clinical relevance is not yet known. Candesartan does not bind to or block other hormone receptors or ion channels known to be important in cardiovascular regulation.

Blockade of the angiotensin II receptor inhibits the negative regulatory feedback of angiotensin II on renin secretion, but the resulting increased plasma renin activity and angiotensin II circulating levels do not overcome the effect of candesartan on blood pressure.

Hydrochlorothiazide is a thiazide diuretic. Thiazides affect the renal tubular mechanisms of electrolyte reabsorption, directly increasing excretion of sodium and chloride in approximately equivalent amounts. Indirectly, the diuretic action of hydrochlorothiazide reduces plasma volume, with consequent increases in plasma renin activity, increases in aldosterone secretion, increases in urinary potassium loss, and decreases in serum potassium. The renin-aldosterone link is mediated by angiotensin II, so co-administration of an angiotensin II receptor antagonist tends to reverse the potassium loss associated with these diuretics.

The mechanism of the antihypertensive effect of thiazides is unknown.

### Pharmacokinetics

#### General

##### *Candesartan Cilexetil*

Candesartan cilexetil is rapidly and completely bioactivated by ester hydrolysis during absorption from the gastrointestinal tract to candesartan, a selective AT<sub>1</sub> subtype

angiotensin II receptor antagonist. Candesartan is mainly excreted unchanged in urine and feces (via bile). It undergoes minor hepatic metabolism by O-deethylation to an inactive metabolite. The elimination half-life of candesartan is approximately 9 hours. After single and repeated administration, the pharmacokinetics of candesartan are linear for oral doses up to 32 mg of candesartan cilexetil. Candesartan and its inactive metabolite do not accumulate in serum upon repeated once-daily dosing. Following administration of candesartan cilexetil, the absolute bioavailability of candesartan was estimated to be 15%. After tablet ingestion, the peak serum concentration ( $C_{\max}$ ) is reached after 3 to 4 hours. Food with a high fat content does not affect the bioavailability of candesartan after candesartan cilexetil administration.

#### *Hydrochlorothiazide*

When plasma levels have been followed for at least 24 hours, the plasma half-life has been observed to vary between 5.6 and 14.8 hours.

### **Metabolism and Excretion**

#### *Candesartan Cilexetil*

Total plasma clearance of candesartan is 0.37 mL/min/kg, with a renal clearance of 0.19 mL/min/kg. When candesartan is administered orally, about 26% of the dose is excreted unchanged in urine. Following an oral dose of  $^{14}\text{C}$ -labeled candesartan cilexetil, approximately 33% of radioactivity is recovered in urine and approximately 67% in feces. Following an intravenous dose of  $^{14}\text{C}$ -labeled candesartan, approximately 59% of radioactivity is recovered in urine and approximately 36% in feces. Biliary excretion contributes to the elimination of candesartan.

#### *Hydrochlorothiazide:*

Hydrochlorothiazide is not metabolized but is eliminated rapidly by the kidney. At least 61% of the oral dose is eliminated unchanged within 24 hours.

### **Distribution**

#### *Candesartan Cilexetil*

The volume of distribution of candesartan is 0.13 L/kg. Candesartan is highly bound to plasma proteins (>99%) and does not penetrate red blood cells. The protein binding is constant at candesartan plasma concentrations well above the range achieved with recommended doses. In rats, it has been demonstrated that candesartan crosses the blood-brain barrier poorly, if at all. It has also been demonstrated in rats that candesartan passes across the placental barrier and is distributed in the fetus.

#### *Hydrochlorothiazide*

Hydrochlorothiazide crosses the placental but not the blood-brain barrier and is excreted in breast milk.

### **Special Populations**

#### *Pediatric*

The pharmacokinetics of candesartan cilexetil have not been investigated in patients <18 years of age.

#### *Geriatric*

The pharmacokinetics of candesartan have been studied in the elderly ( $\geq 65$  years). The plasma concentration of candesartan was higher in the elderly ( $C_{\max}$  was approximately 50% higher, and AUC was approximately 80% higher) compared to younger subjects

administered the same dose. The pharmacokinetics of candesartan were linear in the elderly, and candesartan and its inactive metabolite did not accumulate in the serum of these subjects upon repeated, once-daily administration. No initial dosage adjustment is necessary. (See DOSAGE AND ADMINISTRATION.)

#### *Gender*

There is no difference in the pharmacokinetics of candesartan between male and female subjects.

#### *Renal Insufficiency*

In hypertensive patients with renal insufficiency, serum concentrations of candesartan were elevated. After repeated dosing, the AUC and  $C_{max}$  were approximately doubled in patients with severe renal impairment (creatinine clearance  $<30$  mL/min/1.73m<sup>2</sup>) compared to patients with normal kidney function. The pharmacokinetics of candesartan in hypertensive patients undergoing hemodialysis are similar to those in hypertensive patients with severe renal impairment. Candesartan cannot be removed by hemodialysis. Thiazide diuretics are eliminated by the kidney, with a terminal half-life of 5-15 hours. In a study of patients with impaired renal function (mean creatinine clearance of 19 mL/min), the half-life of hydrochlorothiazide elimination was lengthened to 21 hours. (See DOSAGE AND ADMINISTRATION.)

Safety and effectiveness of candesartan cilexetil and hydrochlorothiazide in patients with severe renal impairment (CrCL  $\leq 30$  mL/min) have not been established. No dose adjustment is required in patients with mild CrCL 60-90 mL/min) or moderate (CrCL 30-60) renal impairment.

#### *Hepatic Insufficiency*

The pharmacokinetics of candesartan were compared in patients with mild (Child-Pugh A) or moderate (Child-Pugh B) hepatic impairment to matched healthy volunteers following a single dose of 16 mg candesartan cilexetil. The AUC for candesartan in patients with mild and moderate hepatic impairment was increased 30% and 145% respectively. The  $C_{max}$  for candesartan was increased 56% and 73% respectively. The pharmacokinetics of candesartan in severe hepatic impairment have not been studied. No dose adjustment is recommended for patients with mild hepatic impairment. In patients with moderate hepatic impairment, candesartan cilexetil and hydrochlorothiazide tablets are not recommended for initiation because the appropriate starting dose, 8 mg, cannot be given. (See DOSAGE AND ADMINISTRATION). Monitor patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma.

## **Pharmacodynamics**

#### *Candesartan Cilexetil*

Candesartan inhibits the pressor effects of angiotensin II infusion in a dose-dependent manner. After 1 week of once-daily dosing with 8 mg of candesartan cilexetil, the pressor effect was inhibited by approximately 90% at peak with approximately 50% inhibition persisting for 24 hours.

Plasma concentrations of angiotensin I and angiotensin II, and plasma renin activity (PRA), increased in a dose-dependent manner after single and repeated administration of candesartan cilexetil to healthy subjects and hypertensive patients. ACE activity was not altered in healthy subjects after repeated candesartan cilexetil administration. The

once-daily administration of up to 16 mg of candesartan cilexetil to healthy subjects did not influence plasma aldosterone concentrations, but a decrease in the plasma concentration of aldosterone was observed when 32 mg of candesartan cilexetil was administered to hypertensive patients. In spite of the effect of candesartan cilexetil on aldosterone secretion, very little effect on serum potassium was observed.

In multiple-dose studies with hypertensive patients, there were no clinically significant changes in metabolic function including serum levels of total cholesterol, triglycerides, glucose, or uric acid. In a 12-week study of 161 patients with non-insulin-dependent (type 2) diabetes mellitus and hypertension, there was no change in the level of HbA<sub>1c</sub>.

#### *Hydrochlorothiazide*

After oral administration of hydrochlorothiazide, diuresis begins within 2 hours, peaks in about 4 hours and lasts about 6 to 12 hours.

### **Clinical Trials**

#### *Candesartan Cilexetil-Hydrochlorothiazide*

Of 12 controlled clinical trials involving 4588 patients, 5 were double-blind, placebo controlled and evaluated the antihypertensive effects of single entities vs the combination. These 5 trials, of 8 to 12 weeks duration, randomized 3037 hypertensive patients. Doses ranged from 2 to 32 mg candesartan cilexetil and from 6.25 to 25 mg hydrochlorothiazide administered once daily in various combinations.

The combination of candesartan cilexetil- hydrochlorothiazide resulted in placebo-adjusted decreases in sitting systolic and diastolic blood pressures of 14-18/8-11 mm Hg at doses of 16 mg/12.5 mg and 32 mg/12.5 mg. The combination of candesartan cilexetil and hydrochlorothiazide 32 mg/25 mg resulted in placebo-adjusted decreases in sitting systolic and diastolic blood pressures of 16-19/9-11 mm Hg. The placebo corrected trough to peak ratio was evaluated in a study of candesartan cilexetil- hydrochlorothiazide 32 mg/12.5 mg and was 88%.

Most of the antihypertensive effect of the combination of candesartan cilexetil and hydrochlorothiazide was seen in 1 to 2 weeks with the full effect observed within 4 weeks. In long-term studies of up to 1 year, the blood pressure lowering effect of the combination was maintained. The antihypertensive effect was similar regardless of age or gender, and overall response to the combination was similar in black and non-black patients. No appreciable changes in heart rate were observed with combination therapy in controlled trials.

### **INDICATIONS & USAGE**

Candesartan cilexetil and hydrochlorothiazide tablets are indicated for the treatment of hypertension, to lower blood pressure. Lowering blood pressure reduces the risk of fatal and non-fatal cardiovascular events, primarily strokes and myocardial infarctions. These benefits have been seen in controlled trials of antihypertensive drugs from a wide variety of pharmacologic classes including the class to which this drug principally belongs. There are no controlled trials demonstrating risk reduction with Candesartan cilexetil and hydrochlorothiazide tablets.

Control of high blood pressure should be part of comprehensive cardiovascular risk management, including, as appropriate, lipid control, diabetes management, antithrombotic therapy, smoking cessation, exercise, and limited sodium intake. Many patients will require more than one drug to achieve blood pressure goals. For specific advice on goals and management, see published guidelines, such as those of the

National High Blood Pressure Education Program's Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC). Numerous antihypertensive drugs, from a variety of pharmacologic classes and with different mechanisms of action, have been shown in randomized controlled trials to reduce cardiovascular morbidity and mortality, and it can be concluded that it is blood pressure reduction, and not some other pharmacologic property of the drugs, that is largely responsible for those benefits. The largest and most consistent cardiovascular outcome benefit has been a reduction in the risk of stroke, but reductions in myocardial infarction and cardiovascular mortality also have been seen regularly. Elevated systolic or diastolic pressure causes increased cardiovascular risk, and the absolute risk increase per mmHg is greater at higher blood pressures, so that even modest reductions of severe hypertension can provide substantial benefit. Relative risk reduction from blood pressure reduction is similar across populations with varying absolute risk, so the absolute benefit is greater in patients who are at higher risk independent of their hypertension (for example, patients with diabetes or hyperlipidemia), and such patients would be expected to benefit from more aggressive treatment to a lower blood pressure goal. Some antihypertensive drugs have smaller blood pressure effects (as monotherapy) in black patients, and many antihypertensive drugs have additional approved indications and effects (e.g., on angina, heart failure, or diabetic kidney disease). These considerations may guide selection of therapy. This fixed dose combination is not indicated for initial therapy (see DOSAGE AND ADMINISTRATION).

## **CONTRAINDICATIONS**

Candesartan cilexetil and hydrochlorothiazide tablets are contraindicated in patients who are hypersensitive to candesartan, to hydrochlorothiazide or to other sulfonamide-derived drugs.

Do not co-administer aliskiren with candesartan cilexetil and hydrochlorothiazide in patients with diabetes (See PRECAUTIONS, Drug Interactions).

Candesartan cilexetil and hydrochlorothiazide is contraindicated in patients with anuria.

## **WARNINGS**

### **Fetal Toxicity**

Use of drugs that act on the renin-angiotensin system during the second and third trimesters of pregnancy reduces fetal renal function and increases fetal and neonatal morbidity and death. Resulting oligohydramnios can be associated with fetal lung hypoplasia and skeletal deformations. Potential neonatal adverse effects include skull hypoplasia, anuria, hypotension, renal failure, and death. When pregnancy is detected, discontinue candesartan cilexetil and hydrochlorothiazide tablets as soon as possible. These adverse outcomes are usually associated with use of these drugs in the second and third trimester of pregnancy. Most epidemiologic studies examining fetal abnormalities after exposure to antihypertensive use in the first trimester have not

distinguished drugs affecting the renin-angiotensin system from other antihypertensive agents. Appropriate management of maternal hypertension during pregnancy is important to optimize outcomes for both mother and fetus.

In the unusual case that there is no appropriate alternative to therapy with drugs affecting the renin-angiotensin system for a particular patient, apprise the mother of the potential risk to the fetus. Perform serial ultrasound examinations to assess the intra-amniotic environment. If oligohydramnios is observed, discontinue candesartan cilexetil and hydrochlorothiazide tablets, unless it is considered lifesaving for the mother. Fetal testing may be appropriate, based on the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury. Closely observe infants with histories of in utero exposure to candesartan cilexetil and hydrochlorothiazide tablets for hypotension, oliguria, and hyperkalemia. (see PRECAUTIONS, Pediatric Use.)

There was no evidence of teratogenicity or other adverse effects on embryo-fetal development when pregnant mice, rats or rabbits were treated orally with candesartan cilexetil alone or in combination with hydrochlorothiazide. For mice, the maximum dose of candesartan cilexetil was 1000 mg/kg/day (about 150 times the maximum recommended daily human dose [MRHD]<sup>1</sup>). For rats, the maximum dose of candesartan cilexetil was 100 mg/kg/day (about 31 times the MRHD<sup>1</sup>). For rabbits, the maximum dose of candesartan cilexetil was 1 mg/kg/day (a maternally toxic dose that is about half the MRHD<sup>1</sup>). In each of these studies, hydrochlorothiazide was tested at the same dose level (10 mg/kg/day, about 4, 8, and 15 times the MRHD<sup>1</sup> in mouse, rats, and rabbit, respectively). There was no evidence of harm to the rat or mouse fetus or embryo in studies in which hydrochlorothiazide was administered alone to the pregnant rat or mouse at doses of up to 1000 and 3000 mg/kg/day, respectively.

Thiazides cross the placental barrier and appear in cord blood. There is a risk of fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions that have occurred in adults.

---

<sup>1</sup> Doses compared on the basis of body surface area. MRHD considered to be 32 mg for candesartan cilexetil and 12.5 mg for hydrochlorothiazide.

## **Hypotension**

Candesartan cilexetil and hydrochlorothiazide can cause symptomatic hypotension. Symptomatic hypotension is most likely to occur in patients who have been volume and/or salt depleted as a result of prolonged diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. Patients with symptomatic hypotension may require temporarily reducing the dose of candesartan cilexetil and hydrochlorothiazide or volume repletion. Volume and/or salt depletion should be corrected before initiating therapy with candesartan cilexetil and hydrochlorothiazide tablets.

In patients with heart failure, candesartan cilexetil and hydrochlorothiazide may cause excessive hypotension, which may lead to oliguria, azotemia, and (rarely) with acute renal failure and death (see WARNINGS, Impaired Renal Function). In such patients, candesartan cilexetil and hydrochlorothiazide therapy should be started under close medical supervision; they should be followed closely for the first 2 weeks of treatment and whenever the dose of candesartan or diuretic is increased.



## **Impaired Renal Function**

Monitor renal function periodically in patients treated with candesartan cilexetil and hydrochlorothiazide. Changes in renal function including acute renal failure can be caused by drugs that inhibit the renin-angiotensin system and by diuretics. Patients whose renal function may depend in part on the activity of the renin-angiotensin system (e.g., patients with renal artery stenosis, chronic kidney disease, severe heart failure, or volume depletion) may be at particular risk of developing oliguria, progressive azotemia, or acute renal failure on candesartan cilexetil and hydrochlorothiazide. Consider withholding or discontinuing therapy in patients who develop a clinically significant decrease in renal function on candesartan cilexetil and hydrochlorothiazide.

## **Potassium Abnormalities**

Drugs that inhibit the renin-angiotensin system can cause hyperkalemia.

Hydrochlorothiazide can cause hypokalemia and hyponatremia. Hypomagnesemia can result in hypokalemia which appears difficult to treat despite potassium repletion.

Monitor serum electrolytes periodically.

In clinical trials of various doses of candesartan cilexetil and hydrochlorothiazide, the incidence of hypertensive patients who developed hypokalemia (serum potassium <3.5 mEq/L) was 2.5% versus 2.1% for placebo; the incidence of hyperkalemia (serum potassium >5.7 mEq/L) was 0.4% versus 1.0% for placebo. No patient receiving candesartan cilexetil and hydrochlorothiazide 16 mg/12.5 mg or 32 mg/12.5 mg was discontinued due to increases or decreases in serum potassium.

## **Acute Myopia and Secondary Angle-Closure Glaucoma**

Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss. The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

## **Hypersensitivity Reaction**

Hypersensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma, but are more likely in patients with such a history.

## **PRECAUTIONS**

### **Metabolic Disturbances**

Hydrochlorothiazide may alter glucose tolerance and raise serum levels of cholesterol and triglycerides.

Hydrochlorothiazide may raise the serum uric acid level due to reduced clearance of uric acid and may cause or exacerbate hyperuricemia and precipitate gout in susceptible patients.

Thiazides decrease urinary calcium excretion and may cause elevation of serum calcium. Avoid using candesartan cilexetil and hydrochlorothiazide tablet in patients with hypercalcemia.

## **Systemic Lupus Erythematosus**

Thiazide diuretics have been reported to cause exacerbation or activation of systemic lupus erythematosus.

### **INFORMATION FOR PATIENTS**

#### *Pregnancy*

Female patients of childbearing age should be told about the consequences of exposure to candesartan cilexetil and hydrochlorothiazide tablet during- pregnancy. Discuss treatment options with women planning to become pregnant. Patients should be asked to report pregnancies to their physicians as soon as possible.

#### *Symptomatic Hypotension*

Tell patients receiving candesartan cilexetil and hydrochlorothiazide tablet that lightheadedness can occur, especially during the first days of therapy, and that it should be reported to the prescribing physician. Tell patients that if syncope occurs, discontinue candesartan cilexetil and hydrochlorothiazide tablet until the physician has been consulted.

Tell all patients that inadequate fluid intake, excessive perspiration, diarrhea, or vomiting can lead to an excessive fall in blood pressure, with the same consequences of lightheadedness and possible syncope.

#### *Hyperkalemia*

Tell patients receiving candesartan cilexetil and hydrochlorothiazide tablet not to use potassium supplements, salt substitutes containing potassium, or other drugs that may increase serum potassium levels without consulting the prescribing physician.

#### *Non-melanoma Skin Cancer*

Instruct patients taking hydrochlorothiazide to protect skin from the sun and undergo regular skin cancer screening.

### **DRUG INTERACTIONS**

Because candesartan is not significantly metabolized by the cytochrome P450 system and at therapeutic concentrations has no effects on P450 enzymes, interactions with drugs that inhibit or are metabolized by those enzymes would not be expected.

#### ***Interactions common to both Candesartan Cilexetil and Hydrochlorothiazide***

*Non-Steroidal Anti-Inflammatory Agents including Selective Cyclooxygenase-2 Inhibitors (COX-2 Inhibitors)* In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, co-administration of NSAIDs, including selective COX-2 inhibitors, with angiotensin II receptor antagonists, including candesartan, may result in deterioration of renal function, including possible acute renal

failure. These effects are usually reversible. Monitor renal function periodically in patients receiving candesartan and NSAID therapy.

The antihypertensive effect of angiotensin II receptor antagonists, including candesartan may be attenuated by NSAIDs including selective COX-2 inhibitors.

*Lithium* – Increases in serum lithium concentrations and lithium toxicity have been reported during concomitant administration of lithium with angiotensin II receptor antagonists or hydrochlorothiazide. Monitor serum lithium levels during concomitant use.

### ***Interactions with Candesartan Cilexetil***

#### **Dual Blockade of the Renin-Angiotensin System (RAS)**

Dual blockade of the RAS with angiotensin receptor blockers, ACE inhibitors, or aliskiren is associated with increased risks of hypotension, hyperkalemia, and changes in renal function (including acute renal failure) compared to monotherapy. Closely monitor blood pressure, renal function and electrolytes in patients on candesartan cilexetil and hydrochlorothiazide and other agents that affect the RAS.

Coadministration of candesartan cilexetil and hydrochlorothiazide with potassium sparing diuretics, potassium supplements, potassium-containing salt substitutes or other drugs that raise serum potassium levels may result in hyperkalemia. Monitor serum potassium in such patients.

Do not co-administer aliskiren with candesartan cilexetil and hydrochlorothiazide in patients with diabetes. Avoid use of aliskiren with candesartan cilexetil and hydrochlorothiazide in patients with renal impairment (GFR <60 ml/min) (see CONTRAINDICATIONS).

### ***Interactions with Hydrochlorothiazide***

Alcohol, barbiturates, or narcotics - Potentiation of orthostatic hypotension may occur.

Antidiabetic drugs (oral agents and insulin) - Dosage adjustment of the antidiabetic drug may be required.

Diazoxide - the hyperglycemic effect of diazoxide may be enhanced by thiazides.

Ion Exchange resins - Single doses of either cholestyramine or colestipol resins bind the hydrochlorothiazide and reduce its absorption from the gastrointestinal tract by up to 85% and 43%, respectively. Stagger the dosage of hydrochlorothiazide and ion exchange resins such that hydrochlorothiazide is administered at least 4 hours before or 4-6 hours after the administration of resins.

Skeletal muscle relaxants, nondepolarizing (e.g., tubocurarine) - Possible increased responsiveness to muscle relaxants such as curare derivatives.

Digitalis - Thiazide-induced hypokalemia or hypomagnesemia may predispose to digoxin toxicity.

Noradrenaline - Thiazides may decrease arterial responsiveness to noradrenaline, but not enough to preclude effectiveness of the pressor agent for therapeutic use.

Steroids or Adrenocorticotrophic Hormone - Hypokalemia may develop during concomitant use of steroids or adrenocorticotrophic hormone (ACTH).

Cytotoxic products - Thiazides may reduce the renal excretion of cytotoxic medicinal products (e.g. cyclophosphamide, methotrexate) and potentiate their myelosuppressive effects.

Cyclosporine - Concomitant treatment with cyclosporine may increase the risk of hyperuricemia and gout-type complications.

## **CARCINOGENESIS & MUTAGENESIS & IMPAIRMENT OF FERTILITY**

No carcinogenicity studies have been conducted with the combination of candesartan cilexetil and hydrochlorothiazide. There was no evidence of carcinogenicity when candesartan cilexetil was orally administered to mice and rats for up to 104 weeks at doses up to 100 and 1000 mg/kg/day, respectively.

Rats received the drug by gavage whereas mice received the drug by dietary administration. These (maximally-tolerated) doses of candesartan cilexetil provided systemic exposures to candesartan AUCs) that were, in mice, approximately 7 times and, in rats, more than 70 times the exposure in man at the maximum recommended daily human dose (32 mg). Two-year feeding studies in mice and rats conducted under the auspices of the National Toxicology Program (NTP) uncovered no evidence of a carcinogenic potential of hydrochlorothiazide in female mice (at doses of up to approximately 600 mg/kg/day) or in male and female rats (at doses of up to approximately 100 mg/kg/day). The NTP, however, found equivocal evidence for hepatocarcinogenicity in male mice.

Candesartan cilexetil or candesartan (the active metabolite), in combination with hydrochlorothiazide, tested positive *in vitro* in the Chinese hamster lung (CHL) chromosomal aberration assay and mouse lymphoma mutagenicity assay. The candesartan cilexetil/hydrochlorothiazide combination tested negative for mutagenicity in bacteria (Ames test), for unscheduled DNA synthesis in rat liver, for chromosomal aberrations in rat bone marrow and for micronuclei in mouse bone marrow.

Both candesartan and its O-deethyl metabolite tested positive for genotoxicity in the *in vitro* CHL chromosomal aberration assay. Neither compound tested positive in the Ames microbial mutagenesis assay or in the *in vitro* mouse lymphoma cell assay. Candesartan (but not its O-deethyl metabolite) was also evaluated *in vivo* in the mouse micronucleus test and *in vitro* in the Chinese hamster ovary (CHO) gene mutation assay, in both cases with negative results. Candesartan cilexetil was evaluated in the Ames test, the *in vitro* mouse lymphoma cell assay, the *in vivo* rat hepatocyte unscheduled DNA synthesis assay and the *in vivo* mouse micronucleus test, in each case with negative results. Candesartan cilexetil was not evaluated in the CHL chromosomal aberration or CHO gene mutation assays.

When hydrochlorothiazide was tested alone, positive results were obtained *in vitro* in the CHO sister chromatid exchange (clastogenicity) and mouse lymphoma cell (mutagenicity) assays and in the *Aspergillus nidulans* non-disjunction assay.

Hydrochlorothiazide was not genotoxic *in vitro* in the Ames test for point mutations and the CHO test for chromosomal aberrations, or *in vivo* in assays using mouse germinal cell chromosomes, Chinese hamster bone marrow chromosomes, and the *Drosophila* sex-linked recessive lethal trait gene.

No fertility studies have been conducted with the combination of candesartan cilexetil and hydrochlorothiazide. Fertility and reproductive performance were not affected in studies with male and female rats given oral doses of up to 300 mg candesartan cilexetil/kg/day (83 times the maximum daily human dose of 32 mg on a body surface area basis). Hydrochlorothiazide had no adverse effects on the fertility of mice and rats of either sex in studies wherein these species were exposed, via their diet, to doses of up to 100 and 4 mg/kg, respectively, prior to conception and throughout gestation.

## **NURSING MOTHERS**

It is not known whether candesartan is excreted in human milk, but candesartan has been shown to be present in rat milk. Thiazides appear in human milk. Because of the potential for adverse effects on the nursing infant, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

## **PEDIATRIC USE**

Neonates with a history of in utero exposure to candesartan cilexetil and hydrochlorothiazide:

If oliguria or hypotension occurs, direct attention toward support of blood pressure and renal perfusion. Exchange transfusions or dialysis may be required as a means of reversing hypotension and/or substituting for disordered renal function.

Safety and effectiveness in pediatric patients have not been established.

## **ADVERSE REACTIONS**

### *Candesartan Cilexetil-Hydrochlorothiazide*

Candesartan cilexetil and hydrochlorothiazide tablets has been evaluated for safety in more than 2800 patients treated for hypertension. More than 750 of these patients were studied for at least six months and more than 500 patients were treated for at least one year. Adverse experiences have generally been mild and transient in nature and have only infrequently required discontinuation of therapy. The overall incidence of adverse events reported with candesartan cilexetil and hydrochlorothiazide tablets was comparable to placebo. The overall frequency of adverse experiences was not related to dose, age, gender, or race.

In placebo-controlled trials that included 1089 patients treated with various combinations of candesartan cilexetil (doses of 2 mg to 32 mg) and hydrochlorothiazide (doses of 6.25 mg -25 mg) and 592 patients treated with placebo, adverse events, whether or not attributed to treatment, occurring in greater than 2% of patients treated with candesartan cilexetil and hydrochlorothiazide tablets and that were more frequent for candesartan cilexetil and hydrochlorothiazide tablets than placebo were:

**Respiratory System Disorder:** upper respiratory tract infection (3.6% vs 3.0%);

**Body as a Whole:** back pain (3.3% vs 2.4%); influenza-like symptoms (2.5% vs 1.9%);  
**Central/Peripheral Nervous System:**

dizziness (2.9% vs 1.2%).

### **Post-Marketing Experience**

The following have been very rarely reported in post-marketing experience with candesartan cilexetil:

**Digestive:** Abnormal hepatic function and hepatitis.

**Hematologic:** Neutropenia, leukopenia, and agranulocytosis.

**Immunologic:** Angioedema

**Metabolic and Nutritional Disorders:** Hyperkalemia, hyponatremia.

**Respiratory System Disorders:** Cough

**Skin and Appendages Disorders:** Pruritus, rash and urticaria.

Rare reports of rhabdomyolysis have been reported in patients receiving angiotensin II receptor blockers.

### **Hydrochlorothiazide**

Other adverse experiences that have been reported with hydrochlorothiazide, without regard to causality, are listed below:

**Gastrointestinal:** pancreatitis, jaundice (intrahepatic cholestatic jaundice), sialadenitis, cramping, constipation, gastric irritation, anorexia

**Hematologic:** aplastic anemia, agranulocytosis, leukopenia, hemolytic anemia, thrombocytopenia

**Hypersensitivity:** anaphylactic reactions, necrotizing angiitis (vasculitis and cutaneous vasculitis), respiratory distress including pneumonitis and pulmonary edema, photosensitivity, urticaria, purpura

**Musculoskeletal:** muscle spasm

**Non-melanoma Skin Cancer:** Hydrochlorothiazide is associated with an increased risk of nonmelanoma skin cancer. In a study conducted in the Sentinel System,

increased risk was predominantly for squamous cell carcinoma (SCC) and in white patients taking large cumulative doses. The increased risk for SCC in the overall population was approximately 1 additional case per 16,000 patients per year, and for white patients taking a cumulative dose of  $\geq 50,000$  mg the risk increase was approximately 1 additional SCC case for every 6,700 patients per year.

**Skin:** erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis, alopecia

**Special Senses:** transient blurred vision, xanthopsia

**Urogenital:** impotence.

## OVERDOSAGE

### *Candesartan Cilexetil – Hydrochlorothiazide*

No lethality was observed in acute toxicity studies in mice, rats and dogs given single oral doses of up to 2000 mg/kg of candesartan cilexetil or in rats given single oral doses of up to 2000 mg/kg of candesartan cilexetil in combination with 1000 mg/kg of hydrochlorothiazide. In mice given single oral doses of the primary metabolite, candesartan, the minimum lethal dose was greater than 1000 mg/kg but less than 2000 mg/kg.

Limited data are available in regard to overdosage with candesartan cilexetil in humans. The most likely manifestations of overdosage with candesartan cilexetil would be hypotension, dizziness, and tachycardia; bradycardia could occur from parasympathetic (vagal) stimulation. If symptomatic hypotension should occur, supportive treatment should be initiated. For hydrochlorothiazide, the most common signs and symptoms observed are those caused by electrolyte depletion (hypokalemia, hypochloremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

Candesartan cannot be removed by hemodialysis. The degree to which hydrochlorothiazide is removed by hemodialysis has not been established.

### *Treatment*

To obtain up-to-date information about the treatment of overdose, consult your Regional Poison Control Center.

Telephone numbers of certified poison control centers are listed in the *Physician's Desk Reference (PDR)*. In managing overdose, consider the possibilities of multiple-drug overdoses, drug-drug interactions, and altered pharmacokinetics in your patient.

## DOSAGE & ADMINISTRATION

The usual recommended starting dose of candesartan cilexetil is 16 mg once daily when it is used as monotherapy in patients who are not volume depleted. Candesartan cilexetil can be administered once or twice daily with total daily doses ranging from 8 mg to 32 mg. Patients requiring further reduction in blood pressure should be titrated to 32 mg. Doses larger than 32 mg do not appear to have a greater blood pressure lowering

effect.

Hydrochlorothiazide is effective in doses of 12.5 to 50 mg once daily.

Use in Renal Impairment: Dosing recommendations for candesartan cilexetil and hydrochlorothiazide in patients with creatinine clearance < 30 mg/min cannot be provided (see SPECIAL POPULATIONS, Renal Insufficiency).

Use in moderate to severe Hepatic Impairment: Candesartan cilexetil and hydrochlorothiazide is not recommended for initiation because the appropriate starting dose, 8 mg, cannot be given. (see SPECIAL POPULATIONS, Hepatic Insufficiency).

**Replacement Therapy:** The combination may be substituted for the titrated components.

**Dose Titration by Clinical Effect:** A patient whose blood pressure is not controlled on 25 mg of hydrochlorothiazide once daily can expect an incremental effect from candesartan cilexetil and hydrochlorothiazide tablets 16 mg/12.5 mg. A patient whose blood pressure is controlled on 25 mg of hydrochlorothiazide but is experiencing decreases in serum potassium can expect the same or incremental blood pressure effects from candesartan cilexetil and hydrochlorothiazide tablets 16 mg/12.5 mg and serum potassium may improve.

A patient whose blood pressure is not controlled on 32 mg of candesartan cilexetil can expect incremental blood pressure effects from candesartan cilexetil and hydrochlorothiazide tablets 32 mg/12.5 mg and then 32 mg/25 mg. The maximal antihypertensive effect of any dose of candesartan cilexetil and hydrochlorothiazide tablets can be expected within 4 weeks of initiating that dose.

Candesartan cilexetil and hydrochlorothiazide tablets may be administered with other antihypertensive agents.

Candesartan cilexetil and hydrochlorothiazide tablets may be administered with or without food.

## **HOW SUPPLIED**

**Candesartan cilexetil and hydrochlorothiazide tablets 16 mg/12.5 mg**, are Pink, oval, biconvex, uncoated tablets debossed with "ML 71" on one side and break line on both sides They are supplied as follows:

**NDC** 33342-131-12 unit dose package of 100

**NDC** 33342-131-10 bottle of 90

**NDC** 33342-131-51 bottle of 300

**NDC** 33342-131-15 bottle of 500

**Candesartan cilexetil and hydrochlorothiazide tablets 32 mg/12.5 mg**, are Yellow oval, biconvex, uncoated tablets debossed with "ML 58" on one side and break line on both sides. They are supplied as follows:



**NDC 33342-132-12** unit dose package of 100

**NDC 33342-132-10** bottle of 90

**NDC 33342-132-51** bottle of 300

**NDC 33342-132-15** bottle of 500

**Candesartan cilexetil and hydrochlorothiazide tablets 32 mg/25 mg**, are Pink, oval, biconvex, uncoated tablets debossed with "ML 57" on one side and break line on both sides. They are supplied as follows:

**NDC 33342-133-12** unit dose package of 100

**NDC 33342-133-10** bottle of 90

**NDC 33342-133-51** bottle of 300

**NDC 33342-133-15** bottle of 500

**Storage:**

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15 to 30°C (59 to 86°F) [see USP Controlled Room Temperature]. Keep container tightly closed.

Manufactured for :

**Macleods Pharma USA, Inc.**

Princeton, NJ 08540

Manufactured by :

Macleods Pharmaceuticals Ltd.

Daman (U.T.), INDIA

**Rev: 10/2022**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**PRINCIPAL DISPLAY PANEL**




NDC 33342-131-10

**Candesartan Cilexetil and Hydrochlorothiazide Tablets**

16 mg/12.5 mg Bottle Label

Rx only

90 Tablets

	 NDC 33342-131-10	Each tablet contains: Candesartan Cilexetil, USP 16 mg Hydrochlorothiazide, USP 12.5 mg	
	<b>Candesartan Cilexetil and Hydrochlorothiazide Tablets</b> <b>16 mg/12.5 mg</b>	<b>USUAL DOSAGE :</b> See prescribing information for Dosage and administration.  Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].  Keep container tightly closed.	
Rx Only	90 Tablets	Manufactured for : <b>Macleods Pharma USA, Inc.</b> Princeton, NJ 08540	Manufactured by: Macleods Pharmaceuticals Ltd. Daman (U.T.), INDIA
		Code No.: DD/DRUGS/DD/375 PM01845503	

## PRINCIPAL DISPLAY PANEL




NDC 33342-131-51

### Candesartan Cilexetil and Hydrochlorothiazide Tablets

16 mg/12.5 mg Bottle Label

Rx only

300 Tablets

	 NDC 33342-131-51	Each tablet contains: Candesartan Cilexetil, USP 16 mg Hydrochlorothiazide, USP 12.5 mg	
	<b>Candesartan Cilexetil and Hydrochlorothiazide Tablets</b> <b>16 mg/12.5 mg</b>	<b>USUAL DOSAGE :</b> See prescribing information for Dosage and administration.  Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].  Keep container tightly closed.	
Rx Only	300 Tablets	Manufactured for : <b>Macleods Pharma USA, Inc.</b> Princeton, NJ 08540	Manufactured by: Macleods Pharmaceuticals Ltd. Daman (U.T.), INDIA
		Code No.: DD/DRUGS/DD/375 PM01845603	

## PRINCIPAL DISPLAY PANEL


NDC 33342-131-15


### Candesartan Cilexetil and Hydrochlorothiazide Tablets

16 mg/12.5 mg Bottle Label

Rx only

500 Tablets





NDC 33342-131-15

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**


**16 mg/12.5 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 16 mg  
Hydrochlorothiazide, USP 12.5 mg

**USUAL DOSAGE :**  
See prescribing information for Dosage and administration.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Keep container tightly closed.



OPZ AREA  
45 X 24 mm

R<sub>x</sub> Only      500 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375  
PM01901603

## PRINCIPAL DISPLAY PANEL


NDC 33342-132-10


### Candesartan Cilexetil and Hydrochlorothiazide Tablets

32 mg/12.5 mg Bottle Label

Rx only

90 Tablets





NDC 33342-132-10

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**


**32 mg/12.5 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 32 mg  
Hydrochlorothiazide, USP 12.5 mg

**USUAL DOSAGE :**  
See prescribing information for Dosage and administration.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Keep container tightly closed.



OPZ AREA  
43 X 15 mm

R<sub>x</sub> Only      90 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375  
PM01845703


## PRINCIPAL DISPLAY PANEL

NDC 33342-132-51


### Candesartan Cilexetil and Hydrochlorothiazide Tablets

32 mg/12.5 mg Bottle Label

Rx only  
300 Tablets



3 33342132512



NDC 33342-132-51

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**


**32 mg/12.5 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 32 mg  
Hydrochlorothiazide, USP 12.5 mg

USUAL DOSAGE :  
See prescribing  
information for Dosage  
and administration.

Store at 20° to 25°C  
(68° to 77°F); excursions  
permitted to 15° to 30°C  
(59° to 86°F) [See USP  
Controlled Room Temperature].

Keep container tightly closed.



←

OPZ AREA  
43 X 24 mm

R<sub>x</sub> Only

300 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375  
PM01845803

## PRINCIPAL DISPLAY PANEL


NDC 33342-132-15

### Candesartan Cilexetil and Hydrochlorothiazide Tablets


32 mg/12.5 mg Bottle Label

Rx only

500 Tablets



3 33342132154



NDC 33342-132-15

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**


**32 mg/12.5 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 32 mg  
Hydrochlorothiazide, USP 12.5 mg

USUAL DOSAGE :  
See prescribing information for  
Dosage and administration.

Store at 20° to 25°C  
(68° to 77°F); excursions  
permitted to 15° to 30°C  
(59° to 86°F) [See USP  
Controlled Room Temperature].

Keep container tightly closed.



←

OPZ AREA  
43 X 25 mm

R<sub>x</sub> Only

500 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375  
PM01901703

## PRINCIPAL DISPLAY PANEL




NDC 33342-133-10

### Candesartan Cilexetil and Hydrochlorothiazide Tablets

32 mg/25 mg Bottle Label

Rx only

90 Tablets

 33342133106	 NDC 33342-133-10	Each tablet contains: Candesartan Cilexetil, USP 32 mg Hydrochlorothiazide, USP 25 mg	 ←
	<b>Candesartan Cilexetil and Hydrochlorothiazide Tablets</b> <b>32 mg/25 mg</b>	USUAL DOSAGE : See prescribing information for Dosage and administration.  Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].  Keep container tightly closed.	
R <sub>x</sub> Only	90 Tablets	Manufactured for : <b>Macleods Pharma USA, Inc.</b> Princeton, NJ 08540	Manufactured by: Macleods Pharmaceuticals Ltd. Daman (U.T.), INDIA
		Code No.: DD/DRUGS/DD/375 PM01845903	

## PRINCIPAL DISPLAY PANEL

NDC 33342-133-51

### Candesartan Cilexetil and Hydrochlorothiazide Tablets

32 mg/25 mg Bottle Label

Rx only

300 Tablets







NDC 33342-133-51

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**

**32 mg/25 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 32 mg  
Hydrochlorothiazide, USP 25 mg

USUAL DOSAGE :  
See prescribing  
information for Dosage  
and administration.

Store at 20° to 25°C  
(68° to 77°F); excursions  
permitted to 15° to 30°C  
(59° to 86°F) [See USP  
Controlled Room Temperature].

Keep container tightly closed.



OPZ AREA  
43 X 25 mm

R<sub>x</sub> Only

300 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375

PM01846003

## PRINCIPAL DISPLAY PANEL


NDC 33342-133-15


### Candesartan Cilexetil and Hydrochlorothiazide Tablets

32 mg/25 mg Bottle Label

Rx only

500 Tablets





NDC 33342-133-15

**Candesartan  
Cilexetil and  
Hydrochlorothiazide  
Tablets**

**32 mg/25 mg**

Each tablet contains:  
Candesartan Cilexetil, USP 32 mg  
Hydrochlorothiazide, USP 25 mg

USUAL DOSAGE :  
See prescribing information for  
Dosage and administration.

Store at 20° to 25°C  
(68° to 77°F); excursions  
permitted to 15° to 30°C  
(59° to 86°F) [See USP  
Controlled Room Temperature].

Keep container tightly closed.



OPZ AREA  
40 X 25 mm

R<sub>x</sub> Only

500 Tablets

Manufactured for :  
**Macleods Pharma USA, Inc.**  
Princeton, NJ 08540

Manufactured by:  
Macleods Pharmaceuticals Ltd.  
Daman (U.T.), INDIA

Code No.: DD/DRUGS/DD/375

PM01901803

CANDESARTAN CILEXETIL AND HYDROCHLOROTHIAZIDE

candesartan cilexetil and hydrochlorothiazide tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33342-131
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CANDESARTAN CILEXETIL (UNII: R85M2X0D68) (CANDESARTAN - UNII:S8Q36MD2XX)	CANDESARTAN CILEXETIL	16 mg
HYDROCHLOROTHIAZIDE (UNII: 0J48LPH2TH) (HYDROCHLOROTHIAZIDE - UNII:0J48LPH2TH)	HYDROCHLOROTHIAZIDE	12.5 mg

Inactive Ingredients	
Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE CALCIUM (UNII: UTY7PDF93L)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics			
Color	PINK	Score	2 pieces
Shape	OVAL (biconvex)	Size	10mm
Flavor		Imprint Code	ML71
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-131-12	10 in 1 CARTON	03/07/2015	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:33342-	90 in 1 BOTTLE; Type 0: Not a Combination	03/07/2015	

2	131-10	Product	03/07/2015	
3	NDC:33342-131-51	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
4	NDC:33342-131-15	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204100	03/07/2015	

CANDESARTAN CILEXETIL AND HYDROCHLOROTHIAZIDE

candesartan cilexetil and hydrochlorothiazide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33342-132
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CANDESARTAN CILEXETIL (UNII: R85M2X0D68) (CANDESARTAN - UNII:S8Q36MD2XX)	CANDESARTAN CILEXETIL	32 mg
HYDROCHLOROTHIAZIDE (UNII: 0J48LPH2TH) (HYDROCHLOROTHIAZIDE - UNII:0J48LPH2TH)	HYDROCHLOROTHIAZIDE	12.5 mg

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE CALCIUM (UNII: UTY7PDF93L)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	YELLOW	Score	2 pieces
Shape	OVAL (biconvex)	Size	11mm
Flavor		Imprint Code	ML58
Contains			

Packaging



#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-132-12	10 in 1 CARTON	03/07/2015	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:33342-132-10	90 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
3	NDC:33342-132-51	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
4	NDC:33342-132-15	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204100	03/07/2015	

## CANDESARTAN CILEXETIL AND HYDROCHLOROTHIAZIDE

candesartan cilexetil and hydrochlorothiazide tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:33342-133
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CANDESARTAN CILEXETIL</b> (UNII: R85M2X0D68) (CANDESARTAN - UNII:S8Q36MD2XX)	CANDESARTAN CILEXETIL	32 mg
<b>HYDROCHLOROTHIAZIDE</b> (UNII: 0J48LPH2TH) (HYDROCHLOROTHIAZIDE - UNII:0J48LPH2TH)	HYDROCHLOROTHIAZIDE	25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE CALCIUM</b> (UNII: UTY7PDF93L)	
<b>HYDROXYPROPYL CELLULOSE</b> (UNII: RFW2ET671P)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	

### Product Characteristics

Color	PINK	Score	2 pieces
-------	------	-------	----------

Shape	OVAL (biconvex)	Size	11mm	
Flavor		Imprint Code	ML57	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33342-133-12	10 in 1 CARTON	03/07/2015	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:33342-133-10	90 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
3	NDC:33342-133-51	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
4	NDC:33342-133-15	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2015	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA204100		03/07/2015	

**Labeler -** Macleods Pharmaceuticals Limited (862128535)

## Establishment

Name	Address	ID/FEI	Business Operations
Macleods Pharmaceuticals Limited		918608365	ANALYSIS(33342-131, 33342-132, 33342-133) , LABEL(33342-131, 33342-132, 33342-133) , MANUFACTURE(33342-131, 33342-132, 33342-133) , PACK(33342-131, 33342-132, 33342-133)

Revised: 10/2022

Macleods Pharmaceuticals Limited