

**LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated**  
**Walgreens Company**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Levocetirizine dihydrochloride USP, 5 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

**Ask a doctor before use if you have**

- ever had trouble urinating or emptying your bladder

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask doctor if**

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

adults 65 years of age and older	<ul style="list-style-type: none"><li>ask a doctor</li></ul>
adults and children 12-64 years of age	<ul style="list-style-type: none"><li>take 1 tablet (5 mg) once daily in the evening</li><li>do not take more than 1 tablet (5 mg) in 24 hours</li><li>1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms</li></ul>
children 6-11 years of age	<ul style="list-style-type: none"><li>take 1/2 tablet (2.5 mg) once daily in the evening</li><li>do not take more than 1/2 tablet (2.5 mg) in 24 hours</li></ul>
children under 6 years of age	<ul style="list-style-type: none"><li>do not use</li></ul>
consumers with kidney disease	<ul style="list-style-type: none"><li>do not use</li></ul>

## Other information

- store between 20° and 25°C (68° and 77°F)
- (bottles only) safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- (blister only) safety sealed: do not use if seal is broken or if individual blister unit is open or torn

## Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

## Questions or comments?

Call 1-888-375-3784

## Carton Label

**ORIGINAL PRESCRIPTION STRENGTH**

**Compare to Xyzal® Allergy 24HR active ingredient<sup>†‡</sup>**

0363-5529-35

NEW Well at Walgreens

WALGREENS PHARMACIST RECOMMENDED<sup>‡</sup>

ORIGINAL  
PRESCRIPTION  
STRENGTH

**Levocetirizine  
24-Hour Allergy**

**Levocetirizine Dihydrochloride  
Tablets, 5 mg / Antihistamine**

Relief of:

- Sneezing
- Runny nose
- Itchy nose or throat
- Itchy, watery eyes

24 HOUR

35 TABLETS



**Bottle Label**

Well at Walgreens

NDC 0363-5529-35

ORIGINAL  
PRESCRIPTION STRENGTH

**Levocetirizine**  
**24-Hour Allergy**

**Levocetirizine Dihydrochloride**  
**Tablets, 5 mg / Antihistamine**

35 TABLETS



NDC 0363-5529-35

ORIGINAL PRESCRIPTION STRENGTH

# Levocetirizine

24-Hour Allergy

Levocetirizine Dihydrochloride Tablets, 5 mg / Antihistamine

35 TABLETS

### Active ingredient (in each tablet)

Levocetirizine dihydrochloride USP, 5 mg.....Antihistamine

### Purpose

**Uses** temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat **Warnings Do not use** ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine **Ask a doctor before use if you have** ■ ever had trouble urinating or emptying your bladder **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery **Stop use and ask a doctor if** ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use

DISTRIBUTED BY: WALGREEN CO.  
200 WILMUT RD, DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEE W00000-0000-0  
walgreens.com ©2017 Walgreen Co. ORG0118-F-R1  
REV-0318

LOT / EXP 150072353

Peel Here

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Directions** **adults 65 years of age and older** ■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms **children 6-11 years of age** ■ take 1/2 tablet (2.5 mg) once daily in the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers with kidney disease** ■ do not use **Other Information** ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing **Inactive ingredients** colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide **Questions or comments?** call 1-888-375-3784

## LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-5529(NDC:43598-735)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
levocetirizine dihydrochloride (UNII: SOD6A38AGA) (levo cetirizine - UNII:6U5EA9RT2O)	levo cetirizine dihydrochloride	5 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5529-35	1 in 1 CARTON	03/26/2018	
1		35 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-5529-55	1 in 1 CARTON	03/26/2018	
2		55 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-5529-10	2 in 1 CARTON	04/17/2018	
3		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:0363-5529-80	1 in 1 CARTON	03/26/2018	
4		80 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0363-5529-12	1 in 1 CARTON	01/15/2021	
5		120 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210375	03/26/2018	

**Labeler** - Walgreens Company (008965063)

Revised: 7/2020

Walgreens Company