

**NAPROXEN SODIUM- naproxen sodium tablet  
FAMILY DOLLAR**

-----

**ALL DAY**

**PAIN RELIEF**

*Naproxen Sodium Tablets USP, 220 mg (NSAID)\*\**

*Pain Reliever/Fever Reducer*

*Strength to Last 12 Hours*

**Active ingredient (For Tablet)**

*(in each tablet)*

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

**Active ingredient (For Caplet)**

*(in each caplet)*

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Allergy alert:**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives

- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of the reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions (For Tablets)**

- **do not take more than directed**
  - **the smallest effective dose should be used**
  - drink a full glass of water with each dose
- Adults and children 12 years and older
- take 1 tablet every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 tablets within the first hour
  - do not exceed 2 tablets in any 8- to 12- hour period
  - do not exceed 3 tablets in a 24- hour period

Children under 12 years

- ask a doctor

### **Directions (For Caplets)**

- **do not take more than directed**
- **the smallest effective dose should be used**

- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12- hour period
- do not exceed 3 caplets in a 24- hour period

Children under 12 years

- ask a doctor

### **Other information (For Tablet)**

- **each tablet contains:** sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

### **Other information (For Caplet)**

- **each caplet contains:** sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

### **Inactive ingredients**

FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide.

### **Questions or comments?**

1-877-770-3183 Mon - Fri 9:00 AM to 4:00 PM EST.

### **Naproxen Sodium 220 mg Tablets and Caplets**



size 11.785 X 1.797 X 3.7  
 part # 191497180 vlen 191497180  
 material 1.016 889 date 07/18/2018

Distributed by:  
**LLC**  
 5700 Vinton Parkway  
 Chesapeake, VA 23060  
**NOT FOR SALE**  
 Return package and unused product  
 within 30 days to any Family Wellness  
 for a refund (with receipt) or exchange.  
 Contains no synthetic waste from a  
 plastic-containing gram (weight), jar, or  
 or (g)

R-01  
 300000365  
 532000002

**Drug Facts (continued)**  
**Heart attack and stroke warnings:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use**  
 if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 if right before or after heart surgery  
**Ask a doctor before use if**  
 the stomach bleeding warning applies to you  
 you have a history of stomach problems, such as heartburn  
 you have had high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke  
 you are taking a diuretic  
 you have problems or serious side effects from taking pain relievers or fever reducers  
**Ask a doctor or pharmacist before use if you are**  
 under a doctor's care for any serious condition  
 taking any other drug  
**When using this product**  
 take with food or milk, if stomach upset occurs  
**Stop use and ask a doctor if**  
 you experience any of the following signs of stomach bleeding:  
 red, black, or bloody stools  
 vomit blood  
 have bloody or black stools  
 have stomach pain that does not get better  
 you have symptoms of heart problems or stroke:  
 chest pain trouble breathing  
 weakness in one part or side of body  
 slurred speech leg swelling  
 pain gets worse or lasts more than 10 days  
 fever gets worse or lasts more than 3 days  
 you have difficulty swallowing  
 it feels like the pill is stuck in your throat  
 redness or swelling is present in the painful area →

**Drug Facts (continued)**  
**if any new symptoms appear**  
 if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
 do not take more than directed  
 the smallest amount possible should be used  
 drink a full glass of water with each dose  
**Adults and children 12 years and older**  
 take 1 caplet every 8 to 12 hours while symptoms last  
 for the first dose you may take 2 caplets within the first hour  
 do not exceed 2 caplets in any 8- to 12-hour period  
 do not exceed 3 caplets in a 24-hour period  
 ask a doctor  
**Children under 12 years**  
 ask a doctor  
**Other information**  
 each caplet contains: sodium 20 mg  
 is active at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)  
**Inactive ingredients:** FD&C blue #2 aluminum lake, hypromellose 5910, microcrystalline cellulose, polyethylene glycol, polyvinyl K-30, sodium starch glycolate, stearic acid, titanium dioxide.  
**Questions or comments?**  
 1-877-776-8183 Mon-Fri 8:00 AM to 6:00 PM EST.  
 MADE IN INDIA

**50 CAPLETS**  
**Strength to Last 12 Hours**  
**ALL DAY PAIN RELIEF**  
**naproxen sodium tablets USP 220 mg (NSAID)\***  
**Pain Reliever/Fever Reducer**  
**FAMILY Wellness™**  
**CONTAINS 50 ALABAMA CAPLETS**  
**OTC**  
 Lot: Exp:

**KEEP CARTON FOR REFERENCE**  
**DO NOT USE IF CARTON IS OPEN OR IF PILL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN.**  
**Drug Facts**  
**Active ingredient (in each caplet)**  
 naproxen sodium 220 mg (NSAID)\*  
**Purposes**  
 Pain reliever/fever reducer  
 (nonsteroid) anti-inflammatory drug  
**Uses**  
 temporarily relieve minor aches and pains due to:  
 minor pain of arthritis muscular aches  
 backache menstrual cramps  
 headache toothache  
 the common cold  
 temporarily reduce fever  
**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 hives facial swelling in softness (swelling)  
 in shock skin redness in eyes in hives  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 are age 60 or older  
 have had stomach ulcers or bleeding problems  
 take a blood thinning (anticoagulant) or steroid drug  
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, rofecoxib, or others)  
 have 3 or more alcoholic drinks every day while using this product  
 take more or for a longer time than directed →

300000365  
 532000002  
 R-01

SKU 0914123  
 0 32251 43069 5  
 \*This product is not manufactured or distributed by Bayer HealthCare, LLC, distributor of Advil. Name is registered trademark of Bayer HealthCare, LLC.

Lot: Exp:

end user : GRANULES PHARMA	
item : 100 CT TABLET	
code # :	
size : 1.77 X 1.77 X 4.17	
ref # : PP180718A	view : PRINT
material : 016 SDS	date : 07/27

R-01  
5320000367  
300000367

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - feel faint
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

300000367  
532000003  
R-01

**Drug Facts (continued)**

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

**Adults and children 12 years and older**

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

**Children under 12 years**

- ask a doctor

**Other information**

- each caplet contains: sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

**Inactive Ingredients** FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone K-30, sodium starch glycolate, stearic acid, titanium dioxide.

**Questions or comments?**  
1-877-770-3103 Mon-Fri 9:00 AM to 4:00 PM EST.

\*This product is not manufactured or distributed by Bayer HealthCare, LLC, manufacturer of Aleve. Aleve is a registered trademark of Bayer HealthCare, LLC.

Distributed By:  
Midwood Brands, LLC  
500 Volvo Parkway  
Chesapeake, VA 23320

**NOT 100% SATISFIED?**  
Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

Contains no ingredient made from a gluten-containing grain (wheat, barley, or rye)

ALL DAY  
**PAIN RELIEF**  
Naproxen Sodium Tablets USP,  
220 mg (NSAID)\*\*  
Pain Reliever/Fever Reducer

FAMILY  
wellness™

ALL DAY  
**PAIN RELIEF**  
Naproxen Sodium Tablets USP, 220 mg (NSAID)\*\*  
Pain Reliever/Fever Reducer

100 CAPLETS  
Strength to Last 12 Hours

FAMILY  
wellness™

COMPARE TO ALEVE® CAPLETS\*

OTC  
OVER THE COUNTER

MADE IN INDIA

SKU 0902330  
0 32251 43067 1  
MADE IN INDIA

Lot  
Exp.

**KEEP CARTON FOR REFERENCE**

**DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL OR BOTTLE OPENING IS MISSING OR BROKEN.**

**Drug Facts**

Active ingredient (in each caplet)	Purposes
Naproxen sodium 220 mg..... Pain reliever/fever reducer (naproxen 200 mg) (NSAID)**	**nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - backache
  - headache
  - the common cold
  - temporarily reduces fever
  - muscular aches
  - menstrual cramps
  - toothache

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- shock
- skin redness
- asthma (wheezing)
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.





code # :  
 size : 1+3/4 X 1+3/4 X 3+1/2  
 ref # : P9180811B view :  
 material : 016 S93 date :

R-00  
 532000020  
 300000476

**Drug Facts (continued)**  
**Heart attack and stroke warnings:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. This risk is higher if you use more than directed or for longer than directed.  
**Do not use**  
 if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 if you have a history of stomach problems, such as heartburn  
 if you have had high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke  
 if you are taking a diuretic  
 if you have problems or serious side effects from taking pain relievers or fever reducers  
**Ask a doctor or pharmacist before use if you are**  
 under a doctor's care for any serious condition  
 or taking any other drug  
**When using this product**  
 take with food or milk if stomach upset occurs  
**Stop use and ask a doctor if**  
 you experience any of the following signs of stomach bleeding:  
 vomit blood  
 have bloody or black stools  
 have stomach pain that does not get better  
 You have symptoms of heart problems or stroke:  
 chest pain  
 weakness in one part or side of body  
 slurred speech  
 leg swelling  
 pain gets worse or lasts more than 10 days  
 fever gets worse or lasts more than 3 days  
 you have difficulty swallowing  
 it feels like the pill is stuck in your throat  
 or redness or swelling is present in the painful area

**Drug Facts (continued)**  
**if any new symptoms appear**  
 if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
 do not take more than directed  
 the smallest effective dose should be used  
 drink a full glass of water with each dose  
**Adults and children 12 years and older**  
 take 1 tablet every 8 to 12 hours while symptoms last  
 for the first dose you may take 2 tablets within the first hour  
 do not exceed 2 tablets in any 8- to 12-hour period  
 do not exceed 3 tablets in a 24-hour period  
**Children under 12 years**  
 ask a doctor  
**Other information**  
 keep sealed container below 20 mg in store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)  
**Inactive ingredients:** FD&C blue #2 aluminum lake, hypromellose 2910, microcrystalline cellulose, polyethylene glycol, croscarmellose sodium starch glycolate, stearic acid, titanium dioxide.  
**Questions or comments?** 1-877-776-3183 Mon-Fri 9:00 AM to 4:00 PM EST.

15 TABLETS Strength to Last 12 Hours  
 PAIN RELIEF  
 Naproxen sodium tablets USP, 220 mg (NSAID)  
 Pain Reliever/Fever Reducer  
 FAMILY WELLNESS  
 COMBINE TO ALLEVIATE PAIN  
 OTC  
 15 TABLETS  
 Lot  
 Exp.

**KEEP CARTON FOR REFERENCE**  
**DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL OR BOTTLE OPENING IS MISSING OR BROKEN.**  
**Drug Facts**  
**Active Ingredient (in each tablet):** Naproxen sodium 220 mg... Pain reliever/fever reducer  
**Purpose(s):** "nonsteroidal anti-inflammatory drug"  
**Uses**  
 temporarily relieve minor aches and pains due to:  
 minor pain of arthritis  
 backache  
 headache  
 in the common cold  
 temporarily reduces fever  
**Warnings**  
 Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 hives  
 facial swelling  
 chest pain  
 trouble breathing  
 dizziness  
 fainting  
 rash  
 blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 are age 60 or older  
 have had stomach ulcers or bleeding problems  
 take a blood thinning (anticoagulant) or steroid drug  
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or other)  
 have 3 or more alcoholic drinks every day while using this product  
 take more or for a longer time than directed

300000476  
 532000020  
 R-00

SKU 000560  
 0 52251 52290 1

# NAPROXEN SODIUM

naproxen sodium tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-761
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:5Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	



STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

### Product Characteristics

Color	blue	Score	no score
Shape	ROUND (Biconvex Film Coated Tablets)	Size	10mm
Flavor		Imprint Code	220
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-761-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	
2	NDC:55319-761-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	
3	NDC:55319-761-07	15 in 1 BOTTLE; Type 0: Not a Combination Product	10/31/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	10/01/2018	

## NAPROXEN SODIUM

naproxen sodium tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-762
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL (Biconvex Film Coated Tablets)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	220
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-762-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	
2	NDC:55319-762-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	
3	NDC:55319-762-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	10/31/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	10/01/2018	

**Labeler** - FAMILY DOLLAR(024472631)

Revised: 1/2021

FAMILY DOLLAR