

# ACETAMINOPHEN- acetaminophen tablet

## Major Pharmaceuticals

-----  
**5427-Major**

### ***Drug Facts***

#### ***Active Ingredient (in each tablet)***

Acetaminophen 500 mg

#### ***Purpose***

Pain reliever/ Fever reducer

### ***Uses***

Temporarily reduces fever and relieves minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening ,blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ☐
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if** ☐

- pain gets worse or lasts more than 10 days☐
- fever gets worse or lasts more than 3 days ☐
- new symptoms occur ☐

- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed (see overdose warning)**

**adults and children 12 years and over:**

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**children under 12 years:** ask a doctor

### ***Other Information***

store in a dry place at 15° – 30°C (59° – 86°F).

Povidone, pregelatinized starch, sodium starch glycolate, stearic acid

### ***Questions or comments?***

**Call 1-800-231-4670**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

**Distributed by:**

**MAJOR® PHARMACEUTICALS  
Indianapolis, IN 46268**

\*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Tylenol®.

**MAJOR**

THIS LARGE SIZE PACKAGE FOR DISPENSING PURPOSE ONLY

See New Warnings Information and Directions

**Acetaminophen Extra Strength Tablets**

Aspirin Free

Extra Strength Pain reliever

Compare to active ingredient in Extra Strength Tylenol®\* Tablets

1000 tablets

500mg each

# MAJOR®

NDC 0904-6730-80

THIS LARGE SIZE PACKAGE FOR DISPENSING PURPOSE ONLY

**EXTRA STRENGTH**

## Acetaminophen

**TABLETS**

**ASPIRIN FREE**  
**EXTRA STRENGTH**  
**PAIN RELIEVER/  
FEVER REDUCER**

Compare to the active ingredient  
in Extra Strength Tylenol® Tablets\*

**1000 TABLETS**  
**500 MG EACH**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)**  
Acetaminophen 500 mg ..... Pain reliever/fever reducer

**Purpose**  
Pain relief/fever reducer

**Uses** Temporarily reduces fever and relieves minor aches and pains due to:  
 ● headache ● muscular aches ● the common cold ● toothache ● backache  
 ● minor pain of arthritis ● premenstrual and menstrual cramps

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ● more than 4,000 mg of acetaminophen in 24 hours ● with other drugs containing acetaminophen ● 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ● skin redness ● blisters ● rash  
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use ● with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ● if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● new symptoms occur ● redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ● do not take more than directed (see overdose warning) adults and children 12 years and over: ● take 2 tablets every 6 hours while symptoms last ● do not take more than 6 tablets in 24 hours, unless directed by a doctor ● do not take for more than 10 days unless directed by a doctor children under 12 years: ask a doctor

**Other information** store in a dry place at 15°–30°C (59°–86°F).

**Inactive ingredients** povidone, pregelatinized starch, sodium starch glycolate, stearic acid

**Questions or comments? Call 1-800-231-4670**

This product is not manufactured or distributed by Kenue Inc., owner of the registered trademark Tylenol®.

Distributed by:  
**MAJOR® PHARMACEUTICALS** call (800) 616-2471  
 www.majorpharmaceuticals.com  
 Indianapolis, IN 46268

Rec-order No. 700923 M-23 Rev. 09/23 R52941

0904673080 4  
3

## ACETAMINOPHEN

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0904-6730
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	54;27

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6730-60	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2018	
2	NDC:0904-6730-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2018	
3	NDC:0904-6730-59	1 in 1 CARTON	09/12/2018	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0904-6730-61	10 in 1 BOX, UNIT-DOSE	09/12/2018	
4		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/2018	

**Labeler** - Major Pharmaceuticals (191427277)

Revised: 5/2024

Major Pharmaceuticals